



US010760075B2

(12) **United States Patent**
Sommer et al.

(10) **Patent No.:** **US 10,760,075 B2**
(45) **Date of Patent:** **Sep. 1, 2020**

(54) **TREATING AND PREVENTING MICROBIAL INFECTIONS**

(71) Applicant: **SNIPR Biome ApS**, Copenhagen Ø (DK)

(72) Inventors: **Morten Sommer**, Copenhagen Ø (DK); **Virginia Martinez**, Copenhagen Ø (DK); **Eric Van Der Helm**, Copenhagen Ø (DK); **Jakob Krause Haaber**, Copenhagen Ø (DK); **Ana De Santiago Torio**, Copenhagen Ø (DK); **Christian Grøndahl**, Copenhagen Ø (DK); **Jasper Clube**, London (GB)

(73) Assignee: **SNIPR Biome ApS**, Copenhagen Ø (DK)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 146 days.

8,329,867 B2 12/2012 Lazar et al.
8,354,509 B2 1/2013 Carven et al.
8,735,553 B1 5/2014 Li et al.
9,701,964 B2 7/2017 Clube et al.
9,758,583 B2* 9/2017 Wang C07K 16/2818
9,822,372 B2 11/2017 Zhang et al.
9,879,269 B2 1/2018 Barrangou et al.
10,066,233 B2 9/2018 Barrangou et al.
10,136,639 B2 11/2018 Wuest et al.
10,136,649 B2 11/2018 Barrangou et al.
10,195,273 B2 2/2019 Clube et al.
10,300,138 B2 5/2019 Clube
10,300,139 B2 5/2019 Clube
10,363,308 B2 7/2019 Clube
10,463,049 B2 11/2019 Clube
10,506,812 B2 12/2019 Clube
10,524,477 B2 1/2020 Clube
10,596,255 B2 3/2020 Clube
10,603,379 B2 3/2020 Clube
2004/0096974 A1 5/2004 Herron et al.

(Continued)

FOREIGN PATENT DOCUMENTS

(21) Appl. No.: **15/967,484**

(22) Filed: **Apr. 30, 2018**

EP 2 320 940 B1 5/2011
EP 2 840 140 A1 2/2015

(Continued)

(65) **Prior Publication Data**

US 2018/0273940 A1 Sep. 27, 2018

OTHER PUBLICATIONS

(51) **Int. Cl.**

C12N 15/11 (2006.01)
A61P 31/04 (2006.01)
A61K 39/395 (2006.01)
A61K 38/00 (2006.01)
C07K 14/21 (2006.01)
A61K 38/46 (2006.01)
C07K 16/28 (2006.01)

Mayo Clinic, <https://www.mayoclinic.org/diseases-conditions/bacterial-vaginosis/symptoms-causes/syc-20352279>, accessed Jul. 20, 2018.*

(Continued)

(52) **U.S. Cl.**

CPC **C12N 15/11** (2013.01); **A61K 38/465** (2013.01); **A61K 39/3955** (2013.01); **A61K 39/39558** (2013.01); **A61P 31/04** (2018.01); **C07K 16/2818** (2013.01); **C07K 16/2827** (2013.01); **C12N 2310/20** (2017.05); **C12N 2320/30** (2013.01); **C12N 2320/31** (2013.01); **Y02A 50/478** (2018.01)

Primary Examiner — Gary B Nickol

Assistant Examiner — Lalkia J Jackson-Tongue

(74) *Attorney, Agent, or Firm* — Morrison & Foerster LLP

(58) **Field of Classification Search**

CPC **C12N 15/111**; **C12N 2320/30**; **C12N 2320/31**; **A61P 31/04**; **A61K 39/3955**
See application file for complete search history.

(57) **ABSTRACT**

The invention provides methods for treating or preventing microbial (eg, bacterial) infections and means for performing these methods. In particular, treatment of infections requiring rapid and durable therapy is made possible, such as for treating acute conditions such as septicemia, sepsis, SIRS or septic shock. The invention is particularly useful, for example, for treatment of microbes such as for environmental, food and beverage use. The invention relates inter alia to methods of controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate or fluid in an industrial or domestic system. The invention also useful for the treatment of pathogenic bacterial infections in subjects receiving a treatment for a disease or condition, such as a transplant or a treatment for cancer, a viral infection or an autoimmune disease.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,626,504 A 12/1986 Puhler et al.
4,870,287 A 9/1989 Cole et al.
5,633,154 A 5/1997 Schaefer et al.
5,760,395 A 6/1998 Johnstone
5,844,905 A 12/1998 McKay et al.
5,885,796 A 3/1999 Linsley et al.
6,207,156 B1 3/2001 Kuchroo et al.
8,008,449 B2 8/2011 Korman et al.
8,017,114 B2 9/2011 Korman et al.
8,119,129 B2 2/2012 Jure-Kunkel et al.
8,241,498 B2 8/2012 Summer et al.
8,252,576 B2 8/2012 Campbell et al.

19 Claims, 9 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

U.S. PATENT DOCUMENTS

2005/0118719 A1 6/2005 Schmidt et al.
 2009/0155768 A1 6/2009 Scholl et al.
 2010/0093617 A1 4/2010 Barrangou et al.
 2011/0002889 A1 1/2011 Barrangou et al.
 2011/0008369 A1 1/2011 Finnefrock et al.
 2011/0136688 A1 6/2011 Scholl et al.
 2012/0177645 A1 7/2012 Langermann et al.
 2012/0269859 A1 10/2012 Minato
 2012/0294796 A1 11/2012 Johnson et al.
 2013/0011828 A1 1/2013 Barrangou et al.
 2014/0022021 A1 1/2014 Kusachi
 2014/0105912 A1 4/2014 Noelle
 2014/0107092 A1 4/2014 Meyerson et al.
 2014/0199767 A1 7/2014 Barrangou et al.
 2014/0234972 A1 8/2014 Zhang
 2014/0294898 A1 10/2014 Miller et al.
 2014/0341920 A1 11/2014 Noelle
 2015/0031134 A1 1/2015 Zhang et al.
 2015/0050699 A1 2/2015 Siksnys et al.
 2015/0064138 A1 3/2015 Lu et al.
 2015/0125463 A1 5/2015 Cogswell et al.
 2015/0132419 A1 5/2015 Arvik et al.
 2015/0184139 A1 7/2015 Zhang et al.
 2015/0353905 A1 12/2015 Weiss
 2016/0009805 A1 1/2016 Kowanetz et al.
 2016/0009813 A1 1/2016 Themeli
 2016/0024510 A1 1/2016 Bikard et al.
 2016/0115488 A1 4/2016 Zhang et al.
 2016/0115489 A1 4/2016 Zhang et al.
 2016/0159905 A1 6/2016 Abdiche
 2016/0160186 A1 6/2016 Parsley et al.
 2016/0324938 A1 11/2016 Bikard et al.
 2016/0333348 A1 11/2016 Clube et al.
 2016/0345578 A1 12/2016 Barrangou et al.
 2016/0347836 A1 12/2016 Grosso
 2016/0354416 A1 12/2016 Gajewski et al.
 2017/0022499 A1 1/2017 Lu et al.
 2017/0037416 A1 2/2017 Barrangou et al.
 2017/0173085 A1 6/2017 Kovarik et al.
 2017/0174713 A1* 6/2017 Du C07F 9/65583
 2017/0175142 A1 6/2017 Zhang et al.
 2017/0196225 A1 7/2017 Clube et al.
 2017/0246221 A1 8/2017 Clube et al.
 2017/0304443 A1 10/2017 Lebwohl et al.
 2017/0327582 A1 11/2017 Bissonnette et al.
 2017/0340733 A1 11/2017 Cao et al.
 2018/0015131 A1 1/2018 Gajewski et al.
 2018/0055852 A1 3/2018 Kutok et al.
 2018/0064114 A1 3/2018 Clube et al.
 2018/0064115 A1 3/2018 Clube et al.
 2018/0070594 A1 3/2018 Clube et al.
 2018/0084785 A1 3/2018 Clube et al.
 2018/0084786 A1 3/2018 Clube et al.
 2018/0140698 A1* 5/2018 Clube A61P 37/00
 2018/0146681 A1 5/2018 Clube et al.
 2018/0155721 A1 6/2018 Lu et al.
 2018/0155729 A1 6/2018 Beisel et al.
 2018/0179547 A1 6/2018 Zhang et al.
 2018/0200342 A1 7/2018 Bikard et al.
 2018/0303934 A1 10/2018 Clube et al.
 2018/0326057 A1 11/2018 Clube et al.
 2018/0371405 A1 12/2018 Barrangou et al.
 2019/0015441 A1* 1/2019 Shachar A61K 31/7088
 2019/0021343 A1 1/2019 Barrangou et al.
 2019/0070233 A1 3/2019 Yeung
 2019/0133135 A1 5/2019 Clube
 2019/0134194 A1 5/2019 Clube
 2019/0160120 A1 5/2019 Haaber et al.
 2019/0230936 A1 8/2019 Clube
 2019/0240325 A1 8/2019 Clube
 2019/0240326 A1 8/2019 Clube
 2019/0256900 A1 8/2019 Zhang
 2019/0321468 A1 10/2019 Clube et al.
 2019/0321469 A1 10/2019 Clube et al.

2019/0321470 A1 10/2019 Clube et al.
 2020/0030444 A1 1/2020 Clube et al.
 2020/0157237 A1* 5/2020 Regev C07K 16/2818

FOREIGN PATENT DOCUMENTS

RU 2531343 C2 10/2014
 WO WO-1995/001994 A1 1/1995
 WO WO-1998/042752 A1 10/1998
 WO WO-2000/037504 A2 6/2000
 WO WO-2000/037504 A3 6/2000
 WO WO-2001/014424 A2 3/2001
 WO WO-2001/014424 A3 3/2001
 WO WO-2005/003168 A2 1/2005
 WO WO-2005/003168 A3 1/2005
 WO WO-2005/009465 A1 2/2005
 WO WO-2005/046579 A2 5/2005
 WO WO-2005/046579 A3 5/2005
 WO WO-2006/003179 A2 1/2006
 WO WO-2006/003179 A3 1/2006
 WO WO-2006/072625 A2 7/2006
 WO WO-2006/072625 A3 7/2006
 WO WO-2006/072626 A1 7/2006
 WO WO-2006/121168 A1 11/2006
 WO WO-2007/025097 A2 3/2007
 WO WO-2007/042573 A2 4/2007
 WO WO-2007/042573 A3 4/2007
 WO WO-2008/084106 A2 7/2008
 WO WO-2008/084106 A9 7/2008
 WO WO-2008/108989 A2 9/2008
 WO WO-2008/132601 A1 11/2008
 WO WO-2009/044273 A2 4/2009
 WO WO-2009/044273 A3 4/2009
 WO WO-2009/101611 A1 8/2009
 WO WO-2009/114335 A2 9/2009
 WO WO-2009/114335 A3 9/2009
 WO WO-2010/011961 A1 1/2010
 WO WO-2010/027827 A2 3/2010
 WO WO-2010/027827 A3 3/2010
 WO WO-2010/065939 A1 6/2010
 WO WO-2010/075424 A2 7/2010
 WO WO-2011/014438 A1 2/2011
 WO WO-2011/066342 A2 6/2011
 WO WO-2011/066342 A3 6/2011
 WO WO-2012/071411 A2 5/2012
 WO WO-2012/071411 A3 5/2012
 WO WO-2012/160448 A2 11/2012
 WO WO-2012/160448 A3 11/2012
 WO WO-2012/164565 A1 12/2012
 WO WO-2013/006490 A2 1/2013
 WO WO-2013/006490 A3 1/2013
 WO WO-2013/025779 A1 2/2013
 WO WO-2013/067492 A1 5/2013
 WO WO-2013/176772 A1 11/2013
 WO WO-2014/018423 A2 1/2014
 WO WO2014015252 A1 1/2014
 WO WO-2014/093595 A1 6/2014
 WO WO-2014/093661 A2 6/2014
 WO WO-2014/093661 A3 6/2014
 WO WO-2014/093661 A9 6/2014
 WO WO-2014/124226 A1 8/2014
 WO WO-2015/016718 A1 2/2015
 WO WO-2015/034872 A2 3/2015
 WO WO-2015/069682 A2 5/2015
 WO WO-2015/071474 A2 5/2015
 WO WO-2015/075688 A1 5/2015
 WO WO-2015/088643 A1 6/2015
 WO WO-2015/136541 A2 9/2015
 WO WO-2015/136541 A3 9/2015
 WO WO-2015/148680 A1 10/2015
 WO WO-2015/153940 A1 10/2015
 WO WO-2015/155686 A2 10/2015
 WO WO-2015/159068 A1 10/2015
 WO WO-2016/063263 A2 4/2016
 WO WO-2016/063263 A3 4/2016
 WO WO2016084088 A1 6/2016
 WO WO-2016/177682 A1 11/2016
 WO WO-2016/196361 A1 12/2016
 WO WO-2016/196605 A1 12/2016

(56) **References Cited**

FOREIGN PATENT DOCUMENTS

WO WO-2016/205276 A1 12/2016
 WO WO-2017/009399 A1 1/2017
 WO WO-2017/042347 A1 3/2017
 WO WO-2017/058751 A1 4/2017
 WO WO-2017/112620 A1 6/2017
 WO WO-2017/211753 A1 * 6/2017
 WO WO-2018/064165 A2 4/2018
 WO WO-2018/081502 A1 5/2018
 WO WO-2018/115519 A1 6/2018
 WO WO-2018/222969 A1 12/2018

OTHER PUBLICATIONS

Mayo clinic, <https://www.mayoclinic.org/diseases-conditions/ meningitis/symptoms-causes/syc-20350508>, accessed Jul. 20, 2018.*
 Mayo clinic, <https://www.mayoclinic.org/diseases-conditions/ pneumonia/symptoms-causes/syc-20354204>, accessed Jul. 20, 2018.*
 Mayo Clinic, <https://www.mayoclinic.org/diseases-conditions/cystitis/ symptoms-causes/syc-20371306>, accessed Jul. 20, 2018.*
 healthline, <https://www.healthline.com/health/acute-nephritic- syndrome#types>, accessed Jul. 20, 2018.*
 Merck Manual, <https://www.merckmanuals.com/home/skin-disorders/ bacterial-skin-infections/overview-of-bacterial-skin-infections>, accessed Jul. 20, 2018.*
 Colgate, <https://www.colgate.com/en-us/oral-health/conditions/mouth- sores-and-infections/eight-common-oral-infections-0615>, accessed Jul. 20, 2018.*
 Zhang et al., *Oncotarget*, 2016; 7(45): 73068-73079.*
 Beisel et al., *Genome Biology*, 2014; 15: 516-519.*
 Clubbe, Jasper U.S. Appl. No. 16/028,311; projected PG publica- tion: Oct. 25, 2018.*
 Denham et al., Case Reports in Infectious Diseases, 2018; Article ID 8438701: 1-3 (Year: 2018).*
 Mancha-Agresti et al., *Molecular Therapy: Methods & Clinical Development*, 2017; 4: 83-91. (Year: 2017).*
 YourGenome, CRISPR/CAS9, <https://www.yourgenome.org/facts/ what-is-crispr-cas9>, accessed Jan. 6, 2020 (Year: 2020).*
 Abernethy, J. K. et al, (Mar. 2015, e-pub. Jan. 14, 2015). "Thirty Day All-Cause Mortality in Patients With *Escherichia coli* Bacteraemia in England," *Clin. Microbiol. Infect.* 21:251.e1-251.e8).
 Aklujkar et al. (2010) "Interference With Histidyl-tRNA Synthetase by a CRISPR Spacer Sequence as a Factor in the Evolution of *Pelobacter carbinolicus*," *BMC Evolutionary Biology* 10:203, 15 pages.
 Anatoliotaki, M. et al. (2004). "Bloodstream Infections in Patients with Solid Tumors: Associated Factors, Microbial Spectrum and Outcome," *Infection* 2004, 32(2):65-71.
 Ang, Y.L.E. et al. (2015). "Best Practice in the Treatment of Advanced Squamous Cell Lung Cancer," *Ther. Adv. Respir. Dis.* 9(5):224-235.
 Arnold, I.C. et al. (Apr. 8, 2015, e-pub. Mar. 4, 2015). "*Helicobacter hepaticus* Infection in BALB/c Mice Abolishes Subunit-Vaccine- Induced Protection Against *M. tuberculosis*," *Vaccine* 33(15):1808-1814.
 Arslan, Z. et al. (May 7, 2013). "RcsB-BglJ-Mediated Activation of Cascade Operon Does Not Induce the Maturation of CRISPR RNAs in *E. coli* K12," *RNA Biology* 10(5):708-715.
 Arumugam et al. (May 12, 2011). "Enterotypes of the human gut microbiome," *Nature* 473(7346):174-180, 16 pages.
 Belizario et al. (Oct. 6, 2015). "Human Microbiomes and Their Roles in Dysbiosis, Common Diseases, and Novel Therapeutic Approaches," *Frontiers in Microbiology* 6(1050):1-16.
 Bikard et al. (Aug. 16, 2012). "CRISPR Interference Can Prevent Natural Transformation and Virulence Acquisition during In Vivo Bacterial Infection," *Cell Host & Microbe* 12(2):177-186.
 Bikard et al. (Nov. 2014). "Development of Sequence-Specific Antimicrobials Based on Programmable CRISPR-Cas Nucleases," *Nature Biotechnology* 32(11):1146-1151, 16 pages.

Bikard et al. (2013, e-pub. Jun. 12, 2013). "Programmable Repres- sion and Activation of Bacterial Gene Expression Using an Engi- neered CRISPR-Cas System," *Nucleic Acids Research* 41(15):7429- 7437.
 Broaders et al. (Jul./Aug. 2013). "Mobile Genetic Elements of the Human Gastrointestinal Tract," *Gut Microbes* 4(4):271-280.
 Bugrysheva et al. (Jul. 2011, E-Pub. Apr. 29, 2011). "The Histone- Like Protein Hlp Is Essential for Growth of *Streptococcus pyogenes*: Comparison of Genetic Approaches to Study Essential Genes," *Appl. Environ. Microbiol.* 77(13):4422-28.
 Bullman et al. (Nov. 23 2017). "Analysis of *Fusobacterium* Persist- ence and Antibiotic Response in Colorectal Cancer," *Science* 10 pages.
 Citorik et al. (Nov. 2014, e-pub Sep. 21, 2014). "Sequence-Specific Antimicrobials Using Efficiently Delivered RNA-Guided Nucleases," *Nat. Biotechnol.* 32(11):1141-1145, 18 pages.
 Cochrane, K. et al. (2016, e-pub. Nov. 3, 2015). "Complete Genome Sequences and Analysis of the *Fusobacterium nucleatum* Subspe- cies *Animalis* 7-1 Bacteriophage Φ Funu1 and Φ Funu2," *Anaerobe* 38:125-129, 12 pages.
 Coyne et al. (2014). "Evidence of Extensive DNA Transfer between *Bacteroidales* Species Within the Human Gut," *mBio* 5(3):e01305- 14, 12 pages.
 Daillere, R. et al. (Oct. 18, 2016). "*Enterococcus hirae* and *Barnesiella intestinhominis* Facilitate Cyclophosphamide-Induced Therapeutic Immunomodulatory Effects," *Immunity* 95:931-943.
 Deeks, E.D. (2014, e-pub. Jul. 15, 2014). "Nivolumab: A Review of Its Use in Patients With Malignant Melanoma," *Drugs* 74:1233- 1239.
 Del Castillo, M. et al. (Dec. 1, 2016). The Spectrum of Serious Infections Among Patients Receiving Immune Checkpoint Block- ade for the Treatment of Melanoma *Clin. Infect. Dis.* 63:1490-1493.
 De Filippo et al. (Aug. 33 2010). "Impact of Diet in Shaping Gut Microbiota Revealed by a Comparative Study in Children From Europe and Rural Africa," *Proc. Natl. Acad. Sci. USA* 107(33):14691- 14696, 6 pages.
 De Paepe et al. (Mar. 28, 2014). "Bacteriophages: An Underesti- mated Role in Human and Animal Health?" *Frontiers in Cellular and Infection Microbiology* 4(39):1-11.
 DeRosa, L. et al. (2018, e-pub. Mar. 30, 2018). "Negative Associa- tion of Antibiotics on Clinical Activity of Immune Checkpoint Inhibitors in Patients With Advanced Renal Cell and Non-Small- Cell Lung Cancer," *Annals of Oncology*, abstract only.
 Diez-Villasenor et al. (May 2013). "CRISPR-Spacer Integration Reporter Plasmids Reveal Distinct Genuine Acquisition Specificities Among CROSPR-Cas 1-E Variants of *Escherichia coli*," *RNA Biology* 10(5):792-802.
 Dutilh et al. (Jul. 24, 2014). "A Highly Abundant Bacteriophage Discovered in the Unknown Sequences of Human Faecal Metagenomes," *Nature Communications* 5(4498):1-10.
 Edgar et al. (Dec. 2010). "The *Escherichia coli* CRISPR System Protects From λ Lysogenization, Lysogens, and Prophage Induc- tion," *Journal of Bacteriology* 192(23):6291-6294.
 Foca et al. (2015, e-pub. Apr. 7, 2015). Gut Inflammation and Immunity: What Is the Role of the Human Gut Virome? *Mediators of Inflammation* 2015(326032):1-7.
 Fujita, K. et al. (2017). "Emerging Concern of Infectious Diseases in Lung Cancer Patients Receiving Immune Checkpoint Inhibitor Therapy," *Eur. Resp. J.* 50, OA1478. (Abstract Only).
 Galperin, M.Y. (Dec. 2013). "Genome Diversity of Spore-Forming *Firmicutes*," *Microbiology Spectrum* 1(2):TBS-0015-2012, 27 pages.
 Garon, E.B. et al. (Oct. 2015). "Current Perspectives in Immunotherapy for Non-Small Cell Lung Cancer," *Seminars in Oncology* 42(5 Supp. 2):S11-S18.
 Geller, L.T. et al. (Sep. 15, 2017). "Potential Role of Intratumor Bacteria in Mediating Tumor Resistance to the Chemotherapeutic Drug Gemcitabine," *Cancer*, 6 pages.
 Goldwater, P.N. et al. (2012). "Treatment of Enterohemorrhagic *Escherichia coli* (EHEC) Infection and Hemolytic Uremic Syn- drome (HUS)," *BMC Medicine* 10:12, 8 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Golubovskaya, V. et al. (Mar. 15, 2016). "Different Subsets of T Cells, Memory, Effector Functions, and CAR-T Immunotherapy," *Cancers* 8(36), 12 pages.
- Gomaa et al. (Jan. 28, 2014). "Programmable Removal of Bacterial Strains by Use of Genome-Targeting CRISPR-Cas Systems," *mBio*, 5(1):e000928-13.
- Gopalakrishnan, V. et al. (Jan. 5, 2018). "Gut Microbiome Modulates Response to Anti-PD-1 Immunotherapy in Melanoma Patients," *Science* 359:97-103, 20 pp.
- Gudbergsdottir et al. (2011, e-pub. Nov. 18, 2010). "Dynamic Properties of the *Sulfolobus* CRISPR/Cas and CRISPR/Cmr Systems When Challenged With Vector-Borne Viral and Plasmid Genes and Protospacers," *Molecular Microbiology* 79(1):35-49.
- Gudiol, C. et al. (2016). "Bloodstream Infections in Patients With Solid Tumors," *Virulence* 7(3):298-308.
- Guedan, S. et al. (Aug. 14, 2014). "ICOS-Based Chimeric Antigen Receptors Program Bipolar T_H17/T_H1 Cells," *Blood* 124(7):1070-1080.
- Ha, Y.E. et al. (2013). "Epidemiology and Clinical Outcomes of Bloodstream Infections Caused by Extended-Spectrum β -Lactamase-Producing *Escherichia coli* in Patients With Cancer," *Int. J. Antimicrob. Agen.* 42(5):403-409.
- Hargreaves et al. (Aug. 26, 2014). "Abundant and Diverse Clustered Regularly Interspaced Short Palindromic Repeat Spacers in *Clostridium difficile* Strains and Prophages Target Multiple Phage Types within This Pathogen," *mBio* 5(5):e01045-13.
- Hooper, L.V. et al. (Jun. 8, 2012). "Interactions Between the Microbiota and the Immune System," *Science* 336(6086):1268-1273, 16 pages.
- Huddleston. (Jun. 20, 2014). "Horizontal Gene Transfer in the Human Gastrointestinal Tract: Potential Spread of Antibiotic Resistance Genes," *Infection and Drug Resistance* 7:167-176.
- Hurwitz, A.A. et al. (Aug. 1998). "CTLA-4 Blockade Synergizes With Tumor-Derived Granulocyte-Macrophage Colony-Stimulating Factor for Treatment of an Experimental Mammary Carcinoma," *Proc. Natl. Acad. Sci. USA* 95:10067-10071.
- Ivanov, I.I. et al. (May 2010). "Segmented Filamentous Bacteria Take the Stage," *Mucosal Immunol.* 3(3):209-212, 7 pages.
- Jiang et al. (Nov. 2013). "Demonstration of CRISPR/Cas9/sgRNA-Mediated Targeted Gene Modification in Arabidopsis, Tobacco, Sorghum and Rice," *Nucleic Acids Research* 41(20):e188, 12 pages.
- Jinek et al. (Aug. 17, 2012). "A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity," *Science* 337(6096):816-821.
- Jones, R.B. et al. (2008). "Tim-3 Expression Defines a Novel Population of Dysfunctional T Cells With Highly Elevated Frequencies in Progressive HIV-1 Infection," *J. Exp. Med.* 205(12):2763-2779.
- Karch, H. et al. (Jul. 1999). "Epidemiology and diagnosis of Shiga toxin-producing *Escherichia coli* infections," *Diagnostic Microbiology and Infectious Disease* 34(3):229-243.
- Khoja, L. et al. (2015). "Pembrolizumab," *Journal for ImmunoTherapy of Cancer* 3(36):1-13.
- Kosiewicz, M.M. et al. (2014, e-pub. Mar. 26, 2014). "Relationship Between Gut Microbiota and Development of T Cell Associated Disease," *FEBS Lett.* 588:4195-4206.
- Lopez-Sanchez et al. (2012, e-pub. Jul. 27, 2012). "The Highly Dynamic CRISPR1 System of *Streptococcus agalactiae* Controls the Diversity of its Mobilome," *Molecular Microbiology* 85(6):1057-1071.
- Luo et al. (2015, e-pub. Oct. 17, 2014). "Repurposing Endogenous Type I CRISPR-Cas Systems for Programmable Gene Repression," *Nucleic Acids Research* 43(1):674-681.
- Mahoney, K.M. et al. (2015). "The Next Immune-Checkpoint Inhibitors: PD-1/PD-L1 Blockade in Melanoma," *Clinical Therapeutics* 37(4):764-779.
- Marin, M. et al. (2014). "Bloodstream Infections in Patients With Solid Tumors Epidemiology, Antibiotic Therapy, and Outcomes in 528 Episodes in a Single Cancer Center," *Medicine* 93:143-149.
- Marraffini et al. (Dec. 19, 2008). "CRISPR Interference Limits Horizontal Gene Transfer in *Staphylococci* by Targeting DNA," *Science* 322(5909):1843-1845, 12 pages.
- Matson, V. et al. (2018). "The Commensal Microbiome Is Associated With Anti-PD-1 Efficacy in Metastatic Melanoma Patients," *Science* 359(6371):104-108.
- Matsushiro, A. et al. (Apr. 1999). "Induction of Prophages of Enterohemorrhagic *Escherichia coli* O157:H7 with Norfloxacin," *Journal of Bacteriology* 181(7):2257-2260.
- Medina-Aparicio, L. et al. (Mar. 11, 2011, e-pub. Mar. 11, 2011). "The CRISPR/Cas Immune System Is an Operon Regulated by LeuO, H-NS, and Leucine-Responsive Regulatory Protein in *Salmonella enterica* Serovar Typhi," *Journal of Bacteriology* 193(10):2396-2407.
- Mick et al. (May 2013). "Holding a Grudge: Persisting Anti-Phage CRISPR Immunity in Multiple Human Gut Microbiomes," *RNA Biology* 10(5):900-906.
- Mills et al. (Jan./Feb. 2013). "Movers and Shakers: Influence of Bacteriophages in Shaping the Mammalian Gut Microbiota," *Gut Microbes* 4(1):4-16.
- Nakeamura et al. (Nov. 2008). "Metagenomic Diagnosis of Bacterial Infections," *Emerging Infectious Diseases* 14(11):1784-1786.
- Nale, J.Y. et al. (May 18, 2012). "Diverse Temperate Bacteriophage Carriage in *Clostridium difficile* 027 Strains," *PLoS One* 7(5):e37263, 9 pages.
- Nelson, M.H. et al. (2015). "Harnessing the Microbiome to Enhance Cancer Immunotherapy," *Journal of Immunology Research* 2015:Article 368736, 12 pages.
- Nowak, P. et al. (Nov. 28, 2015). "Gut Microbiota Diversity Predicts Immune Status in HIV-1 Infection," *AIDS* 29(18):2409-2418.
- Okazaki, T. et al. (2007). "PD-1 and PD-1 Ligands: From Discovery to Clinical Application," *Intern. Immun.* 19(7):813-824.
- Pardoll, D.M. (2012). "The Blockade of Immune Checkpoints in Cancer Immunotherapy," *Nat. Rev. Cancer* 12(4): 252-264.
- Patterson et al. (Dec. 15, 2016). "Quorum Sensing Controls Adaptive Immunity Through the Regulation of Multiple CRISPR-Cas Systems," *Mol. Cell* 64(6):1102-1108.
- Patterson, A.G. et al. (Mar. 27, 2017). "Regulation of CRISPR-Cas Adaptive Immune Systems," *Current Opinion in Microbiology* 37:1-7.
- Pawluk et al. (Apr. 15, 2014). "A New Group of Phage Anti-CRISPR Genes Inhibits the Type I-E CRISPR-Cas System of *Pseudomonas aeruginosa*," *mBio*, 5(2):e00896.
- Purdy, D. et al. (2002). "Conjugative Transfer of Clostridial Shuttle Vectors From *Escherichia coli* to *Clostridium difficile* Through Circumvention of the Restriction Barrier," *Molec. Microbiology* 46(2):439-452.
- Richter, C. et al. (Oct. 19, 2012, e-pub. Oct. 19, 2012). "Function and Regulation of Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) / CRISPR Associated (Cas) Systems," *Viruses* 4(12):2291-2311.
- Ridaura et al. (Sep. 6, 2013). "Cultured Gut Microbiota From Twins Discordant for Obesity Modulate Adiposity and Metabolic Phenotypes in Mice," *Science* 341(6150):1241214, 22 pages.
- Routy, B. et al. (Jan. 5, 2018, e-pub. Nov. 2, 2017). "Gut Microbiome Influences Efficacy of PD-1-Based Immunotherapy Against Epithelial Tumors," *Science* 359(6371):91-97.
- Roy, S. et al. (May 2017, e-pub. Mar. 17, 2017). "Microbiota: A Key Orchestrator of Cancer Therapy," *Nat. Rev. Cancer* 17(5):271-285.
- Safdar, N. et al. (Jun. 4, 2002). "The Commonality of Risk Factors for Nosocomial Colonization and Infection With Antimicrobial-Resistant *Staphylococcus aureus*, Enterococcus, Gram-Negative Bacilli, *Clostridium difficile*, and *Candida*," *Ann. Intern. Med.* 136(11):834-844.
- Samonis, G. et al. (Sep. 2013, e-pub. Apr. 27, 2013). "A Prospective Study of Characteristics and Outcomes of Bacteremia in Patients With Solid Organ or Hematologic Malignancies," *Support Care Cancer* 21(9):2521-2526.
- Seed et al. (Feb. 27, 2013). "A Bacteriophage Encodes Its Own CRISPR/Cas Adaptive Response to Evade Host Innate Immunity," *Nature* 494(7438):489-491.
- Selle et al. (Apr. 1, 2015). "Harnessing CRISPR-Cas Systems for Bacterial Genome Editing," *Trends in Microbiology* 23(4):225-232.

(56)

References Cited

OTHER PUBLICATIONS

- Sharan, S.K. et al. (2009). "Recombineering: A Homologous Recombination-Based Method of Genetic Engineering," *Nat. Protoc.* 4(2):206-223, 37 pages.
- Shoemaker et al. (Feb. 2001). "Evidence for Extensive Resistance Gene Transfer Among *Bacteroides* spp. and Among *Bacteroides* and Other Genera in the Human Colon," *Appl. Environ. Microbiol.* 67(2):561-68.
- Sivan, A. et al. (Nov. 27, 2015, e-pub Nov. 5, 2015). "Commensal *Bifidobacterium* Promotes Antitumor Immunity and Facilitates Anti-PD-L1 Efficacy," *Science* 350(6264):1084-1089, 13 pages.
- Slutsker, L. et al. (Apr. 1998). "A Nationwide Case-Control Study of *Escherichia coli* O157:H7 Infection in the United States," *J. Infect. Dis.* 177(4):962-966.
- Soutourina, O.A. et al. (May 9, 2013). "Genome-Wide Identification of Regulatory RNAs in the Human Pathogen *Clostridium difficile*," *PLoS Genet.* 9(5):e1003493, 18 pages.
- Stern et al. (Aug. 2010). Self-Targeting by CRISPR: Gene Regulation or Autoimmunity? *Trends Genet.* 26(8):335-340, 10 pages.
- Stern et al. (2012). "CRISPR Targeting Reveals a Reservoir of Common Phages Associated With the Human Gut Microbiome," *Genome Research* 22(10):1985-1994.
- Stiefel et al. (Aug. 2014, e-pub. May 27, 2014). "Gastrointestinal Colonization With a Cephalosporinase-Producing *Bacteroides* Species Preserves Colonization Resistance Against Vancomycin-Resistant *Enterococcus* and *Clostridium difficile* in Cephalosporin-Treated Mice," *Antimicrob. Agents Chemother.* 58(8):4535-4542.
- Tan, J. (Dec. 17, 2015). "Immunotherapy Meets Microbiota," *Cell* 163:1561.
- Tarr, P.I. et al. (Mar. 19-25, 2005). "Shiga-Toxin-Producing *Escherichia coli* and Haemolytic Uraemic Syndrome," *Lancet* 365(9464):1073-1086.
- Topalian et al. (Jun. 28, 2012). "Safety, Activity, and Immune Correlates of Anti-PD-1 Antibody in Cancer," *N. Engl. J. Med.* 336(26):2443-2454, 19 pages.
- Uchiyama, J. et al. (2013, e-pub. Mar. 8, 2013). "Characterization of *Helicobacter pylori* Bacteriophage KHP30," *Applied and Environmental Microbiology* 79(10):3176-3184.
- Veeranagouda et al. (Jun. 4, 2014). "Identification of Genes Required for the Survival of *B. fragilis* Using Massive Parallel Sequencing of a Saturated Transposon Mutant Library," *BMC Genomics* 15:429, 11 pages.
- Velasco, E. et al. (2006). "Comparative Study of Clinical Characteristics of Neutropenic and Non-Neutropenic Adult Cancer Patients With Bloodstream Infections," *Eur. J. Clin. Microbiol. Infect. Dis.* 25:1-7.
- Vercoe et al. (Apr. 18, 2013). "Cytotoxic Chromosomal Targeting by CRISPR/Cas Systems Can Reshape Bacterial Genomes and Expel or Remodel Pathogenicity Islands," *PLOS Genetics* 9(4):e1003454, 13 pages.
- Vétizou, M. et al. (Nov. 27, 2015, e-pub Nov. 5, 2015). "Anticancer Immunotherapy by CTLA-4 Blockade Relies on the Gut Microbiota," *Science* 350(6264):1079-1084, 13 pages.
- Villarino, N.F. et al. (Feb. 23, 2016, e-pub. Feb. 8, 2016). "Composition of the Gut Microbiota Modulates the Severity of Malaria," *Proc. Natl. Acad. Sci. USA* 113(8):2235-2240.
- Wagner, P.L. (2002). "Bacteriophage Control of Shiga Toxin 1 Production and Release by *Escherichia coli*," *Molecular Microbiology* 44(4):957-970.
- Walters et al. (Nov. 17, 2014). "Meta-Analyses of Human Gut Microbes Associated With Obesity and IBD," *FEBS Letters* 588(22):4223-4233, 34 pages.
- Wei et al. (2015, e-pub. Jan. 14, 2015). "Sequences Spanning the Leader-Repeat Junction Mediate CRISPR Adaptation to Phage in *Streptococcus thermophilus*," *Nucleic Acids Research* 43(3):1749-1758.
- Westra, E.R. et al. (Sep. 1, 2010). "H-NS-c of CRISPR-Based Immunity in *Escherichia coli* K12 Can Be Relieved by the Transcription Activator LeuO:LeuO Activates CRISPR-Based Immunity," *Molecular Microbiology* 77(6):1380-1393.
- Wexler. (Oct. 2007). "*Bacteroides*: the Good, the Bad, and the Nitty-Gritty," *Clinical Microbiology Reviews* 20(4):593-621.
- Wong, C.S. et al. (Jun. 29, 2000). "The Risk of the Hemolytic-Uremic Syndrome After Antibiotic Treatment of *Escherichia coli* O157:H7 Infections," *N. Engl. J. Med.* 342(26):1930-1936, 13 pages.
- Yao, J. et al. (2016, e-pub. May 9, 2016). "A Pathogen-Selective Antibiotic Minimizes Disturbance to the Microbiome," *Antimicrob. Agents Chemother.* 24 pages.
- Yosef et al. (2011). "High-Temperature Protein G Is Essential for Activity of the *Escherichia coli* Clustered Regularly Interspaced Palindromic Repeats (CRISPR)/Cas system," *Proc. Natl. Acad. Sci. USA* 108(50):20136-20141.
- Zembower, T.R. (2004). "Epidemiology of Infections in Cancer Patients," in *Infectious Complications in Cancer Patients*, Springer International Publishing Switzerland, 48 pages.
- Zimmerhackl, L.B. (Jun. 29, 2000). "*E. coli*, Antibiotics, and the Hemolytic-Uremic Syndrome," *N. Engl. J. Med.* 342(26):1990-1991.
- Zitvogel et al. (Jan. 2015). "Cancer and the Gut Microbiota: An Unexpected Link," *Sci. Transl. Med.* 7(271):271ps1, 10 pages.
- Zitvogel, L. et al. (Mar. 2018). "The Microbiome in Cancer Immunotherapy: Diagnostic Tools and Therapeutic Strategies," *Science* 359(6382):1366-1370.
- International Search Report for PCT/EP2016/059803, dated Jun. 30, 2016, filed May 3, 2016, 6 pages.
- International Search Report and the Written Opinion of the International Searching Authority for PCT/EP2018/066954, dated Oct. 23, 2018, filed Jun. 25, 2018, 14 pages.
- Written Opinion for PCT/EP2016/059803, dated Jun. 30, 2016, filed May 3, 2016, 6 pages.
- U.S. Appl. No. 15/985,658, filed May 21, 2018, for Haaber et al. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. § 1.98(a)(2)(iii) issued by the Office on Sep. 21, 2004).
- U.S. Appl. No. 16/041,595, filed Jul. 20, 2018, for Club et al. (Copy not attached) (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. § 1.98(a)(2)(iii) issued by the Office on Sep. 21, 2004).
- Brouns, S.J.J. et al. (Aug. 15, 2008). "Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes," *Science* 321:960-964.
- Brouns, S.J.J. et al. (Aug. 15, 2008). Supplemental Material for "Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes," *Science* 321:960-964.
- Consumer Updates (2019). "Combating Antibiotic Resistance," retrieved from <<https://www.fda.gov/ForConsumers/ConsumerUpdates/ucm092810.htm>>, last visited Jan. 28, 2019.
- Fact Sheet (Oct. 2010). "Antimicrobial Resistance," *National Institutes of Health*, 1-2.
- Gomaa, A.A. et al. (Jan./Feb. 2014). Supplemental Material to "Programmable Removal of Bacterial Strains by Use of Genome-Targeting CRISPR-Cas Systems," *American Society for Microbiology* 5(1):1-9.
- Krom, R.J. et al. (Jul. 8, 2015). "Engineered Phagemids for Nonlytic, Targeted Antibacterial Therapies," *Nano Letters* 15(7):4808-4813.
- Navarre, L. et al. (2007). "Silencing of Xenogeneic DNA by H-NS—Facilitation of Lateral Gene Transfer in Bacteria by a Defense System That Recognizes Foreign DNA," *Genes & Development* 21:1456-1471.
- News (May 22, 2018). "UK Government and Bill & Melinda Gates Foundation Join Carb-X Partnership in Fight Against Superbugs: Millions Earmarked to Boost Research Into New Life-Saving Products to Address the Global Rise of Drug-Resistant Bacteria," *Combating Antibiotic Resistant Bacteria*, 7 pages.
- Norris, J.S. et al. (2000). "Prokaryotic Gene Therapy to Combat Multidrug Resistant Bacterial Infection," *Gene Therapy* 7:723-725.
- Pires, D.P. et al. (Sep. 2016, e-pub. Jun. 1, 2016). "Genetically Engineered Phages: A Review of Advances Over the Last Decade," *Microbiology and Molecular Biology Reviews* 80(3):523-543.
- Rashid, T. et al. (2013). "The Role of *Klebsiella* in Crohn's Disease With a Potential for the Use of Antimicrobial Measures," *International Journal of Rheumatology* 2013(Article ID 610393):1-9.
- Skenneron, C.T. (May 2011). "Phage Encoded H-NS: A Potential Achilles Heel in the Bacterial Defence System," *PLoS One* 6(5):e20095.

(56)

References Cited

OTHER PUBLICATIONS

- Stoebel, D.M. et al. (2008). "Anti-Silencing: Overcoming H-NS-Mediated Repression of Transcription in Gramnegative Enteric Bacteria," *Microbiology* 154:2533-2545.
- Takeda, T. et al. (2011). "Distribution of Genes Encoding Nucleoid-Associated Protein Homologs in Plasmids," *International Journal of Evolutionary Biology* 2001:685015, 31 pages.
- Vega, N.M. et al. (Oct. 2014). "Collective Antibiotic Resistance: Mechanisms and Implications," *Curr. Opin. Microbiol.* 21:28-34, 14 pages.
- Westwater, C. et al. (Apr. 2003). "Use of Genetically Engineered Phage to Deliver Antimicrobial Agents to Bacteria: An Alternative Therapy for Treatment of Bacterial Infections," *Antimicrobial Agents and Chemotherapy* 47(4):1301-1307.
- International Search Report for PCT/EP2018/082053, dated Mar. 14, 2019, filed Nov. 21, 2018, 9 pages.
- Written Opinion for PCT/EP2018/082053, dated Mar. 14, 2019, filed Nov. 21, 2018, 6 pages.
- Ex Parte Re-Exam, mailed Dec. 10, 2018, for U.S. Appl. No. 90/014,184, filed Aug. 10, 2018, for Reexamination U.S. Pat. No. 9,701,964 102 pages.
- Request for Ex Parte Reexamination dated Aug. 10, 2018, for U.S. Appl. No. 15/160,405, now U.S. Pat. No. 9,701,964, 42 pages.
- Request for Ex Parte Reexamination dated Nov. 1, 2018, for U.S. Appl. No. 15/160,405, now U.S. Pat. No. 9,701,964, 35 pages.
- U.S. Appl. No. 16/192,746, filed Nov. 15, 2018, for Clube et al. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. 1.98(a)(2)(iii) issued by the Office on Sep. 21, 2004).
- U.S. Appl. No. 16/389,376, filed Apr. 19, 2019, for Clube. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. 1.98(a)(2)(iii) issued by the Office on Sep. 21, 2004).
- U.S. Appl. No. 16/389,358, filed Apr. 19, 2019, for Clube. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. 1.98(a)(2)(iii) issued by the Office on Sep. 21, 2004).
- U.S. Appl. No. 16/201,736, filed Nov. 27, 2018, for Martinez et al. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. 1.98(a)(2)(iii) issued by the Office on Sep. 21, 2004).
- U.S. Appl. No. 62/168,355, filed May 29, 2015, Barrangou, R. et al. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. 1.98(a)(2)(iii) issued by the Office on Sep. 21, 2004).
- American Lung Association (2019). "Preventing COPD," retrieved from <https://www.lung.org/lung-health-and-diseases/lung-disease-lookup/copd/symptoms-causes-risk-factors/preventing-copd.html>, last visited Aug. 5, 2019, 1 page.
- Anonymous (Apr. 2016). "Checkpoint Inhibition: A Promising Immunotherapeutic Approach for Colorectal Cancer," *Oncology*, 5(3):1-5, retrieved from <http://www.personalizedmedonc.com/publications/prno/april-2016-vol-5-no-3/checkpoint-inhibition-a-promising-immunotherapeutic-approach-for-colorectal-cancer-2/>, last visited Aug. 27, 2019, 5 pages.
- Bikard, D. et al. (2017, e-pub. Sep. 6, 2017). "Using CRISPR-Cas Systems as Antimicrobials," *Current Opinion in Microbiology* 37:155-160.
- Chan, B.K. et al. (2013). "Phage Cocktails and the Future of Phage Therapy," *Future Microbiol.* 8(6):769-783.
- Dickson, R.P. et al. (Jan./Feb. 2017). "Bacterial Topography of the Healthy Human Lower Respiratory Tract," *American Society for Microbiology* 8(1):e02287-6, 12 pages.
- Hartland, E.L. et al. (Apr. 30, 2013). "Enteropathogenic and Enterohemorrhagic *E. coli*: Ecology, Pathogenesis, and Evolution," *Frontiers in Cellular and Infection Microbiology* 3(15):1-3.
- Healthline (2019). "Cystic Fibrosis," retrieved from <https://www.healthline.com/health/cystic-fibrosis#prevention>, last visited Aug. 5, 2019, 14 pages.
- International Search Report and the Written Opinion of the International Searching Authority for PCT/EP2019/057453, dated Aug. 16, 2019, filed Mar. 25, 2019, 21 pages.
- Jin, Y. et al. (2019, e-pub. Apr. 23, 2019). "The Diversity of Gut Microbiome is Associated With Favorable Responses to Anti-Programmed Death 1 Immunotherapy in Chinese Patients With NSCLC," *Journal of Thoracic Oncology* 14(8):1378-1389.
- Manica, A. et al. (2011, e-pub. Mar. 8, 2011). "In vivo Activity of CRISPR-Mediated Virus Defence in a Hyperthermophilic Archaeon," *Molecular Microbiology* 80(2):481-491.
- Mayo Clinic (2019). "Pulmonary Embolism," retrieved from <https://www.mayoclinic.org/diseases-conditions/pulmonary-embolism/symptoms-causes/syc-20354647>, last visited Aug. 5, 2019, 8 pages.
- Mayo Clinic (2020). "Infectious Diseases," retrieved from <https://www.mayoclinic.org/diseases-conditions/infectious-diseases/diagnosis-treatment/drc-20351179>, last visited Jan. 17, 2020, 5 pages.
- Mayo Clinic (2020). "Malaria," retrieved from <https://www.mayoclinic.org/diseases-conditions/malaria/diagnosis-treatment/drc-20351190>, last visited Jan. 17, 2020, 3 pages.
- Mayo Clinic (2020). "Sexually Transmitted Diseases (STDs)," retrieved from <https://www.mayoclinic.org/diseases-conditions/sexually-transmitted-diseases/stds/diagnosis-treatment/drc-20351246>, last visited January 17, 2020, 5 pages.
- Notice of Intent to Issue Ex Parte Reexamination Certificate, mailed Aug. 12, 2019, for U.S. Appl. No. 90/014,184, filed Aug. 10, 2018, 26 pages.
- Park, A. (Oct. 18, 2011). "A Surprising Link Between Bacteria and Colon Cancer," *Cancer* retrieved from <http://healthland.time.com/2011/10/18/a-surprising-link-between-bacteria-and-colon-cancer/>, last visited Aug. 27, 2019, 3 pages.
- Sepsis Alliance, (Dec. 14, 2017). "What Are Vaccines," Retrieved from <https://www.sepsis.org/sepsisand/prevention-vaccinations/>; last visited Jul. 8, 2019, 3 pages.
- Sepsis Alliance; (Jul. 8, 2019). "Prevention," Retrieved from <https://www.sepsis.org/sepsisand/prevention/>; accessed last visited Jul. 8, 2019, 5 pages.
- Takaishi, H. et al. (2008). "Imbalance in Intestinal Microflora Constitution Could Be Involved in the Pathogenesis of Inflammatory Bowel Disease," *Int. J. Med. Microbiol.* 298:463-472.
- U.S. Appl. No. 16/682,889, filed Nov. 13, 2019, for Clube et al. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. 1.98(a)(2)(iii) issued by the Office on Sep. 21, 2004).

* cited by examiner




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3




FIG. 4




FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 7




FIG. 8

Progression-free survival by fecal diversity:




FIG. 9A

Progression-free survival by low/high abundance of certain bacteria:




FIG. 9B

TREATING AND PREVENTING MICROBIAL INFECTIONS

SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

The content of the following submission on ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing (file name: 786212000300SEQLIST.txt, date recorded: Apr. 30, 2018, size: 21 KB).

TECHNICAL FIELD

The invention provides methods for treating or preventing microbial (eg, bacterial) infections and means for performing these methods. In particular, treatment of infections requiring rapid or durable therapy is made possible, such as for treating acute conditions such as septicemia, sepsis, SIRS or septic shock. The invention is also particularly useful, for example, for treatment of microbes for environmental, food and beverage use. The invention relates inter alia to methods of controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate or fluid in an industrial or domestic system.

The invention also useful for the treatment of pathogenic bacterial infections in subjects receiving a treatment for a disease or condition, such as a transplant or a treatment for cancer, a viral infection or an autoimmune disease.

BACKGROUND

Septicaemia is an acute and serious bloodstream infection. It is also known as bacteraemia, or blood poisoning. Septicaemia occurs when a bacterial infection elsewhere in the body, such as in the lungs or skin, enters the bloodstream. This is dangerous because the bacteria and their toxins can be carried through the bloodstream to a subject's entire body. Septicaemia can quickly become life-threatening. It must be rapidly treated, such as in a hospital. If it is left untreated, septicaemia can progress to sepsis.

Septicaemia and sepsis aren't the same. Sepsis is a serious complication of septicaemia. Sepsis is when inflammation throughout the body occurs. This inflammation can cause blood clots and block oxygen from reaching vital organs, resulting in organ failure. The US National Institutes of Health (NIH) estimates that over 1 million Americans get severe sepsis each year. Between 28 and 50 percent of these patients may die from the condition. When the inflammation occurs with extremely low blood pressure, it's called septic shock. Septic shock is fatal in many cases.

The increase in average age of the population, more people with chronic diseases, on immunosuppressive drugs, and increase in the number of invasive procedures being performed has led to an increased rate of sepsis. People over 65 years old, particularly those who have health issues, are even more susceptible to sepsis than any other group. According to a study published in 2006, while people aged 65 years and older make up about 12% of the American population, they make up 65% of sepsis cases in the hospitals.

Septicaemia is caused by an infection in a part of the body. This infection is typically acute. Many types of bacteria can lead to septicaemia. The exact source of the infection often can't be determined. The most common infections that lead to septicaemia are:

- urinary tract infections
- lung infections, such as pneumonia
- kidney infections
- infections in the abdominal area

5 Bacteria from these infections enter the bloodstream and multiply rapidly, causing acute infection and immediate symptoms.

People who are already in the hospital for something else, such as a surgery, are at a higher risk of developing septicaemia. Secondary infections can occur while in the hospital. These infections are often more dangerous because the bacteria may already be resistant to antibiotics. There is a higher risk of developing septicaemia if the subject:

- has severe wounds or burns
- is very young or very old
- has a compromised immune system, which can occur from diseases such as HIV or leukaemia
- has a urinary or intravenous catheter
- is on mechanical ventilation
- is receiving medical treatments that weakens the immune system, such as chemotherapy or steroid injections

The symptoms of septicaemia usually start very quickly. Even in the first stages of the illness, a person can look very sick. They may follow an injury, surgery, or another localized (eg, confined to one location) infection, like pneumonia. The most common initial symptoms are:

- chills
- elevated body temperature (fever)
- very fast respiration
- rapid heart rate

More severe symptoms will begin to emerge as the septicaemia progresses without proper treatment. These include the following:

- confusion or inability to think clearly
- nausea and vomiting
- red dots that appear on the skin
- reduced urine volume
- inadequate blood flow (shock)

Septicaemia that has started to affect the organs or tissue function is an acute medical emergency. It must be rapidly treated at a hospital. Many people with septicaemia are admitted to a hospital's ICU for treatment and recovery. It is recommended to never take a "wait and see" approach or try to treat the problem at home. It is crucial to get to the hospital right away if the subject is showing signs of septicaemia.

Septicaemia has a number of serious complications. These complications may be fatal if left untreated or if treatment is delayed for too long.

50 Septic Shock

One complication of septicaemia is a serious drop in blood pressure. This is called septic shock. Toxins released by the bacteria in the bloodstream can cause extremely low blood flow, which may result in organ or tissue damage. Septic shock is an acute medical emergency. People with septic shock are usually cared for in a hospital's intensive care unit (ICU). The patient may need to be put on a ventilator, or breathing machine, if in septic shock.

Acute Respiratory Distress Syndrome (ARDS)

Another complication of septicaemia is acute respiratory distress syndrome (ARDS). This is a life-threatening condition that prevents enough oxygen from reaching your lungs and blood. According to the National Heart, Lung, and Blood Institute (NHLBI), ARDS is fatal in about one-third of cases. It often results in some level of permanent lung damage. It can also damage the brain, which can lead to memory problems.

Sepsis

Sepsis occurs when the body has a strong immune response to the infection. This leads to widespread inflammation throughout the body. It is called severe sepsis if it leads to organ failure. People with chronic diseases, such as HIV or cancer, are at a higher risk of sepsis. This is because they have a weakened immune system and cannot fight off the infection on their own. Sepsis causes millions of deaths globally each year and is the most common cause of death in people who have been hospitalized. The worldwide incidence of sepsis is estimated to be 18 million cases per year. In the United States sepsis affects approximately 3 in 1,000 people, and severe sepsis contributes to more than 200,000 deaths per year. Sepsis occurs in 1-2% of all hospitalizations and accounts for as much as 25% of ICU bed utilization.

Early diagnosis is necessary to properly manage sepsis, as initiation of rapid therapy is key to reducing deaths from severe sepsis. Within the first three hours of suspected sepsis, diagnostic studies should include white blood cell counts, measuring serum lactate, and obtaining appropriate cultures before starting antibiotics, so long as this does not delay their use by more than 45 minutes

The most common primary sources of infection resulting in sepsis are the lungs, the abdomen, and the urinary tract. Typically, 50% of all sepsis cases start as an infection in the lungs.

Speed of treatment is essential. Two sets of blood cultures (aerobic and anaerobic) should be taken without delaying the initiation of antibiotics. Cultures from other sites such as respiratory secretions, urine, wounds, cerebrospinal fluid, and catheter insertion sites (in-situ more than 48 hours) can be taken if infections from these sites are suspected. In severe sepsis and septic shock, broad-spectrum antibiotics (usually two, a β -lactam antibiotic with broad coverage, or broad-spectrum carbapenem combined with fluoroquinolones, macrolides, or aminoglycosides) are conventional. However, combination of antibiotics is not recommended for the treatment of sepsis without shock and in immunocompromised persons unless the combination is used to broaden the anti-bacterial activity. The administration of antibiotics is important in determining the survival of the person. Some recommend they be given within one hour of making the diagnosis, stating that for every hour of delay in the administration of antibiotics, there is an associated 6% rise in mortality.

Early goal directed therapy (EGDT) is an approach to the management of severe sepsis during the initial 6 hours after diagnosis. It is a step-wise approach, with the physiologic goal of optimizing cardiac preload, afterload, and contractility. It includes giving early antibiotics.

Neonatal sepsis can be difficult to diagnose as newborns may be asymptomatic. If a newborn shows signs and symptoms suggestive of sepsis, antibiotics are immediately started and are either changed to target a specific organism identified by diagnostic testing or discontinued after an infectious cause for the symptoms has been ruled out.

Approximately 20-35% of people with severe sepsis and 30-70% of people with septic shock die. The Surviving Sepsis Campaign (SSC) is a global initiative to bring together professional organizations in reducing mortality from sepsis. Antibiotics are administered within two hours of admission/diagnosis. For every hour a patient is denied antibiotic therapy after the onset of septic shock, the patient's chance of survival is reduced by 7.9% (Survivesepsis.org 2005)

There is, therefore, a need for a rapid treatment of acute microbial infections, such as bacterial infections associated with septicaemia, sepsis or septic shock. It would also be advantageous if the treatment is durable for many hours. Rapid and durable treatment of microbes is also desirable for is for controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate in industrial and domestic systems.

Acute bacterial infections can, in certain circumstances, be health-threatening or even life-threatening. This may be the case, for example, in cancer patients, transplant patients or other subjects. The need for the treatment of the bacterial infection can become urgent and indeed an immediate focus of attention in the medical care. It would be useful to provide methods of treating such pathogenic bacterial infections in a way that does not adversely undermine the efficacy of the cancer or other separate therapy to which the patient also needs to respond.

SUMMARY OF THE INVENTION

The invention provides a solution by using the action of programmable nuclease cutting of microbe genomes; this is different from the metabolic inhibitor and other mechanisms of action used by beta-lactams and other conventional antibiotics for treating infections. The targeted cutting provides selective microbe killing or reduction of growth or proliferation to treat or prevent infection. Moreover, the inventors have surprisingly found a substantial killing (by several logs) can be achieved very rapidly (eg, within 15 minutes) and sustainable effects can be achieved (eg, for more than 1 hour, and even around 3 hours after treatment commenced) in some embodiments. Thus, the invention provides the following configurations.

In a First Configuration

A programmable nuclease for use in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

In a Second Configuration

A plurality of viruses (eg, phage or phagemids for producing phage) for use with a programmable nuclease in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and viruses wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated; wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject;

5

wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

In a Third Configuration

A composition comprising a plurality of nucleic acids for programming a programmable nuclease in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and the nucleic acids wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated; wherein each nucleic acid encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

In a Fourth Configuration

A CRISPR/Cas system comprising a nuclease according to the invention for use in the method of treatment, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of the microbes.

In a Fifth Configuration

A method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

In a Sixth Configuration

A method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a plurality of viruses wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated; wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

In a Seventh Configuration

A method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising

6

exposing the subject to the nuclease and a plurality of nucleic acids wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated; wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein each nucleic acid encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

In a Eighth Configuration

Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of the invention, in the manufacture of a composition for carrying out a method of treatment as defined herein, wherein the subject is an organism other than a human or animal.

In a Ninth Configuration

Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of the invention, in the manufacture of a composition for carrying out an ex vivo or in vitro a method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.

In a Tenth Configuration

Use of a programmable nuclease in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.

In any Configuration:

For example, the infection is an acute infection. For example, the infection is an acute infection that is rapidly treated. For example, the infection is treated rapidly—for example, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. For example, the treatment is durable—for example, the reduction in infection persists for at least 30 minutes immediately after the first 30 minutes of the treatment. Also, optionally a reduction of the infection by at least 100-fold or 1000-fold is maintained for at least 60 minutes (eg, at least 120 minutes) after commencement of the treatment. Exemplification is provided below which surprisingly demonstrates these, such as a rapid killing that was durable around 3 hours after treatment commenced. For example, the method improves survival of the subject, or improves survival rates in humans or human patients suffering from infection by the microbes of the first species or strain.

The invention also provides a solution to the need for effective treatment of pathogenic bacterial infections in subjects undergoing a cancer or other, separate therapy which must also be efficacious. Thus, the invention further provides:—

In an Eleventh Configuration

A method for treating a pathogenic bacterial infection in a human or animal subject caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a programmable nuclease that is programmed to cut the target site, wherein the subject is suffering from a further disease or condition other than the pathogenic bacterial infection and the method comprises administering a therapy to the subject for treating or preventing the further disease or condition, wherein the nuclease treats the infection and the therapy is efficacious in the presence of the programmed nuclease to treat or prevent the disease or condition.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) that are *E. coli*, *Pseudomonas aeruginosa* or *Klebsiella* bacteria, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-PD-1/PD-L1 axis antibody optionally selected from pembrolizumab, nivolumab, atezolizumab, avelumab and durvalumab; and
- (b) The cancer is selected from melanoma; renal cell carcinoma; bladder cancer; a solid tumour; non-small cell lung cancer (NSCLC); forehead and neck squamous cell carcinoma (HNSCC); Hodgkin's lymphoma; a cancer that overexpresses PD-L and no mutations in EGFR or in ALK; colorectal cancer and hepatocellular carcinoma.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a programmable nuclease that is programmed to cut the target site, wherein the subject is suffering from a cancer and the method comprises administering a cancer therapy to the subject for treating the cancer, wherein the nuclease treats the infection and the cancer therapy is efficacious in the presence of the programmed nuclease to treat the cancer.

In a Twelfth Configuration

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nucle-

ase treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer.

In a Thirteenth Configuration

A programmable nuclease for use in the method of the invention.

In a Fourteenth Configuration

A CRISPR/Cas system comprising a nuclease according to the 13th Configuration for use in the method of the 11th or 12th Configuration, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs (gRNAs) or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of first bacteria.

In a Fifteenth Configuration

A guide RNA or a DNA encoding a guide RNA for use in the system or method of treating a pathogenic bacterial infection.

In a Sixteenth Configuration

A nucleic acid vector comprising the guide RNA or DNA.

In a Seventeenth Configuration

A pharmaceutical composition comprising a first nucleic acid vector (or a plurality thereof) encoding the nuclease and a second nucleic acid vector (or a plurality thereof) encoding the guide RNA.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B show time-kill curves for *Escherichia coli* (EHEC) ATCC43888 strain harboring the CGV system. FIG. 1A shows that CRISPR induction killed 99.98% of the population in 30 minutes (black line). Growth in absence of induction is shown in dashed lines. CRISPR was induced at time-point 0 and monitored until 60 minutes. FIG. 2B shows dilution series (10^1 - 10^6) of drop spots (5 μ l) on LB agar plates of *E. coli* ATCC43888 harboring the CGV system after 30 minutes of induction.

FIG. 2 shows CRISPR killing of target strain *Escherichia coli* (EHEC) ATCC43888 in *Galleria mellonella* larvae. *G. mellonella* larvae were delivered injections of bacteria behind the final left proleg. Approximately 1 h after the injection, CRISPR inducers were administered behind the final right proleg. Larvae were incubated at 37° C. for 2 h and sacrificed. Control bacteria carrying an off-target single guide RNA plasmid were also injected in the control group.

FIG. 2 shows CRISPR-kill curves of *Escherichia coli* (EHEC) ATCC43888 in *Galleria mellonella*. *G. mellonella* larvae were delivered injections of bacteria behind the final left proleg. Approximately 1 h after the injection, CRISPR inducers were administered behind the final right proleg. Larvae were incubated at 37° C. and sacrificed at 0, 1, and 2 h after induction.

FIG. 3 shows *E. coli* ATCC43888 count over time.

FIG. 4 shows Kaplan-Meier survival curves of *Galleria mellonella* larvae infected with *Escherichia coli* (EHEC) ATCC43888. CRISPR induction significantly improves survival of the larvae (black line) compared to the off-target control carrying an off-target single guide RNA plasmid (dashed line).

FIGS. 5A-5B show time-kill curves for *Escherichia coli* Nissle 1917 harboring the CGV system targeting pks. FIG. 5A shows that CRISPR induction killed 99.98% of the population in 15 minutes (black line). Growth in absence of induction is shown in dashed lines. CRISPR was induced at time-point 0 and monitored over 3 h. FIG. 5B shows dilution

series (10^1 - 10^6) of drop spots (5 μ l) on LB agar plates of *E. coli* Nissle 1917 harboring the CGV system after 15 minutes of induction.

FIGS. 6A-6B show time-kill curves for *Escherichia coli* Nissle 1917 harboring the CGV system targeting yapH. FIG. 6A shows that CRISPR induction killed 99.98% of the population in 15 minutes (black line). Growth in absence of induction is shown in dashed lines. CRISPR was induced at time-point 0 and monitored over 3 h. FIG. 6B shows dilution series (10^1 - 10^6) of drop spots (5 μ l) on LB agar plates of *E. coli* Nissle 1917 harboring the CGV system after 15 minutes of induction.

FIG. 7 shows complete killing of transconjugant *C. difficile*. The complete precision killing of *Clostridium difficile* using a gRNA-encoding CRISPR array that was delivered from a probiotic carrier bacterial species by conjugative plasmids as vectors is shown. A carrier bacterium (*E. coli* donor strain containing the vectors) was mated with *Clostridium difficile* which was killed upon delivery of the designed array. This harnessed the endogenous Cas3 machinery of *Clostridium difficile*. A 100% killing of *Clostridium difficile* cells was achieved and is shown in this figure.

FIG. 8 shows the antibiotic treatment during ICI therapy has fatal outcomes. Kaplan Meier curve for overall survival of a validation cohort from the Memorial Sloan Ketterin Cancer Center including n=239 advanced NSCLC patients treated with anti-PD-L1/anti-PD-1 mAb who received (ATB, n=68) or not (no ATB, n=171) antibiotics (ATB) two months before the injection of immune checkpoint blockade. There was a median overall survival of 21.9 months in the absence of antibiotic treatment, compared to an overall survival of 9.8 months with antibiotic treatment. So, the median overall survival in patients treated with classical antibiotics is <50% (or >12 months shorter) that of patients not receiving antibiotic treatment.

FIGS. 9A and 9B show that the gut microbiome modulates the efficacy of anti-PD-1 inhibition in melanoma patients (from Gopalakrishnan et al, *Science* 2018, 359, 97-103).

DETAILED DESCRIPTION

The approach of the present invention is different from conventional antibiotic approaches. The present invention utilizes targeted cutting of microbial genomes using programmed nucleases, whereas conventional antibiotics rely upon metabolic processes and cell replication cycles—and the inhibition of these—for their activity. By focusing instead on nuclease cutting, the invention surprisingly achieves very quick and efficient microbial killing that also is remarkably durable. This is demonstrated in experiments below with different microbes, different nucleases and different delivery approaches. Typically, 99-100% killing was surprisingly observed many times and killing of 3-4 logs was very quickly achieved and with lasting duration.

The invention provides methods for treating or preventing microbial (eg, bacterial) infections and means for performing these methods. In particular, treatment of infections requiring rapid therapy is made possible, such as for treating acute conditions such as septicemia, sepsis, SIRS or septic shock. As explained herein, a rapid response is vital to address microbial infection in many settings. Speed is of the essence for many infection scenarios, such as acute infections requiring hospital admission. Benefits of the invention can be one or more of: the reduction in the spread, severity or progression of the infection in the subject; reduction in the

development, severity or progression of symptoms of the infection (eg, sepsis or septic shock); and an increase in the likelihood of survival in human or animal patients.

The invention uses programmable nuclease cutting of microbe genomes. The targeted cutting provides selective microbe killing or reduction of growth or proliferation to treat or prevent infection, as opposed to more broad-spectrum microbial killing of several different species as seen with conventional antibiotics. Selective killing is advantageous to leave beneficial microbes untargeted by the treatment, which may be beneficial to the patient. Moreover, the inventors have surprisingly found a substantial (by several logs) killing can be achieved very rapidly (eg, within 15 minutes) and sustainable effects can be achieved (eg, for more than 1 hour) in some embodiments. As exemplified below, the inventors surprisingly could remarkably achieve a fast and durable killing for around 2-3 hours.

Thus, the invention provides the following aspects:—

A programmable nuclease for use in a method of treating a microbial infection (eg, an acute bacterial infection) of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

Another aspect provides: A programmable nuclease for use in a method of rapidly treating an acute microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is rapidly treated.

Another aspect provides: A programmable nuclease for use in a method of treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a nucleic acid that programs the nuclease to recognise and cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

Another aspect provides: A programmable nuclease for use in a method of rapidly treating an acute microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a nucleic acid that programs the nuclease to recognise and cut the target site, whereby

genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is rapidly treated.

Another aspect provides: A programmable nuclease for use in a method of durably treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are durably killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

Another aspect provides: A programmable nuclease for use in a method of durably treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are durably killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a nucleic acid that programs the nuclease to recognise and cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

Another aspect provides: A programmable nuclease for use in a method of durably treating an acute microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are durably killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a nucleic acid that programs the nuclease to recognise and cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated.

Surprisingly, as exemplified below, a durable effect of several logs (eg, 3 or 4 logs) using a nuclease (as opposed to conventional means for conventional antibiotic killing) was observed around 3 hours after the first exposure of bacteria with a programmed nuclease. This aspect of the invention, therefore, makes possible dosing regimens for less frequent exposure to a programmed nuclease (ie, less frequent administration of a programmed nuclease, programmable nuclease and/or nucleic acid for programming the nuclease). For example, a Cas and gRNA (or DNA encoding a gRNA) for programming the nuclease are administered with a programmable nuclease (eg, a Cas 9 or Cas3) to the subject at a first time (T1) and at a second time (T2); or gRNA (or DNA encoding a gRNA) is administered on T1 and T2 for programming an endogenous Cas nuclease (eg, a Cas9 or Cas3) of bacteria of said first species or strain, wherein the programmed endogenous Cas cuts the genomes of the bacteria to kill the bacteria or to reduce growth or proliferation, thus treating the infection. Such less frequent dosing is convenient for the healthcare practitioner and patient, as well as provides for economical therapy. Thus, optionally, the nuclease and/or nucleic acid is administered to the subject on T1 and T2, wherein T2 is at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 24 hours after T1. For example, T2 is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 24

hours after T1. For example, T2 is 2-7 hours after T1. For example, T2 is 1 hour after T1. For example, T2 is 2 hours after T1. For example, T2 is 3 hours after T1. For example, T2 is 4 hours after T1. For example, T2 is 5 hours after T1.

Optionally, the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject on T1 and T2, wherein T2 is at least 1 hour (eg, 1, 1.5, 2, 2.5 or 3 hours) after T1.

Another aspect provides: A Cas nuclease for use in a method of treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable with a guide RNA (gRNA) to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising administering to the subject said a nucleic acid, wherein the nucleic acid is the gRNA or a DNA encoding the gRNA, thereby programming the nuclease to recognise and cut the target site of the microbes comprised by the subject, whereby genomes of the microbes are cut and microbial infection of the subject is treated, wherein the method comprises administering the nucleic acid to the subject on at a first time (T1) and at a second time (T2), whereby the subject is exposed to programmed nuclease on T1 and T2, and wherein T2 is no less than 1 hour after T1.

Optionally, T2 is no less than 2 hours after T1; optionally, T2 is no less than 3 hours after T1; optionally, T2 is no less than 4 hours after T1; optionally, T2 is no less than 5 hours after T1; optionally, T2 is no less than 6 hours after T1; optionally, T2 is no less than 7 hours after T1; optionally, T2 is no less than 8 hours after T1; optionally, T2 is no less than 9 hours after T1; optionally, T2 is no less than 10 hours after T1; optionally, T2 is no less than 11 hours after T1; optionally, T2 is no less than 12 hours after T1; optionally, T2 is no less than 13 hours after T1; optionally, T2 is no less than 14 hours after T1; or optionally, T2 is no less than 24 hours after T1. Additionally or alternatively: Optionally, T2 is no more than 7 hours after T1; optionally, T2 is no more than 12 hours after T1; optionally, T2 is no more than 24 hours after T1; optionally, T2 is 2-7 hours after T1; optionally, T2 is 24 hours after T1; optionally, T2 is 7 hours after T1; optionally, T2 is 6 hours after T1; optionally, T2 is 5 hours after T1; optionally, T2 is 4 hours after T1; optionally, T2 is 3 hours after T1; optionally, T2 is 2 hours after T1; optionally, T2 is 1 hour after T1. For example, T2 is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 24 hours after T1. For example, T2 is 1-7 hours after T1; or T2 is 2-7 hours after T1; or T2 is 3-7 hours after T1; or T2 is 4-7 hours after T1; or T2 is 5-7 hours after T1; or T2 is 6-7 hours after T1.

Optionally, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the method comprises reducing the infection at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the method comprises reducing the infection at least 10000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally, the method comprises reducing the infection such that the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection such that a reduction in infection by at least 100-fold persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the

infection such that a reduction in infection by at least 1000-fold persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection such that a reduction in infection by at least 10000-fold persists for 30 minutes immediately after the first 30 minutes of the treatment.

Optionally, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein a reduction in infection by at least 100-fold persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein a reduction in infection by at least 1000-fold persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection at least 10000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein a reduction in infection by at least 10000-fold persists for 30 minutes immediately after the first 30 minutes of the treatment.

Optionally, the method comprises maintaining reduction of the infection by at least 100-fold for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, a reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, the method comprises maintaining reduction of the infection by at least 1000-fold for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, a reduction of the infection by at least 1000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, the method comprises maintaining reduction of the infection by at least 10000-fold for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, a reduction of the infection by at least 10000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.

Optionally, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, the method comprises reducing the infection at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 1000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, the method comprises reducing the infection at least 10000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 10000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.

Optionally, the method comprises reducing the infection at least 10000-fold by the first 15 minutes of the treatment; and wherein reduction of the infection by at least 10000-fold is maintained for at least 45 minutes after exposing the subject to the programmed nuclease. This is exemplified below.

In an example, the infection is durably treated, wherein a reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after commencement of the treatment. In an example, the infection is durably treated, wherein a reduction of the infection by at least 1000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after commencement of the treatment. In an example, the infection is durably treated, wherein a reduction of the infection by at least 10000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after commencement of the treatment.

Optionally, the infection is reduced at least 100,000-fold by the first 30 or 45 minutes of the treatment. Optionally, the infection is reduced at least 100,000-fold by the first 30 or 45 minutes of the treatment and the reduction is maintained until the 60th minute of the treatment.

Optionally, the infection is reduced at least 1000,000-fold by the first 30 or 45 minutes of the treatment. Optionally, the infection is reduced at least 1000,000-fold by the first 30 or 45 minutes of the treatment and the reduction is maintained until the 60th minute of the treatment.

Optionally, the infection is reduced at least 100-fold by the first 15 minutes of the treatment. Optionally, the infection is reduced at least 1000-fold by the first 15 minutes of the treatment. Optionally, the infection is reduced at least 100-fold by the first 15 minutes of the treatment and at least 1000-fold by the first 30 minutes of the treatment.

For example, the reduction is maintained for at least 15 further minutes, eg, the infection is reduced at least 100-fold or at least 1000-fold by the first 15 minutes of the treatment and the reduction is maintained from the 15-30th minute or 15-45th minute of the treatment or 15-60th minute of the treatment.

For example, the infection is reduced at least 100-fold or at least 1000-fold or at least 10000-fold by the first 15 minutes of the treatment in the first 15 minutes and the reduction is maintained for from the 15-30th minute or 15-45th minute of the treatment.

Optionally, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally the method comprises reducing the infection at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally the method comprises reducing the infection at least 10000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally, the method comprises reducing the infection such that the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment, eg, the reduction may persist for at least 60 minutes after the first 30 minutes of the treatment. If the treatment is administered at time zero (T₀), then the reduction in infection may be present at 60 minutes counted after T₀, and indeed may persist after that 60 minutes. In FIGS. 1A, 5A and 6A, for example, reduction is seen at 60-180 minutes after T₀. Optionally, the reduction in infection persists for at least 30 minutes after the first 30 minutes of the treatment.

In an example, the infection is reduced by at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99%, eg, in the first 15 minutes of treatment. In an example, the infection is reduced by at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99%, eg, in the first 30 minutes of treatment.

For determining killing or reduction in growth or proliferation of the target microbes, one can, for example, deter-

mine the difference in the number of microbes of the first species or strain in (i) a sample taken from the subject (eg, a blood, gut or leaf sample) immediately before commencement of the treatment and (ii) a sample (of the same type as the sample of (i), eg, a blood, gut or leaf sample respectively) taken from the subject at 30 minutes of the treatment. For example, if the microbes are bacteria, the samples may be assessed for the difference in colony forming units (CFU)/ml sample, eg, when the samples have been plated on agar in respective petri dishes and incubated under identical conditions. Another example may use microscopic counting of microbes in samples, or other routine methods known to the skilled addressee.

In an example, at least 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% killing of the microbes is achieved by the first 30, 60, 90 or 120 minutes (eg, by the first 30 minutes; or by the first 120 minutes) of the treatment. For example, wherein the subject is a human or animal, the killing is determined comparing the prevalence (eg, by standard colony counting on an agar plate) of the microbes (eg, bacteria) in a blood sample taken immediately before commencement of the treatment versus a sample taken after the first 15 or 30 minutes of the treatment. In an example, at least 80, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% killing of the microbes is achieved by the first 0.5, 1 or 2 hours of the treatment. In an example, at least 99% killing of the microbes is achieved by the first 30 minutes of the treatment. In an example, at least 99% killing of the microbes is achieved by the first 2 hours of the treatment. In an example, 100% killing is achieved. These are exemplified below. In an embodiment, less than 100% of the microbes are killed.

Worked examples of killing in bacteria are shown below. Surprisingly, using a programmed nuclease to target the bacteria of choice, specific cutting resulted in rapid killing of the target bacteria—at least 3 or 4 logs of killing (ie, 1000- or 10,000 fold killing) could be observed in very short spaces of time and surprisingly these were sustained for at least to an hour. Optionally, the infection is reduced at least 1000-fold by the first 15, 30 or 45 minutes of the treatment. Optionally, the infection is reduced at least 1000-fold by the first 15, 30 or 45 minutes of the treatment and the reduction is maintained until the 60th, 120th or 180th minute of the treatment. Optionally, the infection is reduced at least 10,000-fold by the first 15, 30 or 45 minutes of the treatment. Optionally, the infection is reduced at least 10,000-fold by the first 15, 30 or 45 minutes of the treatment and the reduction is maintained until the 60th, 120th or 180th minute of the treatment. See, for example, exemplification in FIG. 5A.

In an example, 100% killing is achieved by 24 hours after commencement of the treatment.

In an example, the infection is reduced at least 1000-fold for 2 hours or more (eg, for 2-3 hours). Optionally also the infection is reduced by at least 1000-fold by the first 15 or 13 minutes of the treatment.

In an example, the infection is reduced at least 10,000-fold for 2 hours or more (eg, for 2-3 hours). Optionally also the infection is reduced by at least 10,000-fold by the first 15 or 13 minutes of the treatment.

In an example, the infection is reduced by at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% for 1 hour; or for 1 hour or more; or for 2 hours or more (eg, for 2-3 hours). Optionally, the infection is reduced by at least 90% for 1 hour; or for 1 hour or more; or for 2 hours or more (eg, for 2-3 hours), and optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the infection is reduced by at least 90% for 1 hour or more, and by the first

30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the infection is reduced by least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% by the first 15 or 13 minutes of the treatment. Optionally, the infection is reduced by least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% by the first 15 or 13 minutes of the treatment; and wherein the reduction is maintained for 1 hour or more (eg, for 2 hours or more; or for 3 hours or more; or for about 2 hours; or for 2 hours; or for about 3 hours; or for 3 hours). Exemplification below is provided, wherein the bacteria are *E. coli*.

Optionally, the subject is a human or animal and the microbes are bacteria (eg, *E. coli* or *C. difficile*), wherein blood infection of the subject by the bacteria is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the subject is a human or animal and the microbes are bacteria (eg, *E. coli* or *C. difficile*), wherein blood infection of the subject by the bacteria is reduced at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the subject is a human or animal and the microbes are bacteria (eg, *E. coli* or *C. difficile*), wherein blood infection of the subject by the bacteria is reduced at least 10,000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the *E. coli* are EHEC *E. coli*.

Optionally, the programmed nuclease (eg, a Cas9 or Cas3) is capable of cutting a target site comprised by *E. coli* (EHEC) ATCC43888. Optionally, the programmed nuclease (eg, a Cas9 or Cas3) is capable of cutting a target site comprised by *E. coli* Nissle.

Optionally, the blood of the subject is infected with from 10⁷ to 10¹² CFU/ml (eg, from 10¹⁷ to 10¹¹, from 10⁷ to 10¹⁰, from 10⁷ to 10⁹ or from 10⁷ to 10⁸ CFU/ml) of the bacteria immediately before the treatment.

The worked example below shows improved survival using the method of the invention in an in vivo model. In an example, therefore, the method of the invention is for improving survival of the subject by treating acute microbial infection of a subject. In an example, the programmed nuclease herein is capable of carrying out the killing of bacteria of the first species or strain in *Galleria mellonella* larvae in vivo infection model.

The nuclease may be, for example, a DNase (eg, a Cpf1, Cas9 or Cas3) or a RNase (eg, Cas13b).

In an example, the nuclease is an isolated or recombinant nuclease. For example, the nuclease is a synthetic or non-naturally occurring nuclease.

In an example, the nuclease is ex vivo, eg, in vitro. In an example, the nucleic acid is ex vivo. In an example, the guide RNA or DNA encoding guide RNA(s) herein is ex vivo, eg, in vitro.

Optionally, the nuclease is a Cas nuclease (eg, a Cpf1, CasX, CasY, Cas13b, Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease. In an example, the Cas is a *Streptococcus* (eg, *pyogenes* or *aureus*) Cas9, *Clostridium* (eg, *difficile*), *Salmonella* (eg, *typhimurium*) or *E. coli* Cas3. For example, the Cas is a spCas. In an example, the Cas9 is in combination with a tracrRNA or a DNA encoding a tracrRNA which is operable with the Cas. For example, the tracrRNA is of the same species as the Cas, eg, a *S. pyogenes* tracrRNA or DNA encoding this.

In an example, the nuclease is a Cas 3 encoded by a nucleic acid comprising SEQ ID NO: 9 or a sequence that is at least 80, 85, 90, 95, 96, 97, 98 or 99% identical thereto. Optionally, also the bacteria are *Clostridium* (eg, *C. difficile*) bacteria; or any *Clostridium* shown in Table 1. This is exemplified below.

In an example, the nuclease is a Cas 9 encoded by a nucleic acid comprising SEQ ID NO: 10 or a sequence that is at least 80, 85, 90, 95, 96, 97, 98 or 99% identical thereto. In an example, the nuclease is a Cas 9 comprising SEQ ID NO: 11 or a sequence that is at least 80, 85, 90, 95, 96, 97, 98 or 99% identical thereto. Optionally, also the bacteria are *Clostridium* (eg, *C. difficile*) bacteria; or any *Clostridium* shown in Table 1. Optionally, also the bacteria are *E. coli* (eg, EHEC). This is exemplified below.

Optionally, the method comprises administering to the subject a RNA or a nucleic acid (eg, DNA) that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

Optionally, the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject.

Optionally, subject comprises the nuclease prior to administration of the RNA or nucleic acid to the subject. For example, the nuclease is a Cas nuclease that is an endogenous Cas nuclease of bacterial cells of the first species or strain that are comprised by the subject. Thus, in this example, the RNA or nucleic acid may be administered to the subject and introduced into the bacteria for programming endogenous Cas comprised by the bacteria, thereby forming programmed Cas nuclease that cuts the target site in the genomes of the bacteria, whereby bacteria are killed or growth or proliferation of bacteria is reduced, thus treating or preventing the infection.

Optionally, a plurality of viruses (eg, phage or phagemids) are administered to the subject, wherein each virus comprises a copy (eg, one or more, eg, a plurality of copies) of the nucleic acid, wherein the viruses infect the microbes comprised by the subject to deliver thereto the nucleic acid. For example, viruses herein are phage or phagemids that infect (or are capable of infecting) the bacteria of the first species or strain.

Optionally, the ratio of administered viruses:microbes comprised by the subject is from 10 to 150. For example, the microbes are bacteria and the ratio is from 10 to 100, ie, a multiplicity of infection (MOI) of from 1 to 100 (eg, wherein the viruses are capable of replication, eg, are phage and not phagemid), eg, from 10 to 100. The ratio can be determined, for example, using a sample (eg, a blood or gut sample) from a human or animal subject immediately before the treatment and determining the number of microbes (eg, bacteria per ml of blood or gut sample). The amount of viruses to be administered can then be worked out according to the determination using the sample.

Optionally, the microbes are bacteria. Alternatively, the microbes are archaea. Alternatively, the microbes are viruses. Alternatively, the microbes are fungi. Alternatively, the microbes are algae. Alternatively, the microbes are protozoa.

In an example, the subject is a human and the infection is a nosocomial infection. In an example, the subject is a plant, yeast, protist or amoeba.

Optionally, the subject is a human (eg, an adult, child, neonate, toddler, teenager, male or female) or animal (eg, a dog, cat, horse, cow, sheep, goat, salmon, chicken, turkey, pig, companion animal or livestock animal).

In an example, the subject is a human or animal and: Optionally, the infection is an infection of the lungs, abdomen or urinary tract. In an example, the subject is suffering from a urinary tract infection, lung infections, such as pneumonia, a kidney infection or an abdominal infection. In an example, the subject is a surgery patient. In an example, the subject is a burns patient. In an example, the subject has

an infected wound (eg, a bacterially infected wound). In an example, the patient is suffering from AIDS or is infected by HIV. In an example, the subject is suffering from a cancer, such as a blood cancer, eg, leukaemia, eg, AML or CML or CLL or a lymphoma. In an example, the subject is a tissue or organ transplant patient, eg, a haematopoietic stem cell transplant or bone marrow transplant patient. In an example, the subject has a urinary or intravenous catheter. In an example, the subject is on mechanical ventilation. In an example, the subject has been receiving an immunosuppressant. In an example, the subject is suffering from pneumonia. In an example, the subject is an intensive care unit (ICU) patient. In an example, the subject is an Acute respiratory distress syndrome (ARDS) patient. In an example, the subject is suffering from meningitis, an infection in pregnancy, a ruptured gallbladder (a gallbladder rupture is a medical condition where the gallbladder leaks or bursts. Ruptures are commonly caused by inflammation of the gallbladder), abortion with septic shock (abortion with septic shock can be an acute life-threatening illness), endometritis (endometritis is an inflammatory condition of the lining of the uterus, usually due to an infection), Acute Respiratory Distress Syndrome (Acute respiratory distress syndrome is a lung condition; it occurs when fluid fills up the air sacs in the lungs) or cellulitis.

The increase in average age of the population, more people with chronic diseases, on immunosuppressive drugs, and increase in the number of invasive procedures being performed has led to an increased rate of sepsis. Optionally, the subject has undergone surgery, is on an immunosuppressant medication and/or is suffering from a chronic disease.

Optionally, the subject is a human over 60, 65, 70, 75 or 80 years of age or is a paediatric patient. In an alternative, the subject is a paediatric patient (eg, a human baby or child) or adolescent. In an example, the method treats or prevents neonatal sepsis in the subject. In an example the subject is an immune-compromised human or animal, eg, suffering from an acute viral infection, such as HIV infection; or the subject is suffering from a cancer, eg, a blood cancer, such as a leukaemia; or the patient is a transplant patient, eg, that has received an organ, tissue or bone marrow transplant. In an example, the subject is a human or animal that is positive for gram negative bacterial lipopolysaccharide or lipid A. In an example, the subject is a human or animal that is positive for gram positive bacterial cell wall lipoteichoic acid.

Optionally, the method treats or prevents septicaemia and/or sepsis (eg, septic shock) in the subject.

SIRS (Systemic Inflammatory Response Syndrome) criteria has been used to define sepsis.

SIRS is the presence of two or more of the following: abnormal body temperature, heart rate, respiratory rate, or blood gas, and white blood cell count. Sepsis is, for example, SIRS in response to an infectious process. Severe sepsis is, for example, sepsis with sepsis-induced organ dysfunction or tissue hypoperfusion (manifesting as hypotension, elevated lactate, or decreased urine output). Septic shock is, for example, severe sepsis plus persistently low blood pressure, despite the administration of intravenous fluids.

In an embodiment, the method prevents or delays progression of end-organ dysfunction in the subject (when the subject is a human or animal).

Examples of end-organ dysfunction include the following:

Lungs: acute respiratory distress syndrome (ARDS) ($\text{PaO}_2/\text{FiO}_2 < 300$)

Brain: encephalopathy symptoms including agitation, confusion, coma; causes may include ischemia, bleed-

ing, formation of blood clots in small blood vessels, microabscesses, multifocal necrotizing leukoencephalopathy

Liver: disruption of protein synthetic function manifests acutely as progressive disruption of blood clotting due to an inability to synthesize clotting factors and disruption of metabolic functions leads to impaired bilirubin metabolism, resulting in elevated unconjugated serum bilirubin levels

Kidney: low urine output or no urine output, electrolyte abnormalities, or volume overload

Heart: systolic and diastolic heart failure, likely due to chemical signals that depress myocyte function, cellular damage, manifest as a troponin leak (although not necessarily ischemic in nature)

More specific definitions of end-organ dysfunction exist for SIRS in pediatrics.

Cardiovascular dysfunction (after fluid resuscitation with at least 40 ml/kg of crystalloid)

hypotension with blood pressure <5th percentile for age or systolic blood pressure <2 standard deviations below normal for age, or

vasopressor requirement, or

two of the following criteria:

unexplained metabolic acidosis with base deficit >5 mEq/l

lactic acidosis: serum lactate 2 times the upper limit of normal

oliguria (urine output <0.5 ml/kg/h)

prolonged capillary refill >5 seconds

core to peripheral temperature difference >3° C.

Respiratory dysfunction (in the absence of cyanotic heart disease or known chronic lung disease)

the ratio of the arterial partial-pressure of oxygen to the fraction of oxygen in the gases inspired (PaO₂/FiO₂) <300 (the definition of acute lung injury), or

arterial partial-pressure of carbon dioxide (PaCO₂) >65 torr (20 mmHg) over baseline PaCO₂ (evidence of hypercapnic respiratory failure), or

supplemental oxygen requirement of greater than FiO₂ 0.5 to maintain oxygen saturation ≥92%

Neurologic dysfunction

Glasgow Coma Score (GCS) ≤11, or

altered mental status with drop in GCS of 3 or more points in a person with developmental delay/intellectual disability

Hematologic dysfunction

platelet count <80,000/mm³ or 50% drop from maximum in chronically thrombocytopenic, or

international normalized ratio (INR) >2

Disseminated intravascular coagulation

Kidney dysfunction

serum creatinine ≥2 times the upper limit of normal for age or 2-fold increase in baseline creatinine in people with chronic kidney disease

Liver dysfunction (only applicable to infants >1 month)

total serum bilirubin ≥4 mg/dl, or

alanine aminotransferase (ALT) ≥2 times the upper limit of normal

Table 2 sets out the criteria for a positive diagnosis of sepsis.

Optionally, the method reduces one or more symptoms in the patient selected from fever, low body temperature, rapid breathing, elevated heart rate, confusion, confusion, metabolic acidosis, respiratory alkalosis, low blood pressure, dysfunction of blood coagulation (such as blood clotting in

one or more organs, or bruising) and oedema. Optionally, the method reduces septic shock. Optionally, the sepsis is severe sepsis.

Optionally, at the start of the treatment, the subject (eg, a human) has a temperature of <36° C. or >38° C.; a heart rate of >90/min, a respiratory rate of >20 breaths/min or PaCO₂ <4.3 kPa; and white blood cell count of <4000/mm³ or >12,000/mm³.

Optionally, at the start of the treatment, the subject (eg, a human) has presence of two or more of the following: abnormal body temperature, abnormal heart rate, abnormal respiratory rate, abnormal blood gas and abnormal white blood cell count.

Optionally, the subject is a plant. In an example, the subject is a protist, eg, amoeba. Optionally in this example, the microbes are viruses (eg, large or giant viruses, eg, Mimiviruses). The nuclease, for example, is a Cas and is programmable using a guide RNA delivered by a viroplasm that infects the virus microbes

In an example the microbes are yeast, eg, *Candida*.

Preferably, the microbes are bacteria. Optionally, the bacteria are gram positive bacteria. Optionally, the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonnorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E. coli* (eg, EHEC *E. coli*), *C. difficile*, *V. cholera*, *Staphylococcus* (eg, *S. aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).

Optionally, the first species is selected from the species in Table 1.

Optionally, the first species is enterohemorrhagic *E. coli* (EHEC), *E. coli* Serotype O157:H7 or Shiga-toxin producing *E. coli* (STEC). In an example, the bacteria are selected from

Shiga toxin-producing *E. coli* (STEC) (STEC may also be referred to as Verocytotoxin-producing *E. coli* (VTEC); Enterohemorrhagic *E. coli* (EHEC) (this pathotype is the one most commonly heard about in the news in association with foodborne outbreaks);

Enterotoxigenic *E. coli* (ETEC);
Enteropathogenic *E. coli* (EPEC);
Enterohemorrhagic *E. coli* (EHEC);
Enteroinvasive *E. coli* (EIEC); and
Diffusely adherent *E. coli* (DAEC).

Enterohemorrhagic *Escherichia coli* (EHEC) serotype O157:H7 is a human pathogen responsible for outbreaks of bloody diarrhoea and haemolytic uremic syndrome (HUS) worldwide. Conventional antimicrobials trigger an SOS response in EHEC that promotes the release of the potent Shiga toxin that is responsible for much of the morbidity and mortality associated with EHEC infection. Cattle are a natural reservoir of EHEC, and approximately 75% of EHEC outbreaks are linked to the consumption of contaminated bovine-derived products. EHEC causes disease in humans but is asymptomatic in adult ruminants. Characteristics of *E. coli* serotype O157:H7 (EHEC) infection includes abdominal cramps and bloody diarrhoea, as well as the life-threatening complication haemolytic uremic syndrome (HUS). Currently there is a need for a treatment for EHEC infections (Goldwater and Bettelheim, 2012). The use of conventional antibiotics exacerbates Shiga toxin-mediated cytotoxicity. In an epidemiology study conducted by the Centers for Disease Control and Prevention, patients treated with antibiotics for EHEC enteritis had a higher risk of developing HUS (Slutsker et al., 1998). Additional studies support the contraindication of antibiotics in EHEC

infection; children on antibiotic therapy for hemorrhagic colitis associated with EHEC had an increased chance of developing HUS (Wong et al., 2000; Zimmerhackl, 2000; Safdar et al., 2002; Tarr et al., 2005). Conventional antibiotics promote Shiga toxin production by enhancing the replication and expression of six genes that are encoded within a chromosomally integrated lambdoid prophage genome. The approach of the present invention relies on nuclease cutting. Stx induction also promotes phage-mediated lysis of the EHEC cell envelope, allowing for the release and dissemination of Shiga toxin into the environment (Karch et al., 1999; Matsushiro et al., 1999; Wagner et al., 2002). Thus, advantageously, the invention provides alternative means for treating EHEC in human and animal subjects. This is exemplified below with surprising results on the speed and duration of anti-EHEC action produced by nuclease action (as opposed to conventional antibiotic action).

In an example, the subject (eg, a human) is suffering from or at risk of haemolytic uremic syndrome (HUS), eg, the subject is suffering from an *E coli* infection, such as an EHEC *E coli* infection.

An aspect of the invention provides: A plurality of viruses (eg, phage or phagemids for producing phage) for use with the nuclease of the invention in the method of treatment, wherein each virus comprises a copy of a nucleic acid described herein, wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

A aspect of the invention provides: A plurality of viruses (eg, phage or phagemids for producing phage) for use with a programmable nuclease in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated;

wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

Optionally, the method is for durable treatment, eg, as described herein; and/or optionally, the infection is acute infection.

Optionally, the method is for rapid treatment, eg, as described herein; and/or optionally, the infection is acute infection.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, the nucleic acid is according to any nucleic acid of the invention herein.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, the nucleic acid is according to any nucleic acid of the invention herein.

In an alternative, when the microbes are viruses, the plurality of viruses are phage that are capable of infecting host cells harbouring the microbes, wherein the nucleic acids are introduced into the host cells for expression therein of the RNA. The RNA complexes with the nuclease in the

host cells to guide the nuclease to cut a target site of the microbes (ie, to cut viral RNA or DNA), thereby inactivating the viral microbes. For example, the microbes are viruses (eg, in amoeba; or in human or animal or plant cells) and the viruses of said plurality of viruses are capable of targeting the microbes, whereby the nuclease is programmed to cut the microbes (eg, in the amoeba or in said cells).

An aspect of the invention provides: A composition comprising a plurality of nucleic acids for programming the nuclease of the invention in the method of treatment, wherein each nucleic acid is a nucleic acid as defined herein.

An aspect of the invention provides: A composition comprising a plurality of nucleic acids for programming a programmable nuclease in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and the nucleic acids wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated; wherein each nucleic acid encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

Optionally, the method is for durable treatment, eg, as described herein; and/or optionally, the infection is acute infection.

Optionally, the method is for rapid treatment, eg, as described herein; and/or optionally, the infection is acute infection.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, each nucleic acid is according to any nucleic acid of the invention herein.

Optionally, the composition is a pharmaceutical composition comprising the nucleic acids and a pharmaceutically acceptable diluent, carrier or excipient. Optionally, the composition is for oral, intravenous, pulmonary, rectal, topical, buccal, ocular, intranasal, or subcutaneous administration to a human or animal subject. Optionally, the composition is a herbicide or pesticide or insecticide or nematocide or arnicide. Optionally, the composition is toxic to yeast. Optionally, the composition is toxic to giant viruses.

An aspect of the invention provides: A CRISPR/Cas system comprising a nuclease according to the invention for use in the method of treatment, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9 or any other Cas mentioned herein) and the system comprises one or more guide RNAs or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of the microbes.

In an example, each guide RNA mentioned herein is a single guide RNA (ie, a chimaeric guide RNA). In another example, each guide RNA comprises a crRNA that is hybridised to a tracrRNA.

In an example, a target site mentioned herein is comprised by an essential gene, virulence gene or antibiotic resistance gene of the bacteria. In an example, a target site mentioned herein is comprised by a multi-copy sequence (ie, a sequence that is present in more than one (eg, 2, 3, 4, 5, 6, 7, 8 or 9, or more) copies in each bacterial genome). For example, the target site is comprised by a ribosomal RNA gene. In an example, a target site mentioned herein is

comprised by a ribosomal RNA gene (eg, a 23S ribosomal RNA gene), a yapH gene; or a pks gene; or homologue or orthologue thereof.

Optionally, each guide RNA herein is capable of hybridizing to a protospacer sequence comprising the target site, wherein the protospacer sequence is 15-45 nucleotides in length, eg, 15-25; 18-21; 20; or about 20 nucleotides in length. Optionally, each guide RNA herein comprises a spacer sequence that is 15-45 nucleotides in length, eg, 15-25; 18-21; 20; or about 20 nucleotides in length.

Optionally, each guide RNA herein is cognate to a 5'-NGG protospacer adjacent motif (PAM), eg, wherein the bacteria are *E. coli*. Optionally, each guide RNA herein is cognate to a 5'-CCA or 5'-CCT protospacer adjacent motif (PAM), eg, wherein the bacteria are *C. difficile*.

An aspect of the invention provides: A guide RNA or a DNA encoding a guide RNA for use in the system of the invention for use in the method of treating an acute microbial infection in the subject, eg, septicaemia or sepsis.

An aspect of the invention provides: A nucleic acid vector comprising the guide RNA or DNA.

Optionally, the vector is a phage, phagemid, viriophage, virus, plasmid (eg, conjugative plasmid) or transposon. The example below shows almost complete killing can be achieved using a conjugative plasmid as the vector. Thus, in an embodiment, each vector is a conjugative plasmid that is delivered from carrier bacteria eg, probiotic carrier bacteria for administration to the human or animal subject. In an example, the carrier bacteria are *Lactobacillus* (eg, *L. reuteri*) or *E. coli*. This is exemplified below and achieved complete (100%) killing.

An aspect of the invention provides: An anti-sepsis or anti-septicaemia composition for administration to a human or animal for treating sepsis or septicaemia, the composition comprising a plurality of vectors, wherein each vector a vector of the invention.

An aspect of the invention provides: A method of treating (eg, rapidly and/or durably treating) an acute microbial infection of a subject, wherein the method is as defined herein.

An aspect of the invention provides: A method of treating (eg, rapidly and/or durably treating) an acute microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated (eg, rapidly and/or durably treated).

An aspect of the invention provides: A method of treating (eg, rapidly and/or durably treating) an acute microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a plurality of viruses wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated (eg, rapidly and/or durably treated); wherein each virus comprises a copy of a

nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, the nucleic acid is according to any nucleic acid of the invention herein.

An aspect of the invention provides: A method of treating (eg, rapidly and/or durably treating) an acute microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a plurality of nucleic acids wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated (eg, rapidly and/or durably treated); wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein each nucleic acid encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, each nucleic acid is according to any nucleic acid of the invention herein.

In an example, the invention is for medical or dental or ophthalmic use (eg, for treating or preventing an infection in an organism or limiting spread of the infection in an organism).

In an example, the invention is for cosmetic use (eg, use in a cosmetic product, eg, make-up), or for hygiene use (eg, use in a hygiene product, eg, soap).

In an example, the vectors and/or nuclease prior to administration to the subject are comprised by a composition which is as any of the following (host here refers to the microbes of the first species or strain): In an example, the composition is a medical, ophthalmic, dental or pharmaceutical composition (eg, comprised by an anti-host vaccine). In an example, the composition is an antimicrobial composition, eg, an antibiotic or antiviral, eg, a medicine, disinfectant or mouthwash. In an example, the composition is a cosmetic composition (eg, face or body make-up composition). In an example, the composition is a herbicide. In an example, the composition is a pesticide (eg, when the host is a *Bacillus* (eg, *thuringiensis*) host). In an example, the composition is a beverage (eg, beer, wine or alcoholic beverage) additive. In an example, the composition is a food additive (eg, where the host is an *E. coli*, *Salmonella*, *Listeria* or *Clostridium* (eg, botulinum) host). In an example, the composition is a water additive. In an example, the composition is an additive for aquatic animal environments (eg, in a fish tank). In an example, the composition is an oil or petrochemical industry composition or comprised in such a composition (eg, when the host is a sulphate-reducing bacterium, eg, a *Desulfovibrio* host). In an example, the composition is an oil or petrochemical additive. In an example, the composition is a chemical additive. In an example, the composition is a disinfectant (eg, for sterilizing equipment

for human or animal use, eg, for surgical or medical use, or for baby feeding). In an example, the composition is a personal hygiene composition for human or animal use. In an example, the composition is a composition for environmental use, eg, for soil treatment or environmental decontamination (eg, from sewage, or from oil, a petrochemical or a chemical, eg, when the host is a sulphate-reducing bacterium, eg, a *Desulfovibrio* host). In an example, the composition is a plant growth stimulator. In an example, the composition is a composition for use in oil, petrochemical, metal or mineral extraction. In an example, the composition is a fabric treatment or additive. In an example, the composition is an animal hide, leather or suede treatment or additive. In an example, the composition is a dye additive. In an example, the composition is a beverage (eg, beer or wine) brewing or fermentation additive (eg, when the host is a *Lactobacillus* host). In an example, the composition is a paper additive. In an example, the composition is an ink additive. In an example, the composition is a glue additive. In an example, the composition is an anti-human or animal or plant parasitic composition. In an example, the composition is an air additive (eg, for air in or produced by air conditioning equipment, eg, where the host is a *Legionella* host). In an example, the composition is an anti-freeze additive (eg, where the host is a *Legionella* host). In an example, the composition is an eyewash or ophthalmic composition (eg, a contact lens fluid). In an example, the composition is comprised by a dairy food (eg, the composition is in or is a milk or milk product; eg, wherein the host is a *Lactobacillus*, *Streptococcus*, *Lactococcus* or *Listeria* host). In an example, the composition is or is comprised by a domestic or industrial cleaning product (eg, where the host is an *E coli*, *Salmonella*, *Listeria* or *Clostridium* (eg, botulinum) host). In an example, the composition is comprised by a fuel. In an example, the composition is comprised by a solvent (eg, other than water). In an example, the composition is a baking additive (eg, a food baking additive). In an example, the composition is a laboratory reagent (eg, for use in biotechnology or recombinant DNA or RNA technology). In an example, the composition is comprised by a fibre retting agent. In an example, the composition is for use in a vitamin synthesis process. In an example, the composition is an anti-crop or plant spoiling composition (eg, when the host is a saprotrophic bacterium). In an example, the composition is an anti-corrosion compound, eg, for preventing or reducing metal corrosion (eg, when the host is a sulphate-reducing bacterium, eg, a *Desulfovibrio* host, eg for use in reducing or preventing corrosion of oil extraction, treatment or containment equipment; metal extraction, treatment or containment equipment; or mineral extraction, treatment or containment equipment). In an example, the composition is an agricultural or farming composition or comprised in such a composition. In an example, the composition is a silage additive. The invention provides a CRISPR array, gRNA-encoding nucleotide sequence, vector or plurality of vectors described herein for use in any of the compositions described in this paragraph or for use in any application described in this paragraph, eg, wherein the host cell is a bacterial or archaeal cell. The invention provides a method for any application described in this paragraph, wherein the method comprises combining a CRISPR array, gRNA-encoding nucleotide sequence, vector or plurality of the invention with a host cell (eg, bacterial or archaeal cell). In an embodiment, the host cell is not present in or on a human (or human embryo) or animal.

Any aspect of the present invention is, for example, for an industrial or domestic use, or is used in a method for such

use. For example, it is for or used in agriculture, oil or petroleum industry, food or drink industry, clothing industry, packaging industry, electronics industry, computer industry, environmental industry, chemical industry, aerospace industry, automotive industry, biotechnology industry, medical industry, healthcare industry, dentistry industry, energy industry, consumer products industry, pharmaceutical industry, mining industry, cleaning industry, forestry industry, fishing industry, leisure industry, recycling industry, cosmetics industry, plastics industry, pulp or paper industry, textile industry, clothing industry, leather or suede or animal hide industry, tobacco industry or steel industry.

Host cells herein refers to the microbes of the first species or strain. Optionally, any host cell(s) herein is/are bacterial or archaeal cells. In an example, the cell(s) is/are in stationary phase. In an example, the cell(s) is/are in exponential phase. In an example, the cell(s) is/are in lag phase. In an example, the cell(s) is/are wild-type cells or naturally-occurring cells, eg, comprised by a naturally-occurring microbiome, eg, of a human, animal, plant, soil, water, sea, waterway or environment. In an example, the cell(s) is/are artificially genetically modified.

In an example, a plurality of vectors of the invention are introduced into a plurality of said host cells, wherein the host cells are comprised by a bacterial population, eg, ex vivo, in vivo or in vitro. In an example, the host cells are comprised by a microbiota population comprised by an organism or environment (eg, a waterway microbiota, water microbiota, human or animal gut microbiota, human or animal oral cavity microbiota, human or animal vaginal microbiota, human or animal skin or hair microbiota or human or animal armpit microbiota), the population comprising first bacteria that are symbiotic or commensal with the organism or environment and second bacteria comprising said host cells, wherein the host cells are detrimental (eg, pathogenic) to the organism or environment. In an embodiment, the population is ex vivo. In an example, the ratio of the first bacteria sub-population to the second bacteria sub-population is increased. In an example, the first bacteria are *Bacteroides* (eg, *B fragalis* and/or *B thetaiotaomicron*) bacteria. Optionally, the *Bacteroides* comprises one, two, three or more *Bacteroides* species selected from *caccae*, *capillosus*, *cellulosilyticus*, *coprocola*, *coprophilus*, *coprosuis*, *distasonis*, *dorei*, *eggerthii*, *faecis*, *fmegoldii*, *fluxus*, *fragalis*, *intestinalis*, *melaninogenicus*, *nordii*, *oleiciplenus*, *oxalis*, *ovatus*, *pectinophilus*, *plebeius*, *stercoris*, *thetaitaomicron*, *uniformis*, *vulgatus* and *xylanisolvens*. For example, the *Bacteroides* is or comprises *B thetaiotaomicron*. For example, the *Bacteroides* is or comprises *B fragalis*.

In an example, the host, first or second cells are any bacterial species disclosed in US20160333348, GB1609811.3, PCT/EP2017/063593 and all US equivalent applications. The disclosures of these species (including specifically, Table 1 of PCT/EP2017/063593), are incorporated herein in their entirety and for potential inclusion of one or more disclosures therein in one or more claims herein.

In an example, the host cell(s) or bacterial population is harboured by a beverage or water (eg, a waterway or drinking water) for human consumption. In an example, the host cell(s) or said population is comprised by a composition (eg, a medicament (eg, bacterial gut transplant), beverage, mouthwash or foodstuff) for administration to a human or non-human animal for populating and rebalancing the gut or oral microbiota thereof (eg, wherein said use of the medicament is to treat or prevent a disease or condition in the human or animal). In an example, the host cell(s) or said

population are on a solid surface or comprised by a biofilm (eg, a gut biofilm or a biofilm on an industrial apparatus). In an example of the invention for in vitro treating an industrial or medical fluid, solid surface, apparatus or container (eg, for food, consumer goods, cosmetics, personal healthcare product, petroleum or oil production); or for treating a waterway, water, a beverage, a foodstuff or a cosmetic, wherein the host cell(s) are comprised by or on the fluid, surface, apparatus, container, waterway, water, beverage, foodstuff or cosmetic.

In an example, the invention provides a container for medical or nutritional use, wherein the container comprises the vectors for use in the method. For example, the container is a sterilised container, eg, an inhaler or connected to a syringe or IV needle.

In an example, the vectors or composition is for administration (or is administered) to the human or non-human animal subject by mucosal, gut, oral, intranasal, intrarectal, intravaginal, ocular or buccal administration.

Optionally, each host cell is of a strain or species found in human microbiota, optionally wherein the host cells are mixed with cells of a different strain or species, wherein the different cells are *Enterobacteriaceae* or bacteria that are probiotic, commensal or symbiotic with humans (eg, in the human gut. In an example, the host cell is an *E coli* or *Salmonella* cell.

The invention is optionally for altering the relative ratio of sub-populations of first and second bacteria in a mixed population of bacteria, eg, for altering human or animal microbiomes, such as for the alteration of the proportion of Bacteroidetes (eg, *Bacteroides*, eg, *fragalis* and/or *thetaita-micron*), Firmicutes and/or gram positive or negative bacteria in microbiota of a human.

In an example, the vectors or composition of the invention comprises a nucleotide sequence for expressing in the host cell an endolysin for host cell lysis, optionally wherein the endolysin is a phage phi11, phage Twort, phage P68, phage phiWMY or phage K endolysin (eg, MV-L endolysin or P-27/HP endolysin).

In an example, the target site is comprised by a chromosome of each microbe host cell, eg, wherein the sequence is comprised by an antibiotic resistance gene, virulence gene or essential gene of the host cell. An example, provides the vectors of the invention in combination with an antibiotic agent (eg, a beta-lactam antibiotic), eg, wherein the vectors target a protospacer sequence comprised by an antibiotic resistance gene comprised by host cell genome or episome (eg, a plasmid comprised by the host cells). In an example, the episome is a plasmid, transposon, mobile genetic element or viral sequence (eg, phage or prophage sequence).

In an example, the target is a chromosomal sequence, an endogenous host cell sequence, a wild-type host cell sequence, a non-viral chromosomal host cell sequence, not an exogenous sequence and/or a non-phage sequence (ie, one more or all of these), eg, the sequence is a wild-type host chromosomal cell sequence such as antibiotic resistance gene or essential gene sequence comprised by a host cell chromosome. In an example, the sequence is a host cell plasmid sequence, eg, an antibiotic resistance gene sequence.

Optionally, the nuclease is a Cas and the target site is comprised by a protospacer sequence that is adjacent a NGG, NAG, NGA, NGC, NGGNG, NNGRRT or NNA-GAAW protospacer adjacent motif (PAM), eg, a AAAGAAA or TAAGAAA PAM (these sequences are written 5' to 3'). In an embodiment, the PAM is immediately adjacent the 3' end of the protospacer sequence. In an

example, the Cas is a *S aureus*, *S theromophilus* or *S pyogenes* Cas. In an example, the Cas is Cpf1 and/or the PAM is TTN or CTA. Optionally, the Cas is a Type I (eg, Type I-A, I-B, I-C, I-D, I-E, or I-F) CRISPR system Cas. Optionally, the Cas is a Type II CRISPR system Cas. Optionally, the Cas is a Type III CRISPR system Cas. Optionally, the Cas is a Type IV CRISPR system Cas. Optionally, the Cas is a Type V CRISPR system Cas. Optionally, the Cas is a Type VI CRISPR system Cas.

Optionally, the nuclease is a Cas and each vector comprises a cognate CRISPR array that comprises multiple copies of the same spacer for targeting the target site. Optionally, there is provide a vector or plurality of vectors of the invention, wherein the vector(s) comprises a plurality of CRISPR arrays of said gRNA-encoding sequences for host cell protospacer sequence targeting, wherein the protospacers comprise the target site. Optionally, the or each vector comprises two, three or more of copies of nucleic acid sequences encoding crRNAs (eg, gRNAs), wherein the copies comprise the same spacer sequence for targeting a host cell target site (eg, a site comprised by a virulence, resistance or essential gene sequence).

In an example, at least two target sequences are modified by Cas, for example an antibiotic resistance gene and an essential gene. Multiple targeting in this way may be useful to reduce evolution of escape mutant host cells.

In an example, the Cas is a wild-type endogenous host cell Cas nuclease. In an example, target site cutting is carried out by a dsDNA Cas nuclease (eg, a Cas9, eg, a spCas9 or saCas9), whereby repair of the cut is by non-homologous end joining (NHEJ); alternatively the Cas is an exonuclease or Cas3

In an example, the array, gRNA-encoding sequence or vector is not in combination with a Cas endonuclease-encoding sequence that is naturally found in a cell together with repeat sequences of the array or gRNA-encoding sequence.

A tracrRNA sequence may be omitted from an array or vector of the invention, for example for Cas systems of a Type that does not use tracrRNA, or an endogenous tracrRNA may be used with the crRNA encoded by the vector.

In an example, the host target site is comprised by at least 5, 6, 7, 8, 9, 10, 20, 30 or 40 contiguous nucleotides.

In an example, the or each vector comprises an exogenous promoter functional for transcription of the crRNA or gRNA in the microbes.

Optionally, each vector is a plasmid, cosmid, virus, a virion, phage, phagemid or prophage. For example, the invention provides a plurality of bacteriophage comprising a plurality of vectors of the invention, eg, wherein the vectors are identical. In an example, the vector is a viral vector. Viral vectors have a particularly limited capacity for exogenous DNA insertion, thus virus packaging capacity needs to be considered. Room needs to be left for sequences encoding vital viral functions, such as for expressing coat proteins and polymerase. In an example, the vector is a phage vector or an AAV or lentiviral vector. Phage vectors are useful where the host is a bacterial cell. In an example, the vector is a virus capable of infecting an archaea host cell.

Optionally, vector components are comprised by a transposon that is capable of transfer into and/or between host cells. The transposon can be a transposon as described in US20160333348, GB1609811.3 and all US equivalent applications; the disclosures of these, including these specific transposon disclosures, are incorporated herein in its entirety and for potential inclusion of one or more disclosures therein in one or more claims herein.

In an example, the or each vector is provided by a nanoparticle or in liposomes.

In an example, the or each host cell (or first and/or second bacteria) is a gram positive cell. In an example, the or each host cell is an *Enterobacteriaceae*, eg, *Salmonella*, *Yersinia pestis*, *Klebsiella*, *Shigella*, *Proteus*, *Enterobacter*, *Serratia*, or *Citrobacter* cells. Optionally, the or each cell is an *E coli* (eg, *E coli* K12) or *Salmonella* (eg, *S enteric* serovar *typhimurium*) cell. Optionally, the or each host cell (or first and/or second bacteria) is a gram negative cell.

Optionally, the host (or first and/or second bacteria) is a *mycoplasma*, *chlamydiae*, *spirochete* or *mycobacterium*. Optionally, the host (or first and/or second bacteria) is a *Streptococcus* (eg, *pyogenes* or *thermophilus*) host. Optionally, the host (or first and/or second bacteria) is a *Staphylococcus* (eg, *aureus*, eg, MRSA) host. Optionally, the host (or first and/or second bacteria) is an *E. coli* (eg, O157: H7) host. Optionally, the host (or first and/or second bacteria) is a *Pseudomonas* (eg, *aeruginosa*) host. Optionally, the host (or first and/or second bacteria) is a *Vibro* (eg, *cholerae* (eg, O139) or *vulnificus*) host. Optionally, the host (or first and/or second bacteria) is a *Neisseria* (eg, *gonorrhoeae* or *meningitidis*) host. Optionally, the host (or first and/or second bacteria) is a *Bordetella* (eg, *pertussis*) host. Optionally, the host (or first and/or second bacteria) is a *Haemophilus* (eg, *influenzae*) host. Optionally, the host (or first and/or second bacteria) is a *Shigella* (eg, *dysenteriae*) host. Optionally, the host (or first and/or second bacteria) is a *Brucella* (eg, *abortus*) host. Optionally, the host (or first and/or second bacteria) is a *Francisella* host. Optionally, the host (or first and/or second bacteria) is a *Xanthomonas* host. Optionally, the host (or first and/or second bacteria) is a *Agrobacterium* host. Optionally, the host (or first and/or second bacteria) is a *Erwinia* host. Optionally, the host (or first and/or second bacteria) is a *Legionella* (eg, *pneumophila*) host. Optionally, the host (or first and/or second bacteria) is a *Listeria* (eg, *monocytogenes*) host. Optionally, the host (or first and/or second bacteria) is a *Campylobacter* (eg, *jejuni*) host. Optionally, the host (or first and/or second bacteria) is a *Yersinia* (eg, *pestis*) host. Optionally, the host (or first and/or second bacteria) is a *Borelia* (eg, *burgdorferi*) host. Optionally, the host (or first and/or second bacteria) is a *Helicobacter* (eg, *pylori*) host. Optionally, the host (or first and/or second bacteria) is a *Clostridium* (eg, *dificile* or *botulinum*) host. Optionally, the host (or first and/or second bacteria) is a *Erlchia* (eg, *chaffeensis*) host. Optionally, the host (or first and/or second bacteria) is a *Salmonella* (eg, *typhi* or *enterica*, eg, serotype *typhimurium*, eg, DT 104) host. Optionally, the host (or first and/or second bacteria) is a *Chlamydia* (eg, *pneumoniae*) host. Optionally, the host (or first and/or second bacteria) is a *Parachlamydia* host. Optionally, the host (or first and/or second bacteria) is a *Corynebacterium* (eg, *amycolatum*) host. Optionally, the host (or first and/or second bacteria) is a *Klebsiella* (eg, *pneumoniae*) host. Optionally, the host (or first and/or second bacteria) is a *Enterococcus* (eg, *faecalis* or *faecim*, eg, linezolid-resistant) host. Optionally, the host (or first and/or second bacteria) is a *Acinetobacter* (eg, *baumannii*, eg, multiple drug resistant) host.

Optionally, the invention is for reducing the growth or proliferation of host cell(s) in an environment (eg, soil, a composition comprising said host cells and yeast cells), human, animal or plant microbiome. This is useful, for example, when the microbiome is naturally-occurring.

Optionally, the nuclease is a Cas and the target is comprised by a protospacer sequence comprising at least 5, 6, 7, 8, 9 or 10 contiguous nucleotides immediately 3' of a

cognate PAM in the genome of the host cell, wherein the PAM is selected from AWG, AAG, AGG, GAG and ATG. Non Medical, Ex Vivo & In Vitro Uses Etc

In certain configurations, the inventive observation of rapid and durable microbial killing and growth or proliferation inhibition using nuclease cutting finds application in subjects other than humans and animals (eg, to treat plants or yeast cultures), or for ex vivo or in vitro treatment of substrates, such as industrial surfaces, fluids and apparatus. Thus, the invention further provides the following Concepts. Any other feature herein of the invention, its configurations, aspects, embodiments, options and examples above and elsewhere herein are combinable mutatis mutandis with these Concepts (including for providing combinations of features in the claims herein).

A Concept provides: Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of the invention, in the manufacture of a composition for carrying out a method of treatment as defined herein, wherein the subject is an organism other than a human or animal.

A Concept provides: Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of the invention, in the manufacture of a composition for carrying out an ex vivo or in vitro a method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.

Herein, treatment of an infection of a substrate may mean the treatment of a bacterial population (eg, one or more colonies) on a surface of the substrate and/or incorporated in the material of the substrate. For example, the treatment may be the treatment to kill bacteria on the surface of an industrial apparatus or equipment (eg, medical equipment, such as a scalpel or medical device or tubing). In another example, the substrate is a fluid (eg, a liquid or a gas), such as a medical fluid or petroleum product in fluid form (eg, an oil or hydrocarbon fluid or liquid).

A Concept provides: Use of a programmable nuclease in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.

Optionally the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject or substrate at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1. T1 and T2 may be as defined herein.

Optionally, the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the infection is maintained by at least

100-fold for at least 60 minutes (eg, at least 120 minutes) after exposing the subject to the programmed nuclease.

Optionally, the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.

Optionally, the method comprises administering to the subject or substrate a RNA or a nucleic acid that encodes an RNA for expression of the RNA in or on the subject or substrate, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject or substrate.

Optionally, the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject or substrate.

Optionally, the subject or substrate comprises the nuclease prior to administration of the RNA or nucleic acid.

Optionally, a plurality of viruses (eg, phage) are administered to the subject or substrate, wherein each virus comprises a copy of the nucleic acid, wherein the viruses infect the microbes comprised by the subject or substrate to deliver thereto the nucleic acid.

Optionally, the ratio of administered viruses:microbes is from 10 to 150.

Optionally, the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally, the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject or substrate to the programmed nuclease.

Optionally, the subject is a plant; or wherein the substrate is a metallic, plastic, concrete, stone, wood, glass or ceramic substrate. Optionally, the subject is a fluid (eg, a liquid or a gas).

Optionally, the microbes are bacteria or archaea. Optionally, the bacteria are gram positive bacteria. Optionally, the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonnorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).

Optionally, the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.

Reference is made to WO2016177682, which discusses aspects of microbiologically influenced corrosion (MIC) or biofouling of substrates and discloses methods for controlling MIC or biofouling of a substrate. The methods, nucleases, arrays, RNAs, vectors and viruses disclosed in that document can be employed in the present invention, for example for carrying out the method or use of the present invention and the disclosures of these parts and the substrates and bacteria disclosed in WO2016177682 are incorporated herein by reference for potentially providing disclosure of features possible to be used in one or more claims herein.

Optionally, the use of the present invention is for controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate in an industrial or domestic system (eg, a system disclosed in WO2016177682, which disclosure is incorporated herein by reference). In an example, the system comprises equipment (eg, for use in an industrial process) and the surface is a surface of said equipment. In an

example, the biofouling comprises microbial biofilm and/or sludge formation, proliferation or maintenance. In an example, the microbes are sessile. In an example "controlling" comprises preventing, reducing or eliminating said MIC or biofouling, or reducing spread of said MIC or biofouling in the system. Cell growth or proliferation or maintenance is, for example, a characteristic of cell viability. Thus, in an example, the method reduces microbe proliferation and/or maintenance.

Optionally, the microbes are comprised by a microbial biofilm that is in contact with said substrate. Optionally, said surface and host cells are in contact with a fluid, such as an aqueous liquid (eg, sea water, fresh water, stored water or potable water).

Fresh water is naturally occurring water on the Earth's surface in ice sheets, ice caps, glaciers, icebergs, bogs, ponds, lakes, rivers and streams, and underground as groundwater in aquifers and underground streams. Fresh water is generally characterized by having low concentrations of dissolved salts and other total dissolved solids. The term specifically excludes sea water and brackish water, although it does include mineral-rich waters such as chalybeate springs. In an example said fresh water is any of these fresh water types. Potable water is water for human or animal (eg, livestock) consumption. In an example, the fluid is selected from industrial cooling water wherein the system is a cooling system; sewage water wherein the system is a sewage treatment or storage system; drinking water wherein the system is a drinking water processing, storage, transportation or delivery system; paper making water wherein the system is a paper manufacture or processing system; swimming pool water wherein the system is a swimming pool or swimming pool water treatment or storage system; fire extinguisher water wherein the system is a fire extinguishing system; or industrial process water in any pipe, tank, pit, pond or channel.

Optionally, the use is for controlling bacterial souring of a liquid in a reservoir or container), wherein the fluid comprises a population of first host cells of a first microbial species that mediates said biofouling, the method comprising

(i) contacting the population with a plurality of vectors that are capable of transforming or transducing the cells, each vector comprising a CRISPR array whereby CRISPR arrays are introduced into the host cells, wherein

- (a) each CRISPR array comprises one or more sequences for expression of a crRNA and a promoter for transcription of the sequence(s) in a host cell; and
- (b) each crRNA is capable of hybridising to a target sequence of a host cell to guide Cas (eg, a Cas nuclease) in the host cell to modify the target sequence (eg, to cut the target sequence); the target sequence being a gene sequence for mediating host cell viability; and

wherein the method comprises allowing expression of said crRNAs in the presence of Cas in host cells, thereby modifying target sequences in host cells, resulting in reduction of host cell viability and control of said biofouling.

In an example, the fluid is a liquid. In an example, the fluid is a gaseous fluid.

Systems:

An example system is selected from the group consisting of a:—

Petrochemical recovery, processing, storage or transportation system; hydrocarbon recovery, processing, storage or transportation system; crude oil recovery, processing, storage or transportation system; natural gas recovery, processing, storage or transportation system, (eg, an oil well, oil rig,

oil drilling equipment, oil pumping system, oil pipeline, gas rig, gas extraction equipment, gas pumping equipment, gas pipeline, oil tanker, gas tanker, oil storage equipment or gas storage equipment); Water processing or storage equipment; water reservoir (eg, potable water reservoir); Air or water conditioning (eg, cooling or heating) equipment, eg, a coolant tube, condenser or heat exchanger; Medical or surgical equipment; Environmental (eg, soil, waterway or air) treatment equipment; Paper manufacturing or recycling equipment; Power plant, eg, a thermal or nuclear power plant; Fuel (eg, hydrocarbon fuel, eg, petroleum, diesel or LPG) storage equipment; Mining or metallurgical, mineral or fuel recovery system, eg, a mine or mining equipment; Engineering system; Shipping equipment; Cargo or goods storage equipment (eg, a freight container); Food or beverage manufacturing, processing or packaging equipment; Cleaning equipment (eg, laundry equipment, eg, a washing machine or dishwasher); Catering (eg, domestic or commercial catering) equipment; Farming equipment; Construction (eg, building, utilities infrastructure or road construction) equipment; Aviation equipment; Aerospace equipment; Transportation equipment (eg, a motor vehicle (eg, a car, lorry or van); a railcar; an aircraft (eg, an aeroplane) or a marine or waterway vehicle (eg, a boat or ship, submarine or hovercraft)); Packaging equipment, eg, consumer goods packaging equipment; or food or beverage packaging equipment; Electronics (eg, a computer or mobile phone or an electronics component thereof); or electronics manufacture or packaging equipment; Dentistry equipment; Industrial or domestic piping (eg, a sub-sea pipe) or storage vessel (eg, a water tank or a fuel tank (eg, gasoline tank, eg, a gasoline tank of a vehicle)); Underground equipment; Building (eg, a dwelling or office or commercial premises or factory or power station); Roadway; Bridge; Agricultural equipment; Factory system; Crude oil or natural gas exploration equipment; Office system; and a Household system.

In an example, the system is used in an industry or business selected from the group consisting of agriculture, oil or petroleum industry, food or drink industry, clothing industry, packaging industry, electronics industry, computer industry, environmental industry, chemical industry, aerospace industry, automotive industry, biotechnology industry, medical industry, healthcare industry, dentistry industry, energy industry, consumer products industry, pharmaceutical industry, mining industry, cleaning industry, forestry industry, fishing industry, leisure industry, recycling industry, cosmetics industry, plastics industry, pulp or paper industry, textile industry, clothing industry, leather or suede or animal hide industry, tobacco industry and steel industry. In an example, the surface or fluid to be treated is a surface or fluid of equipment used in said selected industry. In an example, the system is used in the crude oil industry. In an example, the system is used in the natural gas industry. In an example, the system is used in the petroleum industry. In an example, the system is a sea container, platform or rig (eg, oil or gas platform or rig for use at sea or at sea), ship or boat. In an embodiment, such a system is anchored at sea; eg, non-temporarily anchored at sea, eg, has been anchored at sea for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or more months (eg, contiguous months). In an embodiment, such a system is in the waters of a country or state; eg, non-temporarily at sea in such waters, eg, has been in waters of said country for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or more months (eg, contiguous months).

In an example, the substrate surface to be treated comprises stainless steel, carbon steel, copper, nickel, brass,

aluminium, concrete, a plastic or wood. In an example, the substrate is a metal weld or join. In an example, the surface is a metallic (eg, steel or iron) or non-metallic (eg, plastic, concrete, asphalt, wood, rubber or stone) surface. In an example, the metal is an alloy (eg, stainless steel, brass or a nickel-, zinc-, copper-, nickel- or aluminium-alloy). In an example, the surface is a man-made polymer surface. In an example, the surface is a substrate coating. In an example, the substrate is in contact with soil, fresh water or sea water.

In an example, the fluid is potable water; a waterway; brackish water; or a liquid fuel, eg, gasoline or diesel (eg, for a car or motorized vehicle), LPG, kerosine, an alcohol (eg, ethanol, methanol or butanol), liquid hydrogen or liquid ammonia), in an example, the fuel is stored liquid fuel. In an example the fluid is an oil or non-aqueous liquid. In an example, the fluid is a liquid comprised by a waterway or body of water, eg, sea water, fresh water, potable water, a river, a stream, a pond, a lake, a reservoir, stored water (eg, in a water storage tank or cooling equipment), groundwater, well water, water in a rock formation, soil water or rainwater. In an example, the liquid is sea water. In an example, the substrate is in contact with a liquid mentioned in this paragraph. In an example, the fluid or liquid is selected from the group consisting of an oil, an aqueous solution, a hydraulic fracturing fluid, a fuel, carbon dioxide, a natural gas, an oil/water mixture, a fuel/water mixture, water containing salts, ocean or sea water, brackish water, sources of fresh water, lakes, rivers, stream, bogs, ponds, marshes, runoff from the thawing of snow or ice, springs, groundwater, aquifers, precipitation, any substance that is a liquid at ambient temperature (eg, at rtp) and is hydrophobic but soluble in organic solvents, hexanes, benzene, toluene, chloroform, diethyl ether, vegetable oils, petrochemical oils, crude oil, refined petrochemical products, volatile essential oils, fossil fuels, gasoline, mixtures of hydrocarbons, jet fuel, rocket fuel, biofuels. In an example the fluid is an oil/water mixture.

The terms “microbiologically influenced corrosion” or “MIC” as used herein, unless otherwise specified, refer to processes in which any element (substrate) of a system is structurally compromised due to the action of at least one member of a microbial population, eg, bacterial or archaeal population. The term “biofouling” as used herein, unless otherwise specified, refers to processes in which microorganisms (such as bacteria and/or archaea) accumulate on a substrate surface in contact with a fluid (eg, water or an aqueous liquid, or a hydrocarbon, or a petrochemical). Also included is the undesirable accumulation and proliferation of microorganisms (such as bacteria and/or archaea) in a fluid (eg, water or an aqueous liquid, or a hydrocarbon, or a petrochemical), ie, “souring” of the fluid. In an example, the bacteria are comprised by ship or boat ballast water and the bacteria are environmentally undesirable. The term “substrate” as used herein refers to any type of surface on which cells can attach and a biofilm can form and grow or on which biofouling (eg slime or sludge formation) can occur. The substrate may be an “industrial” substrate such as the surface of equipment in an petrochemical, fuel, crude oil or gas piping system, or a “non-industrial” (eg, domestic, eg, household or office) substrate such as a kitchen counter or a shower substrate or a garden substrate.

In an alternative, instead of a population of host bacterial cells, the population is a population of archaeal cells of a first species.

Optionally, said fluid is an aqueous liquid (eg, sea water, fresh water, stored water or potable water).

In an alternative, instead the microbes are algal cells.

Optionally, the microbes are sulphate reducing bacteria (SRB) cells (eg, *Desulfovibrio* or *Desulfotomaculum* cells). In an example, the cells are selected from the group consisting of *Desulfotomaculum nigrificans*, *Desulfacinum infernum*, *Thermodesulfobacterium mobile*, *Thermodesulforhabdus norvegicus*, *Archaeoglobus fulgidus*, *Desulfomicrobium apsheronum*, *Desulfovibrio gabonensis*, *Desulfovibrio longus*, *Desulfovibrio vietnamensis*, *Desulfobacterium ceticum*, *Desulphomaculum halophilum*, *Desulfobacterium vibrioformis* and *Desulfotomaculum thermocisternum* cells. In an example, the population comprises a mixture of two or more of these cell species.

Optionally, the surface or fluid is comprised by a crude oil, gas or petrochemicals recovery, processing, storage or transportation equipment. Crude oil is one of the most important energetic resources in the world. It is used as raw material in numerous industries, including the refinery-petrochemical industry, where crude oil is refined through various technological processes into consumer products such as gasoline, oils, paraffin oils, lubricants, asphalt, domestic fuel oil, vaseline, and polymers. Oil-derived products are also commonly used in many other chemical processes. In an alternative, the fluid is a said consumer product or the surface is in contact with such a consumer product.

Optionally, the surface is in contact with sea water, a fracking liquid or liquid in a well; or wherein the fluid is sea water, a fracking liquid or liquid in a well.

Optionally, step (i) of the method comprises providing a population of microbial cells of a second species (second host cells), the second cells comprising said vectors, wherein the vectors are capable of transfer from the second host cells to the first host cells; and combining the second host cells with the first host cells, whereby vectors are introduced into the first host cells. In an example, the second cell(s) are environmentally-, industrially-, or domestically-acceptable in an environment (eg, in a water or soil environment) and the first host cell(s) are not acceptable in the environment.

Optionally, the first host cells are comprised by a mixture of microbial cells (eg, comprised by a microbial biofilm) before contact with said vectors, wherein the mixture comprises cells of said second species.

Optionally, said second species is a species of *Bacillus* or nitrate-reducing bacteria or nitrate reducing sulfide oxidizing bacteria (NRB)

Optionally, the NRB is selected from the group consisting of *Campylobacter* sp., *Nitrobacter* sp., *Nitrosomonas* sp., *Thiomicrospira* sp., *Sulfurospirillum* sp., *Thauera* sp., *Paracoccus* sp., *Pseudomonas* sp., *Rhodobacter* sp. and *Desulfovibrio* sp; or comprises at least 2 of said species.

Optionally, the NRB is selected from the group consisting of *Nitrobacter vulgaris*, *Nitrosomonas europea*, *Pseudomonas stutzeri*, *Pseudomonas aeruginosa*, *Paracoccus denitrificans*, *Sulfurospirillum deleyianum*, and *Rhodobacter sphaeroides*.

Optionally, the method comprises contacting the host cells of said first species with a biocide simultaneously or sequentially with said vectors. In an example, the vectors and biocide are provided pre-mixed in a composition that is contacted with the host cells.

Optionally, the biocide is selected from the group consisting of tetrakis hydroxymethyl phosphonium sulfate (THPS), glutaraldehyde, chlorine monoxide, chlorine dioxide, calcium hypochlorite, potassium hypochlorite, sodium hypochlorite, dibromonitropropionamide (DBNPA), methylene bis(thiocyanate) (MBT), 2-(thiocyanomethylthio) benzothiazole (TCMTB), bronopol, 2-bromo-2-nitro-

1,3-propanediol (BNPD), tributyl tetradecyl phosphonium chloride (TTPC), taurinamide and derivatives thereof, phenols, quaternary ammonium salts, chlorine-containing agents, quinaldinium salts, lactones, organic dyes, thiosemicarbazones, quinones, carbamates, urea, salicylamide, carbanilide, guanide, amidines, imidazolines, acetic acid, benzoic acid, sorbic acid, propionic acid, boric acid, dehydroacetic acid, sulfurous acid, vanillic acid, p-hydroxybenzoate esters, isopropanol, propylene glycol, benzyl alcohol, chlorobutanol, phenylethyl alcohol, formaldehyde, iodine and solutions thereof, povidone-iodine, hexamethylenetetramine, noxythiolin, 1-(3-chloroallyl)-3,5,7-triazo-1-azoniaadamantane chloride, taurolidine, taurultam, N-(5-nitro-2-furfurylidene)-1-amino-hydantoin, 5-nitro-2-furaldehyde semicarbazone, 3,4,4'-trichlorocarbanilide, 3,4', 5-tribromosalicylanilide, 3-trifluoromethyl-4,4'-dichlorocarbanilide, 8-hydroxy quinoline, 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid, 1,4-dihydro-1-ethyl-6-fluoro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid, hydrogen peroxide, peracetic acid, sodium oxychlorosene, parachlorometaxyleneol, 2,4,4'-trichloro-2'-hydroxydiphenol, thymol, chlorhexidine, benzalkonium chloride, cetylpyridinium chloride, silver sulfadiazine, silver nitrate, bromine, ozone, isothiazolones, polyoxyethylene (dimethylimino) ethylene (dimethylimino) ethylene dichloride, 2-(tert-butylamino)-4-chloro-6-ethylamino-5'-triazine (terbutylazine), and combinations thereof. In an example the biocide is tetrakis hydroxymethyl phosphonium sulfate (THPS). In an example, the biocide is a quaternary ammonium compound.

Optionally, the system is used in an industry operation selected from the group consisting of mining; shipping; crude oil, gas or petrochemicals recovery or processing; hydraulic fracturing; air or water heating or cooling; potable water production, storage or delivery; transportation of hydrocarbons; and wastewater treatment.

Optionally, the surface is a surface of equipment used in said selected industry; or wherein the fluid is a fluid comprised by equipment used in said selected industry.

Optionally, the surface is a surface of kitchen, bathing or gardening equipment; or wherein the fluid is comprised by kitchen, bathing or gardening equipment. For example, the equipment is used in a domestic setting.

Optionally, the fluid is a potable liquid contained in a container (eg, water tank or bottle) and the surface is a surface of the container in contact with the liquid.

Optionally, each vector comprises a mobile genetic element (MGE), wherein the MGE comprises an origin of transfer (oriT) and a said CRISPR array; wherein the MGE is capable of transfer between a host cell of said first species and a further microbial host cell in said industrial or domestic system. For example, the further cell(s) are environmentally-, industrially-, or domestically-acceptable in an environment (eg, in a water or soil environment) and the first host cell(s) are not acceptable in the environment. Optionally, the oriT is functional in the first and further host cells.

Optionally, the first and further host cells are comprised by a biofilm of fluid in contact with said surface; or wherein said cells are comprised by said fluid.

Optionally, each MGE is or comprises an integrative and conjugative element (ICE); or wherein each vector is a phage that is capable of infecting host cells of said first species and each MGE is a phage nucleic acid that is capable of said transfer between the cells. Optionally, each ICE is a transposon, eg, a conjugative transposon. Optionally, each vector is a plasmid, optionally comprising an MGE as

described herein. Optionally, the sequences are comprised by a conjugative transposon of the first cell and/or further cell.

In an example, the method is a method of controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate comprised by a crude oil, gas or petrochemicals recovery, processing, storage or transportation equipment (eg, a crude oil tanker, oil rig or oil drilling equipment), wherein a surface of the substrate is in contact with a population of first host cells, wherein the first host cells are sulphur- or sulphate-reducing bacteria (SRB), extracellular polymeric substance-producing bacteria (EPSB), acid-producing bacteria (APB), sulphur- or sulphide-oxidizing bacteria (SOB), iron-oxidising bacteria (JOB), manganese-oxidising bacteria (MOB), ammonia producing bacteria (AmPB) or acetate producing bacteria (AcPB) of a first species that mediates MIC or biofouling of the substrate, wherein the surface and cell population are in contact with a liquid selected from sea water, fresh water, a fracking liquid or liquid in a well (eg, oil or natural gas well), the method comprising

(i) contacting the cell population with vectors by mixing the liquid with a plurality of vectors that are capable of transforming or transducing first host cells, each vector comprising a CRISPR array whereby CRISPR arrays are introduced into the host cells, wherein

- (a) each CRISPR array comprises one or more sequences for expression of a crRNA and a promoter for transcription of the sequence(s) in a host cell;
- (b) each crRNA is capable of hybridising to a target sequence of a host cell to guide Cas (eg, a Cas nuclease, eg, a Cas9 or Cfp1) in the host cell to modify the target sequence (eg, to cut the target sequence); the target sequence being a gene sequence for mediating host cell viability;
- (c) wherein each sequence of (a) comprises a sequence R1-S1-R1' for expression and production of the respective crRNA in a first host cell, wherein R1 is a first CRISPR repeat, R1' is a second CRISPR repeat, and R1 or R1' is optional; and S1 is a first CRISPR spacer that comprises or consists of a nucleotide sequence that is 70, 75, 80, 85, 90 or 95% or more identical to a target sequence of a said first host cell and

(ii) allowing expression of said crRNAs in the presence of Cas in host cells, thereby modifying target sequences in host cells, resulting in reduction of host cell viability and control of MIC or biofouling of said substrate. In an embodiment, both R1 and R1' are present.

In an example, the method is a method of controlling bacterial biofouling in ballast water of a ship or boat, wherein the water comprises a population of first host cells of a first microbial species that mediates said biofouling, the method comprising

(i) contacting the population with a plurality of vectors that are capable of transforming or transducing the cells, each vector comprising a CRISPR array whereby CRISPR arrays are introduced into the host cells, wherein

- (a) each CRISPR array comprises one or more sequences for expression of a crRNA and a promoter for transcription of the sequence(s) in a host cell; and
- (b) each crRNA is capable of hybridising to a target sequence of a host cell to guide Cas (eg, a Cas nuclease) in the host cell to modify the target sequence (eg, to cut the target sequence); the target sequence being a gene sequence for mediating host cell viability; and

(ii) allowing expression of said crRNAs in the presence of Cas in host cells, thereby modifying target sequences in host cells, resulting in reduction of host cell viability and control of said biofouling.

Optionally, the first host cells are *Vibrio cholerae*, *E coli* or *Enterococci* sp cells.

Optionally, step (i) comprises mixing the ballast water with the vectors, eg, in the hull of a ship or boat. Optionally, the ship or boat is a marine vehicle and the water is sea water. Optionally, instead of a ship or boat, the ballast water is comprised by a container or a drilling platform at sea, eg, an oil platform or oil rig. In an example, the ship, boat, container, platform or rig is anchored at sea (ie, not temporarily in its location).

In an example, the method is a method of discharging ballast water from a ship or boat, wherein the discharged ballast water comprises water treated by the method. Optionally, the water is discharged into a body of water, eg, a sea, ocean or waterway (eg, a river, canal, lake or reservoir) or into a container.

Paragraphs:

The invention provides the following Paragraphs, which are supported by the Examples below:—

1. A programmable Cas (eg, Cas3 or Cas9) nuclease for use in a method of treating *E coli* or *C difficile* infection of a subject, wherein the Cas nuclease is programmable with a guide RNA to cut a target site comprised by the genomes of *E coli* or *C difficile* bacteria that have infected the subject, whereby *E coli* or *C difficile* cells are killed, or growth or proliferation of the cells is reduced, the treatment method comprising exposing the subject to the Cas nuclease wherein the nuclease is programmed with guide RNA to cut the target site, whereby genomes of the *E coli* or *C difficile* bacteria comprised by the subject are cut and the infection of the subject is reduced by at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
2. A programmable Cas (eg, Cas3 or Cas9) nuclease (optionally according to paragraph 1) for use in a method of treating *E coli* or *C difficile* infection of a subject, wherein the Cas nuclease is programmable with a guide RNA to cut a target site comprised by the genomes of *E coli* or *C difficile* bacteria that have infected the subject, whereby *E coli* or *C difficile* cells are killed, or growth or proliferation of the cells is reduced, the treatment method comprising exposing the subject to the Cas nuclease wherein the nuclease is programmed with guide RNA to cut the target site, whereby genomes of the *E coli* or *C difficile* bacteria comprised by the subject are cut and the infection of the subject is reduced, wherein a reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.
3. The nuclease of any preceding Paragraph, wherein at least 60% of the infection is reduced by 60 minutes after exposing the subject to the programmed nuclease.
4. The nuclease of any preceding Paragraph, wherein the nuclease (eg, programmed nuclease) and/or a nucleic acid encoding the guide RNA is administered to the subject at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1.
5. The nuclease of any preceding Paragraph, wherein the method comprises reducing the infection such that the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.

6. The nuclease of any preceding Paragraph, wherein the method comprises administering to the subject the RNA or a nucleic acid that encodes the RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject. 5
7. The nuclease of any preceding Paragraph, wherein the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid encoding the RNA to the subject. 10
8. The nuclease of Paragraph 7, wherein the subject comprises the nuclease prior to administration of the RNA or nucleic acid to the subject.
9. The nuclease of any preceding Paragraph, wherein a plurality of viruses (eg, phage) are administered to the subject, wherein each virus comprises a copy of a nucleic acid encoding the RNA, wherein the viruses infect the microbes comprised by the subject to deliver thereto the nucleic acid. 15
10. The nuclease of Paragraph 9, wherein the ratio of administered viruses:microbes comprised by the subject is from 10 to 150. 20
11. The nuclease according to any preceding Paragraph, wherein the subject is a human or animal, optionally wherein the subject is a human over 65 years of age or is a paediatric patient. 25
12. The nuclease according to Paragraph 11, wherein the infection is an infection of the lungs, abdomen or urinary tract; or wherein the subject has undergone surgery, is on an immunosuppressant medication and/or is suffering from a chronic disease. 30
13. The nuclease according to any preceding Paragraph, wherein the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment. 35
14. The nuclease according to any preceding Paragraph, wherein the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. 40
15. The nuclease according to any one of Paragraphs 11 to 14, wherein the method treats or prevents septicaemia and/or sepsis (eg, septic shock) in the subject. 45
16. The nuclease of Paragraph 16, wherein at the start of the treatment, the subject (eg, a human) has a temperature of $<36^{\circ}\text{C}$. or $>38^{\circ}\text{C}$.; a heart rate of $>90/\text{min}$, a respiratory rate of >20 breaths/min or $\text{PaCO}_2 < 4.3$ kPa; and white blood cell count of $<4000/\text{mm}^3$ or $>12,000/\text{mm}^3$ 50
17. The nuclease of Paragraph 15 or 16, wherein at the start of the treatment, the subject (eg, a human) has presence of two or more of the following: abnormal body temperature, abnormal heart rate, abnormal respiratory rate, abnormal blood gas and abnormal white blood cell count. 55
18. The nuclease of any preceding Paragraph, wherein the subject is a human or animal and the microbes are bacteria (eg, *E coli* or *C difficile*), wherein blood infection of the subject by the bacteria is reduced at least 100- or 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. 60
19. The nuclease of any one of Paragraphs 11 to 18, wherein the blood of the subject is infected with from 10^7 to 10^{12} CFU/ml of the bacteria immediately before the treatment. 65

20. The nuclease according to any one of Paragraphs 1 to 10, wherein the subject is a plant.
21. The nuclease according to any preceding Paragraph, wherein the bacteria are comprised by a microbiome.
22. The nuclease according to Paragraph 21, wherein the microbiome comprises *Lactobacillus* and/or *Streptococcus* bacteria.
23. The nuclease according to any preceding Paragraph, wherein the *E coli* are EHEC *E coli*.
24. The nuclease according to any preceding Paragraph, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.
25. A plurality of viruses (eg, phage or phagemids for producing phage) for use with the nuclease of any preceding Paragraph in the method of treatment, wherein each virus comprises a copy of a nucleic acid encoding the RNA, wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.
26. A composition comprising a plurality of nucleic acids for programming the nuclease of any one of Paragraphs 1 to 24 in the method of treatment, wherein each nucleic acid is a nucleic acid as defined in any one of Paragraphs 6 to 9.
27. A CRISPR/Cas system comprising a nuclease according to any preceding Paragraph for use in the method of treatment, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of the microbes.
28. A guide RNA or a DNA encoding a guide RNA for use in the system of Paragraph 27 for use in the method of treating an acute microbial infection in the subject, eg, septicaemia or sepsis.
29. A nucleic acid vector comprising the guide RNA or DNA recited in Paragraph 27 or 28.
30. The vector of Paragraph 29 wherein the vector is a phage, phagemid, viriophage, virus, plasmid (eg, conjugative plasmid) or transposon.
31. An anti-sepsis or anti-septicaemia composition for administration to a human or animal for treating sepsis or septicaemia, the composition comprising a plurality of vectors, wherein each vector is according to Paragraph 29 or 30.
32. A method of treating an acute microbial infection of a subject, wherein the method is as defined by any preceding Paragraph.
33. Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of any one of Paragraphs 1 to 25 and 27 to 30, in the manufacture of a composition for carrying out a method of treatment as defined by any preceding Paragraph, wherein the subject is an organism other than a human or animal.
34. Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of any one of Paragraphs 1 to 25 and 27 to 30, in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method com-

- prising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.
35. Use of a programmable nuclease in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.
36. The use of Paragraph 33, 34, 35, wherein the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject or substrate at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1.
37. The use of any one of Paragraphs 33 to 36, wherein the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
38. The use of any one of Paragraphs 33 to 37, wherein the reduction of the infection is maintained by at least 100-fold for at least 60 minutes (eg, at least 120 minutes) after exposing the subject to the programmed nuclease.
39. The use of any one of Paragraphs 33 to 38, wherein the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.
40. The use of any one of Paragraphs 33 to 39, wherein the method comprises administering to the subject or substrate a RNA or a nucleic acid that encodes an RNA for expression of the RNA in or on the subject or substrate, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject or substrate.
41. The use of Paragraph 40, wherein the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject or substrate.
42. The use of Paragraph 40, wherein the subject or substrate comprises the nuclease prior to administration of the RNA or nucleic acid.
43. The use of any one of Paragraphs 40 to 42, wherein a plurality of viruses (eg, phage) are administered to the subject or substrate, wherein each virus comprises a copy of the nucleic acid, wherein the viruses infect the microbes comprised by the subject or substrate to deliver thereto the nucleic acid.
44. The use of Paragraph 43, wherein the ratio of administered viruses:microbes is from 10 to 150.
45. The use of any one of Paragraphs 33 to 44, wherein the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
46. The use of any one of Paragraphs 44 to 45, wherein the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at

- least 120, 145 or 180 minutes) after exposing the subject or substrate to the programmed nuclease.
47. The use of any one of Paragraphs 33 to 46, wherein the subject is a plant; or wherein the substrate is a metallic, plastic, concrete, stone, wood, glass or ceramic substrate.
48. The use of any one of Paragraphs 33 to 47, wherein the microbes are bacteria.
49. The use according to Paragraph 48, wherein the bacteria are gram positive bacteria.
50. The use according to Paragraph 48 or 49, wherein the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonnorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).
51. The use of any one of Paragraphs 33 to 50, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.

Treatment of Pathogenic Bacterial Infections

Infectious complications are a serious cause of morbidity and mortality in cancer patients, especially those with underlying haematological malignancies where autopsy studies demonstrate that approximately 60% of deaths are infection related. Although fewer data exist on infectious mortality in patients with solid organ tumours, approximately 50% of these patients are estimated to have an infection as either the primary or an associated cause of death (“Epidemiology of Infections in Cancer Patients”, in “Infectious Complications in Cancer Patients”, Springer International Publishing Switzerland (2014)). Bacterial infections dominate. These infectious complications remain a significant limitation of cancer treatment modalities.

The detrimental effects of classic antibiotic treatment with broad-spectrum antibiotics have been demonstrated in immune checkpoint inhibitor (ICI)-treated cancer patients. Routy et al investigated how the gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumours (Routy et al Science 2018, 359, 91-97). In this work, the authors also analyzed datasets for infections/antibiotic use in patients with advanced NSCLC (n=140), renal cell carcinoma (n=67), or urothelial carcinoma (n=42) who received antibody ICI against PD-1/PD-L1 interaction after one or several prior therapies. Among these patients, they were prescribed broad-spectrum antibiotics (beta-lactam+/- inhibitors, fluoroquinolones, or macrolides) within 2 months before, or 1 month after, the first administration of PD-1/PD-L1 mAb. Patients generally took antibiotic orally for common indications (dental, urinary, and pulmonary infections). The detrimental effect of treating infections in cancer patients undergoing ICI therapy with classical, broad-spectrum antibiotics was observed. See FIG. 8, which shows that the antibiotic treatment during ICI therapy has fatal outcomes: there was a medial overall survival of 21.9 months in the absence of antibiotic treatment, compared to an overall survival of 9.8 months with antibiotic treatment. So, the median overall survival in patients treated with classical antibiotics is <50% (or >12 months shorter) that of patients not receiving antibiotic treatment.

The work by Gopalakrishnan et al is another recent example lending support to the importance of a “healthy” microbiome in immuno-oncology therapy outcomes (Go-

palakrishnan et al, *Science* 2018, 359, 97-103). See FIGS. 9A and 9B. It was observed that the gut microbiome modulates the efficacy of anti-PD-1 inhibition in melanoma patients.

Several other studies add to the expanding evidence base of the critical link between the microbiome and immunology outcomes:

“Microbiota: a key orchestrator of cancer therapy”, *Nat. Rev. Cancer* 2017, 17, 271-285

Matson et al, *Science* 2018, 359, 104-108

L. Derosa et al *Annals of Oncology* 2018 (epub 30 Mar. 2018)

M. Vétizou et al *Science*. 2015, 350, 1079-84

Sivan et al *Science* 2015, 350, 1084-1089

Another report—claiming to be the first systematic review of infection among patients receiving immune checkpoint blockade for cancer therapy—investigated serious infections in melanoma patients treated with immune checkpoint inhibitors (against CTLA-4, PD-1, and/or PD-L1) (M. Del Castillo et al *Clin. Infect. Dis.* 2016, 63, 1490-1493). Serious infections were defined as infections requiring hospitalization or parenteral antimicrobials. Of 740 patients (898 courses of immune checkpoint blockade), serious infection developed in 54 patients (7.3%). Nine patients (17%) were deemed to have died of an infection. Total number of infections was 58, as some patients developed >1 infection. The majority of infections were bacterial in origin (~80%; i.e., bacterial infections: 80% of 7.3%: 5.8% of patients). Pneumonia and bloodstream infections were the two dominating bacterial infection types.

Immune checkpoint-blocking drugs are associated with immune-related adverse effects (irAEs) related to the upregulated immune system. The complications are managed with immunosuppressive drugs, such as steroids (immunosuppression is a risk factor for subsequent opportunistic infections). Of the 740 patients, 46% received steroids during the course of treatment. Risk of serious infections was 13.5% in the cohort receiving corticosteroids or infliximab (vs. 7.3% in the overall population).

In yet another report, the emerging concern of infectious diseases in lung cancer patients receiving ICI therapy was investigated. Of 84 NSCLC patients receiving nivolumab (a PD-1 inhibitor), 20 patients (23.8%) developed an infectious disease. Bacterial infections accounted for 75% of infections; i.e., bacterial infections in 18% of patients. Most common type of bacterial infection was pneumonia. See K. Fujita et al *Eur. Resp. J.* 2017, 50, OA1478.

The Gram-negative *bacillus E. coli* is one of the most common causes of bacteraemia in patients with cancer. The all-cause 30-day mortality rate for this pathogen is high (~15%) (Y. E. Ha et al *Int. J. Antimicrob. Agen.* 2013, 42, 403-409). Published estimates of 30-day all-cause mortality among *E. coli* bacteraemia patients (cancer/non-cancer) vary from around 10 to 35% (J. K. Abernethy et al *Clin. Microbiol. Infect.* 2015, 21, 251.e1-251.e8), clearly highlighting the high burden associated with just this pathogen. Overall, causative pathogens in bacteraemia are primarily Gram-negative bacteria (65%), with *E. coli* (18.3%), *P. aeruginosa* (18.3%), and *K. pneumoniae* (17.3%) being the most common organisms encountered; the three pathogens together account for 54% of the bacteraemia cases, or 85% of Gram-negative cases, according to a study investigating >100 bacteraemia cases in cancer patients (G. Samonis et al *Support Care Cancer* 2013, 21, 2521-2526). In-hospital mortality was 26.2% in this study. Comparable numbers can be found elsewhere. For example, a study of neutropenic and non-neutropenic adult cancer patients with bloodstream

infections investigated 399 cases of bloodstream infections in 344 cancer patients: The largest causative pathogen group was Gram-negative bacilli (45%). Of the clinical isolates, *E. coli* (35%) accounted for the most cases within Gram-negatives, followed by *K. pneumoniae* (20%) and *P. aeruginosa* (19%) (E. Velasco et al *Eur. J. Clin. Microbiol. Infect. Dis.* 2006, 25, 1-7). The three pathogens collectively account for 33% of the bacteraemia cases (or 74% of Gram-negative cases). The overall 30-day mortality rate was 32% in this study. Two other reports looked at causative agents of bloodstream infection in patients with solid tumours and also found Gram-negative bacteria to be the dominating pathogen type (47-55% of the infections, across several hundred patients) (M. Marin et al *Medicine* 2014, 93, 143-149; M. Anatoliotaki et al *Infection* 2004, 32, 65-71; see also C. Gudiol et al *Virulence* 2016, 7, 298-308). In the larger of the two studies (with more robust numbers for individual pathogens), the three main pathogens within the Gram-negative group were again *E. coli* (55%), *P. aeruginosa* (18%), and *Klebsiella* spp. (11%)—corresponding to 92% of the Gram-negative cases, or 51% of the 528 total cases of bloodstream infections studied.

The above data on specific causative infectious pathogens in cancer patients are summarized in Table 5 below.

Thus, available data on bloodstream infections in cancer indicate that Gram-negative pathogens are involved in 45-65% of the infection cases, with three key pathogens—*E. coli*, *K. pneumoniae*, and *P. aeruginosa*—being the culprits in the vast majority of Gram-negative cases (73-92%).

The inventors, thus, formulated an oncologist’s dilemma: Reduction in efficacy of the cancer therapy is likely due to the reduced microbiome diversity resulting from antibiotic therapy

At least 1/3 of patients on checkpoint inhibitors get serious and life-threatening infections

Not treating these infections could result in death from the infection (1-2 weeks)

Treatment with classic antibiotics leads to reduction in progression free survival after 4 years from >40% to around 10%.

The choice is to treat the immediate need of a potentially fatal infection (which must be addressed) at the risk of seriously undermining the cancer therapy.

The inventors realised, therefore, that there is a need for methods that can treat a bacterial pathogenic infection in a different way that minimizes compromise to the cancer therapy. The inventors realised that this need would also be useful in other therapy settings where the microbiome composition can modulate therapy outcomes, eg, in transplant settings.

Whilst not wishing to be bound by any particular theory, the inventors believe that alleviating the detrimental effect of traditional antibiotic therapy on overall survival in ICI patients using the invention may, in some embodiments, translate to as much as a doubling of overall survival (or >12 months). Capturing a treatment effect of several months in terms of median overall survival is a very substantial achievement in this space. In fact, an effect size of this order of magnitude is comparable to the outcomes reported for ICI trials (i.e., where benefits usually are measured in months, not years). Additionally, PD-1/PD-L1 drugs are projected to dominate the ICI market. In 2023, PD-1/PD-L1 are projected to account for 94% of \$46B USD global sales of ICIs (CTLA-4 blockers only account for 6%), source: “Landscape & Forecast Immune Checkpoint Inhibitors”, Decision Resources, December 2017. Thus, a need for improving treatment using immune checkpoint inhibitors of PD-1 or

PD-L1 is particularly pressing in medicine, and we believe that the present invention finds particular benefit in this respect.

In an example, the method removes the need to administer a classic antibiotic, such as a broad-spectrum antibiotic (or any other one disclosed herein). In another example, the invention reduces the amount or dosing frequency of a classic antibiotic, such as a broad-spectrum antibiotic (or any other one disclosed herein) that is administered to the subject for treating the infection. For example, the subject can be administered a low-dose broad-spectrum antibiotic (eg, 50, 40, 30, 20, 10% or less of a conventional dose) whilst the guided nuclease cutting is used, and thus treatment of the infection in this setting. The invention may be particularly beneficial for patients on immunosuppressants, eg, for cancer patients, transplant patients or patients suffering from a viral infection (eg, HIV (human immunodeficiency virus), CMV (cytomegalovirus) or RSV (respiratory syncytial virus) infection).

The term “broad-spectrum antibiotic” can refer to an antibiotic that acts on the two major bacterial groups, gram-positive and gram-negative, or any antibiotic that acts against a wide range of disease-causing bacteria. These medications are used when a bacterial infection is suspected but the group of bacteria is unknown (also called empiric therapy) or when infection with multiple groups of bacteria is suspected. Although powerful, broad-spectrum antibiotics pose specific risks, particularly the disruption of native, normal bacteria and the development of antimicrobial resistance. Examples of commonly used broad-spectrum antibiotics are: Aminoglycosides (except for streptomycin), Ampicillin, Amoxicillin, Amoxicillin, clavulanic acid (Augmentin), Carbapenems (e.g. imipenem), Piperacillin, tazobactam, Quinolones (e.g. ciprofloxacin), Tetracyclines, Chloramphenicol, Ticarcillin, Trimethoprim and sulfamethoxazole (Bactrim). In veterinary medicine, examples are co-amoxiclav, (eg, in small animals), penicillin, streptomycin, oxytetracycline and potentiated sulfonamides.

Clauses:—

The invention, therefore, in one aspect provides the following Clauses that are directed to the treatment of a pathogenic bacterial infection using a programmed nuclease.

1. A method for treating a pathogenic bacterial infection in a human or animal subject caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a programmable nuclease that is programmed to cut the target site, wherein the subject is suffering from a further disease or condition other than the pathogenic bacterial infection and the method comprises administering a therapy to the subject for treating or preventing the further disease or condition, wherein the nuclease treats the infection and the therapy is efficacious in the presence of the programmed nuclease to treat or prevent the disease or condition.

In an example, Clause 1 provides:—

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the

subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-PD-1 antibody optionally selected from pembrolizumab (or KEYTRUDA™) and nivolumab (or OPDIVO™); and
- (b) The cancer is selected from metastatic melanoma; renal cell carcinoma; bladder cancer; a solid tumour; non-small cell lung cancer (NSCLC); forehead and neck squamous cell carcinoma (HNSCC); Hodgkin’s lymphoma; a cancer that overexpresses PD-L1 and no mutations in EGFR or in ALK; colorectal cancer and hepatocellular carcinoma; and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-PD-L1 antibody optionally selected from atezolimumab (or TECENTRIQ™), avelumab (or BAVENCIO™) and durvalumab (or IMFINZI™); and
- (b) The cancer is selected from metastatic melanoma; renal cell carcinoma; a solid tumour; non-small cell lung cancer (NSCLC); forehead and neck squamous cell carcinoma (HNSCC); Merkel cell carcinoma; Hodgkin’s lymphoma; a cancer that overexpresses PD-L1 and no mutations in EGFR or in ALK; colorectal cancer and hepatocellular carcinoma; and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing

first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-CD52, antibody optionally alemtuzumab (or CAMPATH™); and
- (b) The cancer is B-cell chronic lymphocytic leukemia (CLL); and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-CD20 antibody, optionally of atumumab (or ARZERRA™) or rituximab (or RITUXAN™); and
- (b) The cancer is B-cell chronic lymphocytic leukemia (CLL) (eg, refractory CLL) or non-Hodgkin lymphoma; and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-KIR antibody, optionally lirilumab; and
- (b) The cancer is optionally acute myeloid leukaemia or squamous cell carcinoma of the head and neck (SCCHN); and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus*

(eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-CD19 CAR-T optionally selected from axicabtagene ciloleucel (or YESCARTA™) and tisagenlecleucel (or KYMRIA™); and
- (b) The cancer is selected from a B-cell lymphoma (eg, non-Hodgkin's lymphoma (NHL); diffuse large B-cell lymphoma (DLBCL); primary mediastinal large B-cell lymphoma; or high grade B-cell lymphoma); B-cell acute lymphoblastic leukaemia (ALL); or central nervous system lymphoma; and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

Alternatively, the CAR-T is an anti-CD30, CD38 or CD22 CAR-T. In an example the cancer is large B-cell lymphoma after at least two other kinds of treatment failed. In an example the cancer is high grade B-cell lymphoma and DLBCL arising from follicular lymphoma. In an example the cancer is relapsing/remitting B cell acute lymphoblastic leukaemia. In an example the cancer is primary central nervous system lymphoma.

In an example, the nuclease treats the infection without causing reduction in efficacy of the therapy. In an embodiment, "without causing reduction in efficacy of the therapy" means the efficacy of the therapy compared to a reduction caused in patients by the administration of a broad-spectrum antibiotic (or an antibiotic disclosed herein) that kills a plurality of different species, wherein the plurality comprises the first species. In an embodiment, "without causing reduction in efficacy of the therapy" means the efficacy of the therapy is reduced by no more than 70, 80, 90 or 95% compared to administration of the therapy in the absence of treatment of the pathogenic bacterial infection (or compared to therapy as typically achieved in patients suffering from the disease or condition and receiving said therapy therefor). This may be assessed, for example, by determining the duration of progression-free survival of the subject or treatment of the disease or condition, or overall survival of the subject; and/or by determining a reduction in one or more symptoms of the disease or condition.

In an example, the infection is treated completely or substantially completely. In another example, the infection is reduced (eg, by at least 80, 90 or 95% as determined by a marker of the infection or a symptom thereof). A marker may, for example, be CFUs of bacteria of the first species or strain per ml of a blood sample taken from the patient after the method has been carried out, eg, within 24 hours of that method being carried out, eg, from 1-12 hours or 1-24 hours

after carrying out the method or from 1-12 hours or 1-24 hours after administering a RNA or DNA encoding the RNA to programme the nuclease in the subject. For example, the RNA is a guide RNA and the nuclease is Cas (eg, a Cas3 or a Cas9). The reduction may be compared to a sample taken from the subject immediately prior to the commencement of the method. Alternatively, the sample may be a stool, saliva or urine sample.

In an example, the invention increases overall survival rate in a human subject (compared to median overall survival rate in humans suffering from the same cancer and receiving the same cancer therapy treatment (eg, administration of the same immune checkpoint inhibitor, such as nivolumab, pembrolizumab or another antibody disclosed herein)). In an example any composition, or other product of the invention herein is provided for use in such method of treatment.

In an example, the method is practiced on a population of human subjects and the median overall survival rate for the population is 120-250% (eg, 150-200%) of the median overall survival rate in humans suffering from the same cancer and receiving the same cancer therapy treatment (eg, administration of the same immune checkpoint inhibitor, such as nivolumab, pembrolizumab or another antibody disclosed herein). In an example any composition, or other product of the invention herein is provided for use in such method of treatment.

A "pathogenic bacterial infection" is a health-threatening infection of the subject, for example, a life-threatening infection. In an embodiment, a pathogenic bacterial infection is an infection requiring hospitalization or parenteral antimicrobials. The infection may be an acute bacterial infection, such as a systemic infection or a localised infection. Bacterial pathogens often cause infection in specific areas of the body. Others are generalists. A pathogenic bacterial infection is contrasted with an infection of commensal bacteria, such as commensal gut bacteria; in this case the bacteria do not cause an immediate health- or life-threatening situation.

The infection (or symptom thereof) can be any of the following:—

Bacterial vaginosis: this is caused by bacteria that change the vaginal microbiota caused by an overgrowth of bacteria that crowd out the *Lactobacilli* species that maintain healthy vaginal microbial populations.

Bacterial meningitis: this is a bacterial inflammation of the meninges, that is, the protective membranes covering the brain and spinal cord.

Bacterial pneumonia: this is a bacterial infection of the lungs.

Urinary tract infection: this is predominantly caused by bacteria. Symptoms include the strong and frequent sensation or urge to urinate, pain during urination, and urine that is cloudy. The main causal agent is *Escherichia coli*. Bacteria can ascend into the bladder or kidney and causing cystitis and nephritis.

Bacterial gastroenteritis: this is caused by enteric, pathogenic bacteria. These pathogenic species are usually distinct from the usually harmless bacteria of the normal gut flora. But a different strain of the same species may be pathogenic.

Bacterial skin infections: these include:

Impetigo, which is a highly contagious bacterial skin infection commonly seen in children. It is caused by *Staphylococcus aureus*, and *Streptococcus pyogenes*.

Erysipelas, which is an acute *streptococcus* bacterial infection of the deeper skin layers that spreads via with lymphatic system.

Cellulitis, which is a diffuse inflammation of connective tissue with severe inflammation of dermal and subcutaneous layers of the skin. Cellulitis can be caused by normal skin flora or by contagious contact, and usually occurs through open skin, cuts, blisters, cracks in the skin, insect bites, animal bites, burns, surgical wounds, intravenous drug injection, or sites of intravenous catheter insertion. In most cases it is the skin on the face or lower legs that is affected, though cellulitis can occur in other tissues.

In an example, the first bacteria are *Streptococcus* and the patient is suffering from chest infection, cellulitis or tonsillitis. In an example, the first bacteria are *Enterococcus* and the patient is suffering from bladder infection or septicaemia. In an example, the first bacteria are *Pseudomonas aeruginosa* and the patient is suffering from diarrhoea. In an example, the first bacteria are *E coli* and the patient is suffering from diarrhoea.

2. The method of Clause 1, wherein the subject is a cancer patient and the therapy is a cancer therapy.

3. The method of Clause 2, wherein the therapy comprises administration of a haematopoietic stem cell transplant, chemotherapeutic agent, immune checkpoint inhibitor, immune checkpoint agonist or an immune cell (eg, T-cell and/or NK cell) enhancer; adoptive cell therapy (eg, CAR-T therapy); radiation or surgery.

In an example, the therapy is immunotherapy. Examples of suitable immunotherapy are administration of adoptive cell therapy (eg, CAR-T therapy), an immune checkpoint inhibitor, an immune checkpoint agonist or an immune cell (eg, T-cell and/or NK cell) enhancer. For example, administration of an anti-CTLA4, PD-1, PD-L1, PD-L2, LAG3, OX40, CD28, BTLA, CD137, CD27, HVEM, KIR, TIM-3, VISTA, ICOS, GITR, TIGIT or SIRPa antibody, such as administration of an antibody selected from ipilimumab (or YERVOY™), tremelimumab, nivolumab (or OPDIVO™), pembrolizumab (or KEYTRUDA™), pidilizumab, BMS-936559, durvalumab and atezolizumab, or a CAR-T therapy such as axicabtagene ciloleucel (Yescarta™) or tisagenlecleucel (Kymriah™).

In an example, the immune enhancer comprises an interleukin-2 (IL-2) or fragment or deletion mutant thereof.

In an example, the surgery comprises the removal of necrotic or cancerous tissue.

In an example, the chemotherapy comprises administration of a platinum-containing chemotherapy drug. In an example, the chemotherapy comprises administration of gefitinib.

In an example, the therapy comprises administering Cyclophosphamide, methotrexate and 5-fluorouracil (CMF); or doxorubicin and cyclophosphamide (AC); docetaxel, doxorubicin and cyclophosphamide (TAC); or doxorubicin, bleomycin, vinblastine and dacarbazine (ABVD); or mustine, vincristine, procarbazine and prednisolone (MOPP); cyclophosphamide, doxorubicin, vincristine and prednisolone (CHOP); bleomycin, etoposide and cisplatin (BEP); epirubicin, cisplatin and 5-fluorouracil (ECF); or epirubicin, cisplatin and capecitabine (ECX); methotrexate, vincristine, doxorubicin and cisplatin (MVAC); cyclophosphamide, doxorubicin and vincristine (CAV); or 5-fluorouracil, folinic acid and oxaliplatin (FOLFOX).

In an example, the cancer is breast cancer and the therapy comprises administering CMF or AC. In an example, the cancer is Hodgkin's lymphoma and the therapy comprises

administering TAC, ABVD or MOPP. In an example, the cancer is Non-Hodgkin's lymphoma and the therapy comprises administering CHOP. In an example, the cancer is germ cell cancer and the therapy comprises administering BEP. In an example, the cancer is stomach cancer and the therapy comprises administering ECF or ECX. In an example, the cancer is bladder cancer and the therapy comprises administering MVAC. In an example, the cancer is lung cancer and the therapy comprises administering CAV. In an example, the cancer is colorectal cancer and the therapy comprises administering FOLFOX.

4. The method of Clause 3, wherein the therapy is an immune checkpoint inhibitor antibody.

Optionally the antibody is an anti-CTLA4, PD-1, PD-L1, PD-L2, LAG3, OX40, CD28, BTLA, CD137, CD27, HVEM, KIR, TIM-3, VISTA, ICOS, GITR, TIGIT or SIRPa antibody. In an example, the antibody is an anti-PD-1 antibody. In an example, the antibody is an anti-PD-L1 antibody. In an example, the antibody is an anti-CTLA4 antibody.

5. The method of Clause 3, wherein the therapy is administration of an antibody selected from ipilimumab (or YERVOY™), tremelimumab, nivolumab (or OPDIVO™), pembrolizumab (or KEYTRUDA™), pidilizumab, BMS-936559, durvalumab and atezolizumab.

Optionally, the antibody (eg, anti-PD-L1 antibody) is administered with an anti-CTLA4 antibody (eg, ipilimumab or tremelimumab).

In an example, the an anti-PD-1 antibody herein is selected from nivolumab, pembrolizumab, pidilizumab, OPDIVO®, KEYTRUDA®, AMP-514, REGN2810, CT-011, BMS 936559, MPDL3280A and AMP-224.

In an example, the an anti-CTLA4 antibody herein is selected from tremelimumab, YERVOY® and ipilimumab.

In an example the therapy is administration of an anti-KIR antibody, eg, lirilumab.

In an example, the checkpoint inhibitor is selected from an inhibitor of CTLA-4, PD-1, PD-L1, PD-L2, LAG-3, BTLA, B7H3, B7H4, TIM3, KIR, or A2aR. In certain aspects, the immune checkpoint inhibitor is a human programmed cell death 1 (PD-1) axis-binding antagonist. In some aspects, the PD-1 axis-binding antagonist is selected from the group consisting of a PD-1 binding antagonist, a PD-L1-binding antagonist and a PD-L2-binding antagonist. In certain aspects, the PD-1 axis-binding antagonist is a PD-1-binding antagonist. In some aspects, the PD-1-binding antagonist inhibits the binding of PD-1 to PD-L1 and/or PD-L2.

In some embodiments, the immune checkpoint inhibitor is a PD-L1 antagonist such as durvalumab, also known as MEDI4736, atezolizumab, also known as MPDL3280A, or avelumab, also known as MSB00010118C. In certain aspects, the immune checkpoint inhibitor is a PD-L2 antagonist such as rHIgM12B7. In some aspects, the immune checkpoint inhibitor is a LAG-3 antagonist such as IMP321 or BMS-986016. The immune checkpoint inhibitor may be an adenosine A2a receptor (A2aR) antagonist such as PBF-509.

In some embodiments, the antibody described herein (such as an anti-PD-1 antibody, an anti-PD-L1 antibody, or an anti-PD-L2 antibody) further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group

consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic, CHO, Cos or HEK cells. In a still further specific aspect the minimal effector function results from an "effector-less Fc mutation" or aglycosylation.

For example, the therapy comprises a haemopoietic stem cell transplant, eg, a bone marrow transplant (such as when the patient is a cancer patient, eg, a blood cancer or leukaemia patient).

For example, the therapy comprises a stem cell transplant, a skin graft, or an organ transplant, eg, a heart, liver, kidney or lung transplant.

6. The method of Clause 1 or 2, wherein the therapy is a tissue, organ or cell transplant.

7. The method of any preceding Clause, wherein the treatment of the bacterial infection is carried out simultaneously with the administration of the therapy to the subject.

8. The method of any one of Clauses 1 to 6, wherein the treatment of the bacterial infection is carried out immediately before administering the therapy to the subject.

In an example, the treatment of the bacterial infection is carried out no more than 7, 6, 5, 4, 3, 2, or 1 day, or 24, 12, 6, 5, 4, 3, 2, 1 or 0.5 hours before the therapy of the further disease or condition. In an example, the treatment of the bacterial infection is carried out no more than 7, 6, 5, 4, 3, 2, or 1 day, or 24, 12, 6, 5, 4, 3, 2, 1 or 0.5 hours after the therapy of the further disease or condition.

The treatment of the infection and the administration of the therapy may be carried out simultaneously or sequentially.

9. The method of any one of Clauses 1 to 6, wherein the treatment of the bacterial infection is carried out immediately after administering the therapy to the subject.

10. The method of any preceding Clause, wherein the method comprises administering to the subject a RNA (eg, a gRNA) or a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in first bacteria comprised by the subject, thereby killing the first bacteria.

The RNA or nucleic acid is, for example, administered to the subject or patient orally, by IV injection, by subcutaneous injection or by inhalation.

11. The method of any preceding Clause, comprising administering a vector (eg, phage or plasmids) to the subject, wherein the vector encodes the programmable nuclease.

The nuclease is, for example, administered to the subject or patient orally, by IV injection, by subcutaneous injection or by inhalation.

12. The method of any one of Clauses 1 to 10, wherein the programmable nuclease is an endogenous nuclease (eg, Cas nuclease) of the first cells.

13. The method of any preceding Clause, wherein the efficacy of the therapy in the presence of the programmed nuclease is greater than the efficacy of the therapy in the presence of a broad-spectrum antibiotic.

In an example, the efficacy being greater is assessed by determining the duration of progression-free survival or treatment of the disease or condition; and/or by determining a reduction in one or more symptoms of the disease or condition. For example, this determination is compared to an analogous determination in a patient suffering from the disease or condition as well as the bacterial infection and

being treated with the therapy and the antibiotic (rather than the nuclease killing of first bacteria as per the invention).

14. The method of any preceding Clause, wherein the efficacy of the therapy in the presence of the programmed nuclease is greater than the efficacy of the therapy in the presence of an antibiotic selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbapristin, teicoplanin, cephalosporin; carbapenem; fluoroquinolone; aminoglycoside; colistin; erythromycin; clindamycin; beta-lactam; macrolide; amoxicillin; azithromycin; penicillin; ceftriaxone; azithromycin; ciprofloxacin; isoniazid (INH); rifampicin (RMP); amikacin; kanamycin; capreomycin; trimethoprim; itrofurantoin; cefalexin; amoxicillin; metronidazole (MTZ); cefixime; tetracycline; and meropenem.
15. The method of any preceding Clause, wherein the first bacteria is selected from (i) *Staphylococcus aureus* that is resistant to an antibiotic selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbapristin and teicoplanin; (ii) *Pseudomonas aeruginosa* that is resistant to an antibiotic selected from cephalosporins, carbapenems, fluoroquinolones, aminoglycosides and colistin; (iii) *Klebsiella* species that is resistant to carbapenem; (iv) *Streptococcus* species that is resistant to an antibiotic selected from erythromycin, clindamycin, beta-lactam, macrolide, amoxicillin, azithromycin and penicillin; (v) *Salmonella* species that is resistant to an antibiotic selected from ceftriaxone, azithromycin and ciprofloxacin; (vi) *Shigella* species that is resistant to ciprofloxacin or azithromycin; (vii) *Mycobacterium tuberculosis* that is resistant to an antibiotic selected from Resistance to isoniazid (INH), rifampicin (RMP), fluoroquinolone, amikacin, kanamycin, capreomycin and azithromycin; (viii) *Enterococcus* species that is resistant to vancomycin; (ix) *Enterobacteriaceae* species that is resistant to an antibiotic selected from cephalosporin and carbapenem; (x) *E coli* that is resistant to an antibiotic selected from trimethoprim, itrofurantoin, cefalexin and amoxicillin; (xi) *Clostridium* species that is resistant to metronidazole (MTZ), fluoroquinolone or carbapenem; (xii) *Neisseria gonorrhoea* that is resistant to an antibiotic selected from cefixime, ceftriaxone, azithromycin and tetracycline; (xiii) *Acinetobacter baumannii* that is resistant to an antibiotic selected from beta-lactam, meropenem and carbapenem; and (xiv) *Campylobacter* species that is resistant to ciprofloxacin or azithromycin.

16. The method of any preceding Clause, wherein the treatment of the infection treats or prevents in the subject a condition selected from vaginosis, meningitis, pneumonia, urinary tract infection, cystitis, nephritis, gastroenteritis, a skin infection, impetigo, erysipelas, dental infection and cellulitis.

17. The method of any preceding Clause, wherein the treatment of the infection treats or prevents septicemia or sepsis in the subject.

In an example, the infection is a bloodstream infection.

18. The method of any preceding Clause, wherein the further disease or condition is a cancer; autoimmune disease or condition; viral infection or GI tract disease or condition. In an example, the cancer is metastatic. In an example, the cancer is melanoma. In an example, the cancer is a solid tumour with mismatch repair deficiency or microsatellite instability. In an example, the cancer is NSCLC. In an example, the cancer is

HNSCC. In an example, the cancer is Hodgkin's lymphoma. In an example, the cancer is urothelial cancer. In an example, the cancer is lung cancer. In an example, the cancer is head and neck cancer. In an example, the cancer is head cancer. In an example, the cancer is neck cancer.

In an example, the viral infection is a HIV, CMV or RSV infection.

19. The method of any preceding Clause, wherein the subject comprises bacteria (second bacteria) of one or more strains or species that are different to the first strain or species, wherein the genomes of the second bacteria do not comprise the target site, wherein the genomes of the second bacteria are not cut by the programmed nuclease in the subject, whereby second bacteria survive in the presence of the programmed nuclease in the patient; and wherein the therapy is efficacious in the presence of the second bacteria.

20. The method of Clause 19, wherein reduction in the second bacteria in patients (eg, in the gut microbiome) is associated with reduced efficacy of the therapy.

Optionally, the therapy is efficacious in the presence of the second bacteria in the gut of the subject.

Optionally, the first and/or second bacteria are present in the gut of the subject immediately prior to carrying out the method.

Optionally, the first and/or second bacteria are present in the blood of the subject immediately prior to carrying out the method.

Optionally, the first bacteria are present in the blood of the subject and the second bacteria are present in the gut of the subject immediately prior to carrying out the method.

Optionally, the first bacteria are present in the gut of the subject and the second bacteria are present in the blood of the subject immediately prior to carrying out the method.

Optionally, first bacteria in the blood of the subject is killed.

Optionally, the bacteria are gram positive bacteria. Optionally, the bacteria are gram negative bacteria.

Optionally, the first and second bacteria are capable of being killed by the same antibiotic. Optionally, the method does not comprise administering the antibiotic to the subject.

In an example, the antibiotic is selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbapristin; teicoplanin; cephalosporin; carbapenem; fluoroquinolone; aminoglycoside; colistin; erythromycin; clindamycin; beta-lactam; macrolide; amoxicillin; azithromycin; penicillin; ceftriaxone; azithromycin; ciprofloxacin; isoniazid (INH); rifampicin (RMP); amikacin; kanamycin; capreomycin; trimethoprim; itrofurantoin; cefalexin; amoxicillin; metronidazole (MTZ); cefixime; tetracycline; and meropenem. In an example, the antibiotic is selected from Aminoglycosides, Ampicillin, Amoxicillin, Amoxicillin or clavulanic acid, Carbapenems (e.g. imipenem), Piperacillin or tazobactam, Quinolones (e.g. ciprofloxacin), Tetracyclines, Chloramphenicol, Ticarcillin, Trimethoprim or sulfamethoxazole, penicillin, streptomycin, oxytetracycline and potentiated sulfonamides. In an example, the first bacteria are resistant to an antibiotic selected from Aminoglycosides, Ampicillin, Amoxicillin, Amoxicillin or clavulanic acid, Carbapenems (e.g. imipenem), Piperacillin or tazobactam, Quinolones (e.g. ciprofloxacin), Tetracyclines, Chloramphenicol, Ticarcillin, Trimethoprim or sulfamethoxazole, penicillin, streptomycin, oxytetracycline and potentiated sulfonamides. In an alternative, the antibiotic is selected from a beta-lactam, fluoroquinolone and macrolide.

Optionally, the first and second bacteria are bacteria of the same species, but are different strains of the species.

Optionally, the first and second bacteria are bacteria of the same genus, but are bacteria of different species of the genus.

Optionally, the first and second bacteria are bacteria of the same family, but are bacteria of different genera of the family.

Optionally, the first and second bacteria are gram positive bacteria.

Optionally, the first and second bacteria are gram-negative bacteria.

Optionally, the therapy is efficacious in the presence of the second bacteria.

Optionally, reduction in the second bacteria in patients is associated with reduced efficacy of the therapy. Optionally, reduction in the second bacteria in patients reduces efficacy of the therapy.

Optionally, the presence of the second bacteria in patients is associated with enhanced efficacy of the therapy. Optionally, the presence of the second bacteria in patients enhances efficacy of the therapy. For example, enhanced efficiency is efficiency compared to therapy in the absence or a reduced presence of the second bacteria, such as in the presence of an antibiotic that kills the second bacteria.

In an example, the therapy is efficacious in the presence of the second bacteria, wherein the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the progression of the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein disease-free progression of the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the duration of the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the severity of the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the disease or condition (or a symptom thereof) is reduced for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 21 or 28 days or for at least 1, 2, 3, 4 5, 6 or 12 months in the patient by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the disease or condition (or a symptom thereof) is undetectable for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 21 or 28 days or for at least 1, 2, 3, 4 5, 6 or 12 months in the patient by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the disease or condition (or a symptom thereof) is undetectable for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 21 or 28 days or for at least 1, 2, 3, 4 5, 6 or 12 months in the patient by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%.

21. The method of Clause 19 or 20, wherein the second bacteria are selected from the group consisting of *Akkermansia*, *Alistipes*, *Bacteroides*, *Barnesiella*, *Bifidobacterium*, *Clostridium*, *Collinsella*, *Enterococcus*, *Fusobacterium*, *Lactobacillus*, *Propionibacterium*,

Ruminococcus, *Segmented filamentous* bacteria (SFB); *Veillonella*, *Prevotella*, *Escherichia* and *Streptococcus* bacteria.

In an example the second bacteria that produce short chain fatty acids (eg, butyrate-producing bacteria). In particular aspects, the species of bacteria produce butyrate. For example, the second bacteria are Clostridiales. The Clostridiales bacteria may be substantially or include bacteria in spore form. In particular aspects, the second bacteria are of the family Ruminococcaceae, Christensenellaceae, Clostridiaceae or Coriobacteriaceae. In some embodiments, the Clostridiales (eg, *Clostridium*) bacteria comprise a first family and a second family. In some embodiments, the first family is selected from the group consisting of Ruminococcaceae, Christensenellaceae, Clostridiaceae and Coriobacteriaceae, and the second family is not identical to the first family. In an example, the second bacteria are *Faecalibacterium prausnitzii*, *Ruminococcus albus*, *Ruminococcus bromii*, *Ruminococcus callidus*, *Ruminococcus flavefaciens*, *Ruminococcus champanellensis*, *Ruminococcus faecis*, *Ruminococcus gnavus*, *Ruminococcus gnavus*, *Ruminococcus hanseni*, *Ruminococcus hydrogenotrophicus*, *Ruminococcus lactaris*, *Ruminococcus luti*, *Ruminococcus obeum*, *Ruminococcus palustris*, *Ruminococcus pasteurii*, *Ruminococcus productus*, *Ruminococcus schinkii*, *Ruminococcus torques*, *Subdoligranulum variabile*, *Butyrivibrio fibrisolvens*, *Roseburia intestinalis*, *Anaerostipes caccae*, *Blautia obeum*, *Eubacterium nodatum* or *Eubacterium oxidoreducens*. In particular aspects, the second bacteria are *Faecalibacterium prausnitzii*. In an example the second bacteria are Firmicutes.

In certain embodiments, the first bacteria are Bacteroidia or Prevotellaceae, eg, Bacteroidetes or *Bacteroides*.

In an embodiment, the treatment results in or maintains a microbiome (eg, gut and/or blood microbiome) of the subject, which is beneficial for the immune checkpoint inhibition or other therapy. In an example, the microbiome comprises a high relative abundance of one or more bacterial species from the phylum Firmicutes, class Clostridia, order Clostridiales, family Ruminococcaceae, genus *Ruminococcus*, genus *Hydrogenoanaerobacterium*, genus *Faecalibacterium*, phylum Actinobacteria, class Coriobacteriia, order Coriobacteriales, family Coriobacteriaceae, domain Archaea, phylum Cyanobacteria, phylum Euryarchaeota or family Christensenellaceae. Additionally or alternatively, the microbiome comprises a low relative abundance of bacteria from the genus *Dialister*, family Veillonellaceae, phylum Bacteroidetes, class Bacteroida, order Bacteroidales or family Prevotellaceae. Accordingly, a favorable microbial profile would have a higher relative abundance of one or more bacterial species from the phylum Firmicutes, class Clostridia, order Clostridiales, family Ruminococcaceae, genus *Ruminococcus*, genus *Hydrogenoanaerobacterium*, phylum Actinobacteria, class Coriobacteria, order Coriobacteriales, family Coriobacteriaceae, domain Archaea, phylum Cyanobacteria, phylum Euryarchaeota or family Christensenellaceae, and/or has a decreased abundance of one or more bacterial species from genus *Dialister*, family Veillonellaceae, phylum Bacteroidetes, class Bacteroida, order Bacteroidales and/or family Prevotellaceae.

For example, the microbiome comprises a higher relative abundance of Firmicutes compared to Bacteroidetes, Bacteroida, Bacteroidales or Prevotellaceae. For example, the microbiome comprises a higher relative abundance of Firmicutes compared to Bacteroidetes, Bacteroida, Bacteroidales and Prevotellaceae.

Optionally, the second bacteria are selected from the group consisting of *Akkermansia muciniphila*; *Alistipes shahii*; *Bacteroides fragilis*; *Bacteroides uniformis*; *Barnesiella intestinihominis*; *Bacteroides dorei*; *Bifidobacterium adolescentis*; *Bifidobacterium breve*; *Bifidobacterium longum*; *Clostridium orbiscindens*; *Clostridium novyi*; *Clostridium perfringens*; *Collinsella aerofaciens*; *Enterococcus hirae*; *Fusobacterium nucleatum*; *Lactobacillus casei* Shirota; *L. casei* AO47; *Lactobacillus rhamnosus*; *Propionibacterium granulosum*; *Ruminococcus gnavus*; *Segmented filamentous bacteria* (SFB); *Veillonella*; *Lactobacilli*; *Bacteroides*; *Clostridia*; *Prevotella*; *E. coli* Nissle; *Lactobacillus plantarum*; *Lactobacillus delbrueckii* (eg, subsp. *Bulgarius*); *Lactobacillus paracasei*; *Lactobacillus acidophilus*; *Bifidobacterium infantis*; and *Streptococcus salivarius* (eg, subsp. *Thermophilus*). See “The microbiome in cancer immunotherapy: Diagnostic tools and therapeutic strategies”; Lawrence Zitvogel et al; Science 23 Mar. 2018: Vol. 359, Issue 6382, pp. 1366-1370; DOI: 10.1126/science.aar6918.

In an example, the second bacteria are commensal bacteria in humans.

In an example, the first bacteria are comprised by gut microbiota, skin microbiota, oral cavity microbiota, throat microbiota, hair microbiota, armpit microbiota, vaginal microbiota, rectal microbiota, anal microbiota, ocular microbiota, nasal microbiota, tongue microbiota, lung microbiota, liver microbiota, kidney microbiota, genital microbiota, penile microbiota, scrotal microbiota, mammary gland microbiota, ear microbiota, urethra microbiota, labial microbiota, organ microbiota or dental microbiota.

In an example, the second bacteria are comprised by gut microbiota, skin microbiota, oral cavity microbiota, throat microbiota, hair microbiota, armpit microbiota, vaginal microbiota, rectal microbiota, anal microbiota, ocular microbiota, nasal microbiota, tongue microbiota, lung microbiota, liver microbiota, kidney microbiota, genital microbiota, penile microbiota, scrotal microbiota, mammary gland microbiota, ear microbiota, urethra microbiota, labial microbiota, organ microbiota or dental microbiota.

In an example, the first and/or second bacteria are blood-borne bacteria.

22. The method of any preceding Clause, wherein the first bacteria are selected from the group consisting *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Helicobacter*, *Legionella*, *Heamophilus*, *Ghomorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria.

H pylori has been implicated in gastric cancer and gastric ulcers. Thus, in an example, the first bacteria are *H pylori* and optionally the disease is a cancer, such as gastric cancer. In an embodiment, the therapy is chemotherapy or therapy with an immune checkpoint inhibitor (eg, an antibody). In an example, the first bacteria are *H pylori* and the disease is gastric ulcer(s). In an embodiment, triple therapy for gastric ulcers is administered to the subject.

In an example, the first bacteria are Gram-negative bacteria and optionally the infection is a blood infection. In an example, the first bacteria are selected from *E. coli*, *P. aeruginosa* and *K pneumoniae*, and optionally the infection is a blood infection.

23. The method of Clause 22, wherein the first bacteria are selected from the group consisting of *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Helicobacter pylori*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* bacteria.

In an example, the subject has been administered an immunosuppressant drug, or is on a course of an immunosuppressant drug, eg, a steroid, such as a corticosteroid.

24. A programmable nuclease for use in the method of any preceding Clause.

25. The method or nuclease of any preceding Clause, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.

26. A CRISPR/Cas system comprising a nuclease according to Clause 24 or 25 for use in the method of any one of Clauses 1 to 23, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs (gRNAs) or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of first bacteria.

27. A guide RNA or a DNA encoding a guide RNA for use in the system of Clause 26.

28. A guide RNA or a DNA encoding a guide RNA for use in the method of treating a pathogenic bacterial infection according to any one of Clauses 1 to 23, wherein the guide RNA is capable of programming the nuclease, wherein the nuclease is a Cas nuclease (eg, a Cas9, Cas3, Cas13, CasX, CasY or Cpf1 nuclease).

29. A nucleic acid vector comprising the guide RNA or DNA recited in any one of Clauses 26 to 28.

30. A nucleic acid vector encoding the nuclease of Clause 24 or 25 and optionally the guide RNA of Clause 29.

31. The vector of Clause 29 or 30 wherein the vector is a phage, phagemid plasmid (eg, conjugative plasmid) or transposon.

The phage are capable of infecting first bacteria and the phagemids are capable of producing such phage in the presence of a helper phage.

32. A pharmaceutical composition comprising a first nucleic acid vector (or a plurality thereof) encoding the nuclease of Clause 24 or 25 and a second nucleic acid vector (or a plurality thereof) encoding the guide RNA of Clause 29, the composition further comprising a pharmaceutically acceptable diluent, excipient or carrier.

33. A pharmaceutical composition comprising the CRISPR/Cas system of claim 26 and a pharmaceutically acceptable diluent, excipient or carrier.

34. A pharmaceutical composition comprising the vector of claim 31 and a pharmaceutically acceptable diluent, excipient or carrier.

Preventing a disease or condition herein may, for example, be reducing the risk of the disease or condition in the subject or patient.

In an alternative, instead of first bacteria, the infection is caused by first archaea and in this embodiment all of the features of the method and other configurations of the invention relating to killing first bacteria instead relate mutatis mutandis to killing first archaea.

In an embodiment, the method comprises carrying out the method of treating an acute microbial infection as described herein, and thus features of that method as described herein are combinable with the present method of treating a pathogenic bacterial infection (ie, where the pathogenic bacterial infection is the acute microbial infection in the first method). In an embodiment, the method comprises carrying out the method of durably treating a microbial infection as described herein, and thus features of that method as described herein are combinable with the present method of treating a pathogenic bacterial infection (ie, where the

pathogenic bacterial infection is the microbial infection in the first method). Any of the optional features of the first method herein may apply mutatis mutandis to the present method of treating a pathogenic bacterial infection.

Aspects:—

Thus, the invention provides the following Aspects, which are optional features of Clauses above:—

1. The method of any one of Clauses 1-23, wherein the infection is reduced at least 100-fold by the first 30 minutes of carrying out step (b). Optionally, the infection is reduced at least 1000-fold by the first 30 minutes of carrying out step (b). Optionally, the reduction in infection persists for 30 minutes immediately after the first 30 minutes of carrying out step (b). For example, the reduction can be assessed by determining the difference in the number of bacteria of the first species or strain in (i) a sample taken from the subject (eg, a blood sample) immediately before commencement of the method and (ii) a sample (of the same type as the sample of (i), eg, a blood sample) taken from the subject at 30 minutes of the treatment. For example, the samples may be assessed for the difference in colony forming units (CFU)/ml sample, eg, when the samples have been plated on agar in respective petri dishes and incubated under identical conditions. Another example may use microscopic counting of bacteria in samples, or other routine methods known to the skilled addressee.
 2. The method of any one of Clauses 1-23, wherein blood infection of the subject by the first bacteria is reduced at least 100- or 1000-fold by the first 30 minutes of carrying out step (b).
 3. The method of Aspect 2, wherein the blood is infected with from 10^5 to 10^{12} (eg, 10^7 to 10^{12}) CFU/ml of the first bacteria immediately before the treatment.
 4. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the method comprises administering to the subject a nucleic acid (eg, a RNA) and nuclease, wherein the nucleic acid complexes with the nuclease to program the nuclease to cut the target site in the first bacteria comprised by the subject.
 5. The method of Aspect 4, wherein the nuclease is administered simultaneously or sequentially with the nucleic acid to the subject.
 6. The method of Aspect 4, wherein the subject comprises the nuclease prior to administration of the nucleic acid to the subject.
 7. The method of any one of Aspects 4 to 6, wherein a plurality phage are administered to the subject, wherein each phage comprises a copy of the nucleic acid, wherein the phage infect first bacteria comprised by the subject to deliver thereto the nucleic acid.
 8. The method of Aspect 7, wherein the ratio of administered phage: first bacteria comprised by the subject is from 10 to 150. For example, the ratio is from 10 to 100, ie, a multiplicity of infection (MOI) of from 10 to 100.
- The ratio can be determined, for example, using a sample (eg, a blood or gut sample) from a human or animal subject immediately before the treatment and determining the number of bacteria per ml of blood or gut sample. The amount of phage to be administered can then be worked out according to the determination using the sample.
9. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the infection is an infection of the lungs, brain, skin, abdomen or urinary tract.
 10. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the subject has undergone

surgery, is on an immunosuppressant medication, suffering from burns, suffering from diabetes, suffering from cancer or is suffering from a chronic disease.

11. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the subject is a human over 65 years of age or is a paediatric patient.
12. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the method treats or prevents sepsis in the subject.
13. The method of Clause 12, wherein at the start of the treatment, the subject (eg, a human) has a temperature of $<36^\circ\text{C}$. or $>38^\circ\text{C}$.; a heart rate of $>90/\text{min}$, a respiratory rate of >20 breaths/min or $\text{PaCO}_2 < 4.3\text{ kPa}$; and white blood cell count of $<4000/\text{mm}^3$ or $>12,000/\text{mm}^3$
14. The method of Clause 12 or 13, wherein at the start of the treatment, the subject (eg, a human) has presence of two or more of the following: abnormal body temperature, abnormal heart rate, abnormal respiratory rate, abnormal blood gas and abnormal white blood cell count.

Immune Checkpoint Modulation

Immune checkpoints of the invention either turn up a signal (e.g., co-stimulatory molecules) or turn down a signal. Inhibitory immune checkpoint molecules that may be targeted by immune checkpoint modulation in the invention include adenosine A2A receptor (A2AR), B7-H3 (also known as CD276), B and T lymphocyte attenuator (BTLA), cytotoxic T-lymphocyte-associated protein 4 (CTLA-4, also known as CD 152), indoleamine 2,3-dioxygenase (IDO), killer-cell immunoglobulin (KIR), lymphocyte activation gene-3 (LAG3), programmed death 1 (PD-1), T-cell immunoglobulin domain and mucin domain 3 (TIM-3) and V-domain Ig suppressor of T cell activation (VISTA). In particular, the immune checkpoint inhibitors target the PD-1 axis and/or CTLA-4.

The immune checkpoint inhibitors may be drugs such as small molecules, recombinant forms of ligand or receptors, or, antibodies, such as human antibodies (e.g., WO2015016718; Pardoll, Nat Rev Cancer, 12(4): 252-64, 2012; both incorporated herein by reference). Known inhibitors of the immune checkpoint proteins or analogs thereof may be used, in particular chimerised, humanised or human forms of antibodies may be used. As the skilled person will know, alternative and/or equivalent names may be in use for certain antibodies mentioned in the present disclosure. Such alternative and/or equivalent names are interchangeable in the context of the present invention. For example it is known that lambrolizumab is also known under the alternative and equivalent names MK-3475 and pembrolizumab.

It is contemplated that any of the immune checkpoint inhibitors that are known in the art to stimulate immune responses may be used. This includes inhibitors that directly or indirectly stimulate or enhance antigen-specific T-lymphocytes. These immune checkpoint inhibitors include, without limitation, agents targeting immune checkpoint proteins and pathways involving PD-L2, LAG3, BTLA, B7H4 and TIM3. For example, LAG3 inhibitors known in the art include soluble LAG3 (IMP321, or LAG3-Ig disclosed in WO2009044273) as well as mouse or humanized antibodies blocking human LAG3 (e.g., IMP701 disclosed in WO2008132601), or fully human antibodies blocking human LAG3 (such as disclosed in EP 2320940). Another example is provided by the use of blocking agents towards BTLA, including without limitation antibodies blocking human BTLA interaction with its ligand (such as 4C7 disclosed in WO2011014438). Yet another example is pro-

vided by the use of agents neutralizing B7H4 including without limitation antibodies to human B7H4 (disclosed in WO 2013025779, and in WO2013067492) or soluble recombinant forms of B7H4 (such as disclosed in US20120177645). Yet another example is provided by agents neutralizing B7-H3, including without limitation antibodies neutralizing human B7-H3 (e.g. MGA271 disclosed as BRCA84D and derivatives in US 20120294796). Yet another example is provided by agents targeting TIM3, including without limitation antibodies targeting human TIM3 (e.g. as disclosed in WO 2013006490 A2 or the anti-human TIM3, blocking antibody F38-2E2 disclosed by Jones et al, J Exp Med. 2008; 205(12):2763-79).

A. PD-1 Axis Antagonists

T cell dysfunction or anergy occurs concurrently with an induced and sustained expression of the inhibitory receptor, programmed death 1 polypeptide (PD-1). Thus, therapeutic targeting of PD-1 and other molecules which signal through interactions with PD-1, such as programmed death ligand 1 (PD-L1) and programmed death ligand 2 (PD-L2) is provided herein. PD-L1 is overexpressed in many cancers and is often associated with poor prognosis (Okazaki T et al, Intern. Immun 2007 19(7):813). Thus, improved methods of treating cancer by inhibiting the PD-L1/PD-1 interaction in combination with modulating the microbiome is provided herein.

For example, PD-1 axis binding antagonists include a PD-1 binding antagonist, a PD-L1 binding antagonist and a PD-L2 binding antagonist. Alternative names for "PD-1" include CD279 and SLEB2. Alternative names for "PD-L1" include B7-H1, B7-4, CD274, and B7-H. Alternative names for "PD-L2" include B7-DC, Btcd, and CD273. In some embodiments, PD-1, PD-L1, and PD-L2 are human PD-1, PD-L1 and PD-L2.

In some embodiments, the PD-1 binding antagonist is a molecule that inhibits the binding of PD-1 to its ligand binding partners. In a specific aspect, the PD-1 ligand binding partners are PD-L1 and/or PD-L2. In another embodiment, a PD-L1 binding antagonist is a molecule that inhibits the binding of PD-L1 to its binding partners. In a specific aspect, PD-L1 binding partners are PD-1 and/or B7-1. In another embodiment, the PD-L2 binding antagonist is a molecule that inhibits the binding of PD-L2 to its binding partners. In a specific aspect, a PD-L2 binding partner is PD-1. The antagonist may be an antibody, an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide. Exemplary antibodies are described in U.S. Pat. Nos. 8,735,553, 8,354,509, and 8,008,449, all incorporated herein by reference. Other PD-1 axis antagonists for use in the methods provided herein are known in the art such as described in U.S. Patent Application No. US20140294898, US2014022021, and US20110008369, all incorporated herein by reference.

In some embodiments, the PD-1 binding antagonist is an anti-PD-1 antibody (e.g., a human antibody, a humanized antibody, or a chimeric antibody). In some embodiments, the anti-PD-1 antibody is selected from the group consisting of nivolumab, pembrolizumab, and CT-011. In some embodiments, the PD-1 binding antagonist is an immunoadhesin (e.g., an immunoadhesin comprising an extracellular or PD-1 binding portion of PDL1 or PD-L2 fused to a constant region (e.g., an Fc region of an immunoglobulin sequence). In some embodiments, the PD-1 binding antagonist is AMP-224. Nivolumab, also known as MDX-1106-04, MDX-1106, ONO-4538, BMS-936558, and OPDIVO® is an anti-PD-1 antibody described in WO2006/121168. Pembrolizumab, also known as MK-3475, Merck 3475, lambrolizumab,

KEYTRUDA® and SCH-900475, is an anti-PD-1 antibody described in WO2009/114335. CT-011, also known as hBAT or hBAT-1, is an anti-PD-1 antibody described in WO2009/101611. AMP-224, also known as B7-DCIg, is a PD-L2-Fc fusion soluble receptor described in WO2010/027827 and WO2011/066342. Additional PD-1 binding antagonists include pidilizumab, also known as CT-011, MEDI0680, also known as AMP-514, and REGN2810.

In some embodiments, the immune checkpoint inhibitor is a PD-L1 antagonist such as durvalumab, also known as MEDI4736, atezolizumab, also known as MPDL3280A, or avelumab, also known as MSB00010118C. In certain aspects, the immune checkpoint inhibitor is a PD-L2 antagonist such as rHlgM12B7. In some aspects, the immune checkpoint inhibitor is a LAG-3 antagonist such as, but not limited to, IMP321, and BMS-986016. The immune checkpoint inhibitor may be an adenosine A2a receptor (A2aR) antagonist such as PBF-509.

In some embodiments, any antibody described herein (such as an anti-PD-1 antibody, an anti-PD-L1 antibody, or an anti-PD-L2 antibody) further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic cells. In a still further specific aspect the minimal effector function results from an "effector-less Fc mutation" or aglycosylation. Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose to a hydroxy amino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxy lysine may also be used. Removal of glycosylation sites from an antibody is conveniently accomplished by altering the amino acid sequence such that one of the above-described tripeptide sequences (for N-linked glycosylation sites) is removed. The alteration may be made by substitution of an asparagine, serine or threonine residue within the glycosylation site another amino acid residue (e.g., glycine, alanine or a conservative substitution).

The antibody or antigen binding fragment thereof, may be made using methods known in the art, for example, by a process comprising culturing a host cell containing nucleic acid encoding any of the previously described anti-PD-L1, anti-PD-1, or anti-PD-L2 antibodies or antigen-binding fragment in a form suitable for expression, under conditions suitable to produce such antibody or fragment, and recovering the antibody or fragment.

B. CTLA-4

Another immune checkpoint that can be targeted in the methods provided herein is the cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), also known as CD152. The complete cDNA sequence of human CTLA-4 has the Genbank accession number L15006. CTLA-4 is found on the

surface of T cells and acts as an “off switch when bound to CD80 or CD86 on the surface of antigen-presenting cells. CTLA4 is a member of the immunoglobulin superfamily that is expressed on the surface of Helper T cells and transmits an inhibitory signal to T cells. CTLA4 is similar to the T-cell co-stimulatory protein, CD28, and both molecules bind to CD80 and CD86, also called B7-1 and B7-2 respectively, on antigen-presenting cells. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T cells and may be important to their function. T cell activation through the T cell receptor and CD28 leads to increased expression of CTLA-4, an inhibitory receptor for B7 molecules.

In some embodiments, the immune checkpoint inhibitor is an anti-CTLA-4 antibody (e.g., a human antibody, a humanized antibody, or a chimeric antibody), an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide.

Anti-human-CTLA-4 antibodies (or VH and/or VL domains derived therefrom) suitable for use in the present methods can be generated using methods well known in the art. Alternatively, art recognized anti-CTLA-4 antibodies can be used. For example, the anti-CTLA-4 antibodies disclosed in: U.S. Pat. No. 8,119,129, WO 01/14424, WO 98/42752; WO 00/37504 (CP675, 206, also known as tremelimumab; formerly ticilimumab), U.S. Pat. No. 6,207,156; Hurwitz et al, 1998; can be used in the methods disclosed herein. The teachings of each of the aforementioned publications are hereby incorporated by reference. Antibodies that compete with any of these art-recognized antibodies for binding to CTLA-4 also can be used. For example, a humanized CTLA-4 antibody is described in International Patent Application No. WO2001014424, WO2000037504, and U.S. Pat. No. 8,017,114; all incorporated herein by reference.

An exemplary anti-CTLA-4 antibody is ipilimumab (also known as 10D1, MDX-010, MDX-101, and Yervoy®) or antigen binding fragments and variants thereof (see, e.g., WO01/14424). In other embodiments, the antibody comprises the heavy and light chain CDRs or VRs of ipilimumab. Accordingly, in one embodiment, the antibody comprises the CDR1, CDR2, and CDR3 domains of the VH region of ipilimumab, and the CDR1, CDR2 and CDR3 domains of the VL region of ipilimumab. In another embodiment, the antibody competes for binding with and/or binds to the same epitope on CTLA-4 as the above-mentioned antibodies. In another embodiment, the antibody has at least about 90% variable region amino acid sequence identity with the above-mentioned antibodies (e.g., at least about 90%, 95%, or 99% variable region identity with ipilimumab).

Other molecules for modulating CTLA-4 include soluble CTLA-4 ligands and receptors such as described in U.S. Pat. Nos. 5,844,905, 5,885,796 and International Patent Application Nos. WO1995001994 and WO1998042752; all incorporated herein by reference, and immunoadhesins such as described in U.S. Pat. No. 8,329,867, incorporated herein by reference.

C. Killer Immunoglobulin-Like Receptor (KIR)

Another immune checkpoint inhibitor for use in the present invention is an anti-KIR antibody. Anti-human-KIR antibodies (or VH/VL domains derived therefrom) suitable for use in the present methods can be generated using methods well known in the art.

Alternatively, art recognized anti-KIR antibodies can be used. The anti-KIR antibody can be cross-reactive with

multiple inhibitory KIR receptors and potentiates the cytotoxicity of NK cells bearing one or more of these receptors. For example, the anti-KIR antibody may bind to each of KIR2D2DL1, KIR2DL2, and KIR2DL3, and potentiate NK cell activity by reducing, neutralizing and/or reversing inhibition of NK cell cytotoxicity mediated by any or all of these KIRs. In some aspects, the anti-KIR antibody does not bind KIR2DS4 and/or KIR2DS3. For example, monoclonal antibodies 1-7F9 (also known as IPH2101), 14F1, 1-6F1 and 1-6F5, described in WO 2006/003179, the teachings of which are hereby incorporated by reference, can be used. Antibodies that compete with any of these art-recognized antibodies for binding to KIR also can be used. Additional art-recognized anti-KIR antibodies which can be used include, for example, those disclosed in WO 2005/003168, WO 2005/009465, WO 2006/072625, WO 2006/072626, WO 2007/042573, WO 2008/084106, WO 2010/065939, WO 2012/071411 and WO 2012/160448.

An exemplary anti-KIR antibody is lirilumab (also referred to as BMS-986015 or IPH2102). In other embodiments, the anti-KIR antibody comprises the heavy and light chain complementarity determining regions (CDRs) or variable regions (VRs) of lirilumab. Accordingly, in one embodiment, the antibody comprises the CDR1, CDR2, and CDR3 domains of the heavy chain variable (VH) region of lirilumab, and the CDR1, CDR2 and CDR3 domains of the light chain variable (VL) region of lirilumab. In another embodiment, the antibody has at least about 90% variable region amino acid sequence identity with lirilumab.

Examples of cancers contemplated for treatment include lung cancer, head and neck cancer, breast cancer, pancreatic cancer, prostate cancer, renal cancer, bone cancer, testicular cancer, cervical cancer, gastrointestinal cancer, lymphomas, pre-neoplastic lesions in the lung, colon cancer, melanoma, metastatic melanoma, basal-cell skin cancer, squamous-cell skin cancer, dermatofibrosarcoma protuberans, Merkel cell carcinoma, Kaposi’s sarcoma, keratoacanthoma, spindle cell tumours, sebaceous carcinomas, microcystic adnexal carcinoma, Paget’s disease of the breast, atypical fibroxanthoma, leiomyosarcoma, and angiosarcoma, Lentigo Maligna, Lentigo Maligna Melanoma, Superficial Spreading Melanoma, Nodular Melanoma, Acral Lentiginous Melanoma, Desmoplastic Melanoma, and bladder cancer.

In some embodiments, the subject has cancer that is resistant (has been demonstrated to be resistant) to one or more anti-cancer therapies. In some embodiments, resistance to anti-cancer therapy includes recurrence of cancer or refractory cancer. Recurrence may refer to the reappearance of cancer, in the original site or a new site, after treatment. In some embodiments, resistance to anti-cancer therapy includes progression of the cancer during treatment with the anti-cancer therapy. In some embodiments, the cancer is at early stage or at late stage. The subject may have a cancer that expresses (has been shown to express e.g., in a diagnostic test) PD-L1 biomarker. In some embodiments, the patient’s cancer expresses low PD-L1 biomarker. In some embodiments, the patient’s cancer expresses high PD-L1 biomarker. The PD-L1 biomarker can be detected in the sample using a method selected from the group consisting of FACS, Western blot, ELISA, immunoprecipitation, immunohistochemistry, immunofluorescence, radioimmunoassay, dot blotting, immunodetection methods, HPLC, surface plasmon resonance, optical spectroscopy, mass spectrometry, HPLC, qPCR, RT-qPCR, multiplex qPCR or RT-qPCR, RNA-seq, microarray analysis, SAGE, MassARRAY technique, and FISH, and combinations thereof.

In some embodiments, the cancer has low levels of T cell infiltration. In some embodiments, the cancer has no detectable T cell infiltrate. In some embodiments, the cancer is a non-immunogenic cancer (e.g., non-immunogenic colorectal cancer and/or ovarian cancer).

For example, a therapeutically effective or sufficient amount of the immune checkpoint inhibitor, such as an antibody, is administered to a human will be in the range of about 0.01 to about 50 mg/kg of patient body weight whether by one or more administrations. In some embodiments, the antibody used is about 0.01 to about 45 mg/kg, about 0.01 to about 40 mg/kg, about 0.01 to about 35 mg/kg, about 0.01 to about 30 mg/kg, about 0.01 to about 25 mg/kg, about 0.01 to about 20 mg/kg, about 0.01 to about 15 mg/kg, about 0.01 to about 10 mg/kg, about 0.01 to about 5 mg/kg, or about 0.01 to about 1 mg/kg administered daily, for example. In some embodiments, the antibody is administered at 15 mg/kg. However, other dosage regimens may be useful. In one embodiment, an anti-PD-L1 antibody described herein is administered to a human at a dose of about 100 mg, about 200 mg, about 300 mg, about 400 mg, about 500 mg, about 600 mg, about 700 mg, about 800 mg, about 900 mg, about 1000 mg, about 1100 mg, about 1200 mg, about 1300 mg or about 1400 mg on day 1 of 21-day cycles. The dose may be administered as a single dose or as multiple doses (e.g., 2 or 3 doses), such as infusions. The progress of this therapy is easily monitored by conventional techniques.

Anti-Cancer and Other Therapies

In some embodiments, the immune checkpoint inhibitor may be administered in combination with at least one additional therapeutic. The additional therapy may be a cancer therapy such as radiation therapy, surgery, chemotherapy, gene therapy, DNA therapy, viral therapy, RNA therapy, immunotherapy, bone marrow transplantation, nanotherapy, monoclonal antibody therapy, or a combination of the foregoing. The additional therapy may be in the form of adjuvant or neoadjuvant therapy.

In an example, the therapy of the cancer (whether with or without administration of an immune checkpoint inhibitor) or any other disease (eg, viral infection or autoimmune disease) may radiation therapy, surgery, chemotherapy, gene therapy, DNA therapy, viral therapy, RNA therapy, immunotherapy, bone marrow transplantation, nanotherapy or monoclonal antibody therapy. The therapy may be a combination of the foregoing. An additional therapy may be administered

In some embodiments, the therapy (or the additional cancer therapy) is the administration of a small molecule enzymatic inhibitor or anti-metastatic agent. In some embodiments, the additional therapy is the administration of side-effect limiting agents (e.g., agents intended to lessen the occurrence and/or severity of side effects of treatment, such as anti-nausea agents, etc.).

In some embodiments, the therapy (or the additional cancer therapy) is radiation therapy. In some embodiments, the therapy (or the additional cancer therapy) is surgery. In some embodiments, the therapy (or the additional cancer therapy) is a combination of radiation therapy and surgery. In some embodiments, the therapy (or the additional cancer therapy) is gamma irradiation. In some embodiments, the therapy (or the additional cancer therapy) is therapy targeting PBK/AKT/mTOR pathway, HSP90 inhibitor, tubulin inhibitor, apoptosis inhibitor, and/or chemopreventative agent. The therapy (or the additional cancer therapy) may be one or more of the chemotherapeutic agents known in the art.

Administration of any compound or therapy of the present embodiments to a patient will follow general protocols for the administration of such compounds, taking into account the toxicity, if any, of the agents. Therefore, in some embodiments there is a step of monitoring toxicity that is attributable to combination therapy.

The therapy can comprise or consist of administration to the subject of any of the following:—

1. Chemotherapy

A wide variety of chemotherapeutic agents may be used in accordance with the present embodiments. The term “chemotherapy” refers to the use of drugs to treat cancer. A “chemotherapeutic agent” is used to connote a compound or composition that is administered in the treatment of cancer. These agents or drugs are categorized by their mode of activity within a cell, for example, whether and at what stage they affect the cell cycle. Alternatively, an agent may be characterized based on its ability to directly cross-link DNA, to intercalate into DNA, or to induce chromosomal and mitotic aberrations by affecting nucleic acid synthesis.

Examples of chemotherapeutic agents include alkylating agents, such as thiotepa and cyclophosphamide; alkyl sulfonates, such as busulfan, improsulfan, and piposulfan; aziridines, such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines, including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide, and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatins; callistatin; CC-1065 (eg, its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (eg, the synthetic analogues, KW-2189 and CB 1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards, such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, and uracil mustard; nitrosoureas, such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics, such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gammall and calicheamicin omegall); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores, aclacinomysins, actinomycin, authrarnycin, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (eg, morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, such as mitomycin C, mycophenolic acid, nogalarnycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, and zorubicin; anti-metabolites, such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues, such as denopoterin, pteropterin, and trimetrexate; purine analogs, such as fludarabine, 6-mercaptopurine, thiamiprine, and thioguanine; pyrimidine analogs, such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, encitabine, and floxuridine; androgens, such as calusterone, dromostanolone propionate, epiostanol, mepitiostane, and testolactone; anti-adrenals, such as mitotane and trilostane; folic acid replenisher, such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic

acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids, such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK polysaccharide complex; razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (eg, T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; taxoids, e.g., paclitaxel and docetaxel gemcitabine; 6-thioguanine; mercaptopurine; platinum coordination complexes, such as cisplatin, oxaliplatin, and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (e.g., CPT-11); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids, such as retinoic acid; capecitabine; carboplatin, procarbazine, plicomycin, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatinum, and pharmaceutically acceptable salts, acids, or derivatives of any of the above

2. Radiotherapy

Other factors that cause DNA damage and have been used extensively include what are commonly known as γ -rays, X-rays, and/or the directed delivery of radioisotopes to tumour cells. Other forms of DNA damaging factors are also contemplated, such as microwaves, proton beam irradiation (U.S. Pat. Nos. 5,760,395 and 4,870,287), and UV-irradiation. It is most likely that all of these factors affect a broad range of damage on DNA, on the precursors of DNA, on the replication and repair of DNA, and on the assembly and maintenance of chromosomes. Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 weeks), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

3. Immunotherapy

The skilled artisan will understand that immunotherapies may be used in combination or in conjunction with the methods described herein. In the context of cancer treatment, immunotherapeutics generally rely on the use of immune effector cells and molecules to target and destroy cancer cells. Rituximab (RITUXAN®) is an example of an immunotherapy. The immune effector may be, for example, an antibody specific for a marker on the surface of a tumour cell. The antibody alone may serve as an effector of therapy or it may recruit other cells to actually effect cell killing. The antibody also may be conjugated to a drug or toxin (chemo therapeutic, radionuclide, ricin A chain, cholera toxin, pertussis toxin, etc.) and serve as a targeting agent. Alternatively, the effector may be a lymphocyte carrying a surface molecule that interacts, either directly or indirectly, with a tumour cell target. Various effector cells include cytotoxic T cells and NK cells.

In an example, the immunotherapy comprises adoptive cell therapy, such as CAR-T administration, eg, anti-CD19 or CD20 CAR-T administration.

In an example, the immunotherapy comprises or consists of administration of an IL-2 (eg, a truncated IL-2 or pegylated IL-2 or Fc-fused IL-2).

Antibody-drug conjugates have emerged as a breakthrough approach to the development of cancer therapeutics. Antibody-drug conjugates (ADCs) comprise monoclonal antibodies (MAbs) that are covalently linked to cell-killing drugs. This approach combines the high specificity of MAbs against their antigen targets with highly potent cytotoxic drugs, resulting in "armed" MAbs that deliver the payload (drug) to tumour cells with enriched levels of the antigen. Targeted delivery of the drug also minimizes its exposure in normal tissues, resulting in decreased toxicity and improved therapeutic index. The approval of two ADC drugs, ADCE-TRIS® (brentuximab vedotin) in 2011 and KADCYLA® (trastuzumab emtansine or T-DM1) in 2013 by FDA validated the approach. There are currently more than 30 ADC drug candidates in various stages of clinical trials for cancer treatment. As antibody engineering and linker-payload optimization are becoming more and more mature, the discovery and development of new ADCs are increasingly dependent on the identification and validation of new targets that are suitable to this approach and the generation of targeting MAbs. Two criteria for ADC targets are upregulated/high levels of expression in tumour cells and robust internalization.

In one aspect of immunotherapy, the tumour cell must bear some marker that is amenable to targeting, i.e., is not present on the majority of other cells. Many tumour markers exist and any of these may be suitable for targeting in the context of the present embodiments. Common tumour markers include CD20, carcinoembryonic antigen, tyrosinase (p97), gp68, TAG-72, HMFG, Sialyl Lewis Antigen, MucA, MucB, PLAP, laminin receptor, erb B, and pi 55. An alternative aspect of immunotherapy is to combine anticancer effects with immune stimulatory effects. Immune stimulating molecules also exist including: cytokines, such as IL-2, IL-4, IL-12, GM-CSF, gamma-IFN, chemokines, such as MIP-1, MCP-1, IL-8, and growth factors, such as FLT3 ligand.

4. Surgery

The cancer or other disease or condition may be treated by surgery in the invention.

Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative, and palliative surgery. Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed and may be used in conjunction with other therapies, such as the treatment of the present embodiments, chemotherapy, radiotherapy, hormonal therapy, gene therapy, immunotherapy, and/or alternative therapies. Tumour resection refers to physical removal of at least part of a tumour. In addition to tumour resection, treatment by surgery includes laser surgery, cryosurgery, electro surgery, and microscopically-controlled surgery (Mohs' surgery).

Upon excision of part or all of cancerous cells, tissue, or tumour, a cavity may be formed in the body. Treatment may be accomplished by perfusion, direct injection, or local application of the area with an additional anti-cancer therapy. Such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. These treatments may be of varying dosages as well.

5. Bacterial Transplants

In an embodiment, the therapy comprises administering to the subject a bacterial transplant, eg, a faecal microbial transplant, comprising defined bacteria. For example, the transplant is any composition disclosed in WO2018064165, the disclosure of which (especially the compositions therein)

are incorporated herein by reference in its entirety for possible application in the present invention. For example, the transplant is according to any of the following Paragraphs (tables and sequence numbers referring to the tables and sequences in WO2018064165, which are explicitly incorporated herein for possible use in the Claims):—

1. A composition comprising at least one isolated or purified population of bacteria belonging to one or more of the families Ruminococcaceae, Clostridiaceae, Lachnospiraceae, Micrococcaceae, and/or Veilonellaceae.

2. A composition comprising at least two isolated or purified populations of bacteria belonging to one or more of the families Ruminococcaceae, Clostridiaceae, Lachnospiraceae, Micrococcaceae, and/or Veilonellaceae.

3. The composition of Paragraph 1 or Paragraph 2, wherein each of the populations of bacteria is present in the composition at a concentration of at least 10^3 CFU.

4. The composition of Paragraph 1 or Paragraph 2, wherein the composition is a live bacterial product or a live biotherapeutic product.

5. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria are provided as bacterial spores.

6. The composition of Paragraph 1 or Paragraph 2, wherein the at least one population of bacteria or the at least two isolated or purified populations of bacteria belong to Clostridiales Family XII and/or Clostridiales Family XIII.

7. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria belong to the family Ruminococcaceae and/or of the family Clostridiaceae.

8. The composition of Paragraph 1 or Paragraph 2, wherein the population of bacteria belonging to the family Ruminococcaceae is further defined as a population of bacteria belonging to the genus *Ruminococcus*.

9. The composition of Paragraph 8, wherein the population of bacteria belonging to the genus *Ruminococcus* is further defined as a population of bacteria belonging to the species *Ruminococcus bromii*.

10. The composition of Paragraph 1 or Paragraph 2, wherein the population of bacteria belonging to the family Ruminococcaceae is further defined as a population of bacteria belonging to the genus *Faecalibacterium*.

11. The composition of Paragraph 10, wherein the population of bacteria belonging to the genus *Faecalibacterium* is further defined as a population of bacteria belonging to the species *Faecalibacterium prausnitzii*.

12. The composition of Paragraph 1 or Paragraph 2, wherein the population of bacteria belonging to the family Micrococcaceae is further defined as a population of bacteria belonging to the genus *Rothia*.

13. The composition of Paragraph 1 or Paragraph 2, wherein the composition further comprises a population of bacteria belonging to the species *Porphyromonas pasteri*, the species *Clostridium hungatei*, the species *Phascolarctobacterium faecium*, the genus *Peptomphilus*, and/or the class Mollicutes.

14. The composition of Paragraph 1 or Paragraph 2, wherein the composition is essentially free of populations of bacteria belonging to the order Bacteroidales.

15. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria belongs to one or more of the species, subspecies or bacterial

strains selected from the group consisting of the species in Table 1 with an enrichment index (ei) greater than 0.5.

16. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria are selected from the group consisting of the species in Table 1 with an “ei” equal to 1.

17. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria comprise a 16S ribosomal RNA (rRNA) nucleotide sequence that is at least 90% identical (eg, at least 91, 92, 93, 94, 95, 96, 97, 98 or 99% identical) to the 16S rRNA nucleotide sequence of bacteria identified by NCBI Taxonomy IDs selected from the group consisting of NCBI Taxonomy ID: 717959, 587, 758823, 649756, 44749, 671218, 1264, 1122135, 853, 484018, 46503, 54565, 290052, 216931, 575978, 433321, 1796646, 213810, 228924, 290054, 1509, 1462919, 29375, 337097, 1298596, 487174, 642492, 1735, 1297424, 742766, 46680, 132925, 411467, 1318465, 1852367, 1841857, 169679, 1175296, 259063, 172901, 39488, 57172, 28118, 166486, 28133, 1529, 694434, 1007096, 84030, 56774, 102148, 626947, 216933, 1348613, 1472417, 100176, 824, 1471761, 1297617, 288966, 1317125, 28197, 358743, 264639, 1265, 1335, 66219, 69473, 115117, 341220, 1732, 873513, 396504, 1796619, 45851, 2741, 105841, 86332, 1349822, 84037, 180311, 54291, 1217282, 762984, 1185412, 154046, 663278, 1543, 398512, 69825, 1841867, 1535, 1510, 84026, 1502, 1619234, 39497, 1544, 29343, 649762, 332095, 536633, 1033731, 574930, 742818, 177412, 1121308, 419208, 1673717, 55779, 28117, 626937, 180332, 1776382, 40519, 34062, 40518, 74426, 1216062, 293826, 850, 645466, 474960, 36835, 115544, 1515, 88431, 216932, 1417852, 39492, 1583, 420247, 118967, 169435, 37658, 138595, 31971, 100886, 1197717, 234908, 537007, 319644, 168384, 915173, 95159, 1816678, 626940, 501571, 1796620, 888727, 1147123, 376806, 1274356, 1267, 39495, 404403, 1348, 253314, 258515, 33033, 1118061, 357276, 214851, 320502, 217731, 246787, 29371, 649764, 901, 29374, 33043, 39778, 682400, 871665, 160404, 745368, 408, 1584, 333367, 47246, 1096246, 53342, 438033, 351091, 1796622, 1776384, 817, 48256, 720554, 500632, 36849, 301302, 879970, 655811, 264463, 1532, 285, 995, 242750, 29539, 1432052, 622312, 1796636, 1337051, 328814, 28446, 1492, 820, 39496, 52786, 1549, 1796618, 582, 46507, 109327, 1531, 1382, 33039, 311460, 230143, 216935, 539, 35519, 1681, 328813, 214853, 89014, 1121115, 1585974, 29466, 1363, 292800, 270498, 214856, 142877, 133926, 209880, 179628, 1121102, 105612, 1796615, 39777, 29353, 1579, 163665, 53443, 261299, 1302, 1150298, 938289, 358742, 471875, 938278, 1796613, 1118057, 1077144, 1737, 218205, 1121298, 684066, 433659, 52699, 204516, 706562, 253257, 328812, 1280, 147802, 58134, 1335613, 891, 585394, 1582, 235931, 308994, 1589, 1682, 1736, 28129, 178001, 551788, 2051, 856, 118562, 101070, 515619, 40215, 187979, 82979, 29363, 1776391, 1285191, 84112, 157688, 38304, 36850, 341694, 287, 75612, 818, 371674, 338188, 88164, 588581, 676965, 546271, 1236512, 178338, 862517, 157687, 158, 51048, 1583331, 529, 888745, 394340, 40545, 855, 553973, 938293, 93063, 708634, 179995, 1351, 476652, 1464038, 555088, 237576, 879566, 1852371, 742727, 1377, 35830, 997353, 218538, 83771, 1605, 28111, 131109, 46609, 690567, 46206, 155615, 51616, 40542, 203, 294, 1034346, 156456, 80866, 554406, 796942, 1002367, 29347, 796944, 61592, 487175,

1050201, 762948, 137732, 1211819, 1019, 272548, 1717, 384636, 216940, 2087, 45634, 466107, 1689, 47678, 575, 979627, 840, 1660, 1236517, 617123, 546, 28135, 82171, 483, 501496, 99656, 1379, 84032, 39483, 1107316, 584, 28124, 1033744, 657309, 536441, 76123, 1118060, 89152, 76122, 303, 1541, 507751, 515620, 38302, 53419, 726, 40324, 1796610, 988946, 1852370, 1017, 1168289, 76936, 94869, 1161098, 215580, 1125779, 327575, 549, 1450648 and 478.

18. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population of bacteria or the at least two isolated or purified populations of bacteria are a species, subspecies or bacterial strains comprising a 16S rRNA gene sequence at least 80% identical (eg, at least 85, 90, 95 or 98% identical) to any one of the sequences of SEQ ID NOs: 1-876 in WO2018064165.

19. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population of bacteria or the at least two isolated or purified populations of bacteria belong to the species, subspecies or bacterial strains selected from the group consisting of *Bacteroides coagulans*, *Clostridium aldenense*, *Clostridium aldrichii*, *Clostridium alkalicellulosi*, *Clostridium amygdalinum*, *Clostridium asparagiforme*, *Clostridium cellulosi*, *Clostridium citroniae*, *Clostridium clariflavum* DSM 19732, *Clostridium clostridioforme*, *Clostridium colinum*, *Clostridium fimetarium*, *Clostridium hiranonis*, *Clostridium hungatei*, *Clostridium hylemonae* DSM 15053, *Clostridium indolis*, *Clostridium lactatifermentans*, *Clostridium leptum*, *Clostridium methylpentosum*, *Clostridium oroficum*, *Clostridium papyrosolvans* DSM 2782, *Clostridium populeti*, *Clostridium propionicum*, *Clostridium saccharolyticum*, *Clostridium scindens*, *Clostridium sporosphaeroides*, *Clostridium stercorarium*, *Clostridium straminisolvens*, *Clostridium sufflavum*, *Clostridium termifidis*, *Clostridium thermosuccino* genes, *Clostridium viride*, *Clostridium xylanolyticum*, *Desulfotomaculum guttoidium*, *Eubacterium rectale* ATCC 33656, *Eubacterium dolichum*, *Eubacterium eligens* ATCC 27750, *Eubacterium hallii*, *Eubacterium infirmum*, *Eubacterium siraeum*, *Eubacterium tenue*, *Ruminococcus torques*, *Acetanaerobacterium elongatum*, *Acetatifactor muris*, *Acetivibrio cellulolyticus*, *Acetivibrio ethanolgignens*, *Acholeplasma brassicae* 0502, *Acholeplasma parvum*, *Acholeplasma vituli*, *Acinetobacter junii*, *Actinobacillus porcinus*, *Actinomyces bowdenii*, *Actinomyces dentalis*, *Actinomyces odontolyticus*, *Acutalibacter muris*, *Aerococcus viridans*, *Aeromicrobium fasfidiosum*, *Alistipes finegoldii*, *Alistipes obesi*, *Alistipes onderdonkii*, *Alistipes putredinis*, *Alistipes shahii*, *Alistipes shahii* WAL 8301, *Alistipes timonensis* JC136, *Alkalibacter saccharofermentans*, *Alkaliphilus metalliredigens* QYMF, *Allisonella histaminiformans*, *Allobaculum stercoricans* DSM 13633, *Alloprevotella cava*, *Alloprevotella tannerae*, *Anaerobacterium chartisolvens*, *Anaerobiospirillum thomasi*, *Anaerobium acetethylcum*, *Anaerococcus octavius* NCTC 9810, *Anaerococcus provenciensis*, *Anaerococcus vaginalis* ATCC 51170, *Anaerocolumna jejuensis*, *Anaerofilum agile*, *Anaerofustis stercorihominis*, *Anaeroglobus geminatus*, *Anaeromassilibacillus senegalensis*, *Anaeroplasmabactoclasficum*, *Anaerorhabdus furcosa*, *Anaerosporebacter mobilis*, *Anaerostipes butyraticus*, *Anaerostipes caccae*, *Anaerostipes hadrus*,

Anaerotruncus colihominis, *Anaerovorax odorimutans*, *Anoxybacillus rupiensis*, *Aquabacterium limnoficum*, *Arcobacter butzleri*, *Arthrospira platensis*, *Asaccharobacter celatus*, *Atopobium parvulum*, *Bacteroides caccae*, *Bacteroides caecimuris*, *Bacteroides cellulosityticus*, *Bacteroides clarus* YIT 12056, *Bacteroides dorei*, *Bacteroides eggerthii*, *Bacteroides finegoldii*, *Bacteroides fragilis*, *Bacteroides gallinarum*, *Bacteroides massiliensis*, *Bacteroides oleiciplenus* YIT 12058, *Bacteroides plebeius* DSM 17135, *Bacteroides rodentium* JCM 16496, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*, *Bacteroides xylanisolvens* XB1A, *Bacteroides xylanolyticus*, *Barnesiella intestinihominis*, *Beduini massiliensis*, *Bifidobacterium bifidum*, *Bifidobacterium denfium*, *Bifidobacterium longum* subsp. *infantis*, *Blaufia caecimuris*, *Blaufia coccoides*, *Blaufia faecis*, *Blaufia glucerasesa*, *Blaufia hanseni* DSM 20583, *Blaufia hydrogenotrophica*, *Blaufia luti*, *Blaufia luti* DSM 14534, *Blaufia wexlerae* DSM 19850, *Budvicia aquatica*, *Butyricoccus pullicaecorum*, *Butyricimonas paravirosa*, *Butyrvibrio crossotus*, *Caldicoprobacter oshimai*, *Caloramator coolhaasii*, *Caloramator proteoclasficus*, *Caloramator quimbayensis*, *Campylobacter gracilis*, *Campylobacter rectus*, *Campylobacter ureolyticus* DSM 20703, *Capnocytophaga gingivalis*, *Capnocytophaga leadbetteri*, *Capnocytophaga sputigena*, *Casaltella massiliensis*, *Catabacter hongkongensis*, *Catenibacterium mitsuokai*, *Christensenella minuta*, *Christensenella timonensis*, *Chryseobacterium taklimakanense*, *Citrobacter freundii*, *Cloacibacillus porcorum*, *Clostridioides difficile* ATCC 9689=DSM 1296, *Clostridium amylolyticum*, *Clostridium bowmanii*, *Clostridium butyricum*, *Clostridium cadaveris*, *Clostridium colicanis*, *Clostridium gasigenes*, *Clostridium lentocellum* DSM 5427, *Clostridium oceanicum*, *Clostridium oryzae*, *Clostridium paraputrificum*, *Clostridium pascui*, *Clostridium perfringens*, *Clostridium quini*, *Clostridium saccharobutylicum*, *Clostridium sporogenes*, *Clostridium ventriculi*, *Collinsella aerofaciens*, *Comamonas testosteroni*, *Coprobacter fasfidiosus* NSB1, *Coprococcus eutactus*, *Corynebacterium diphtheriae*, *Corynebacterium durum*, *Corynebacterium mycetoides*, *Corynebacterium pyruviciproducens* ATCC BAA-1742, *Corynebacterium tuberculostearicum*, *Culturomica massiliensis*, *Cuneatibacter caecimuris*, *Desfluvitalea saccharophila*, *Delftia acidovorans*, *Desulfitobacterium chlororespirans*, *Desulfitobacterium metallireducens*, *Desulfosporosinus acididurans*, *Desulfotomaculum halophilum*, *Desulfotomaculum intricatum*, *Desulfotomaculum togense*, *Desulfovibrio desulfuricans* subsp. *desulfuricans*, *Desulfovibrio idahonensis*, *Desulfovibrio litoralis*, *Desulfovibrio piger*, *Desulfovibrio simplex*, *Desulfovibrio zosteriae*, *Desulfuromonas acetoxidans*, *Dethiobacter alkaliphilus* AHT 1, *Dethiosulfafibacter aminovorans*, *Dialister invisus*, *Dialister propionificaciens*, *Dielma fasfidiosa*, *Dietzia alimentaria* 72, *Dorea longicatena*, *Dysgonomonas gadei* ATCC BAA-286, *Dysgonomonas mossii*, *Eggerthella lenta*, *Eikenella corrodens*, *Eisenbergiella tayi*, *Emergencia timonensis*, *Enorma massiliensis* phi, *Enterococcus faecalis*, *Enterorhabdus muris*, *Ethanoligenes harbinense* YUAN-3, *Eubacterium coprostanoligenes*, *Eubacterium limosum*, *Eubacterium oxidoreducens*, *Eubacterium sulci* ATCC 35585, *Eubacterium uniforme*, *Eubacterium ventriosum*,

Eubacterium xylanophilum, *Extibacter muris*, *Ezak-
iella peruensis*, *Faecalibacterium prausnitzii*, *Faecali-
coccus acidiformans*, *Faecalitalea cylindroides*, *Fili-
factor villosus*, *Flavonifractor plautii*, *Flinfibacter
butyricus*, *Frisingococcus caecimuris*, *Fucophilus* 5
fucoidanalyticus, *Fusicatenibacter saccharivorans*,
Fusobacterium mortiferum, *Fusobacterium nucleatum*
subsp. *vincentii*, *Fusobacterium simiae*, *Fusobacterium
varium*, *Garciella nitratreducens*, *Gemella haemoly-
sans*, *Gemmiger formicilis*, *Gordonibacter urolithinfa-
ciens*, *Gracilibacter thermotolerans* JW/YJL-S1, 10
Granulicatella elegans, *Guggenheimella bovis*, *Haemophilus
haemolyticus*, *Helicobacter typhlonius*, *Hes-
pelliella stercorisuis*, *Holdemanella bififormis*, *Holdema-
nia massiliensis* AP2, *Howardella ureilytica*, 15
Hungatella Hungatella hathewayi, *Hydrogenoanaero-
bacterium saccharovorans*, *Ihubacter massiliensis*,
Intestinibacter bartlettii, *Intestinimonas butyriciprod-
ucens*, *Irregularibacter muris*, *Kiloniella laminariae*
DSM 19542, *Kroppenstedtia guangzhouensis*, *Lachno-
anaerobaculum orale*, *Lachnoanaerobaculum
umaense*, *Lachnoclostridium phytofermentans*, *Lacto-
bacillus acidophilus*, *Lactobacillus algidus*, *Lactoba-
cillus animalis*, *Lactobacillus casei*, *Lactobacillus del-
brueckii*, *Lactobacillus fomicalis*, *Lactobacillus iners*, 25
Lactobacillus pentosus, *Lactobacillus rogosae*, *Lacto-
coccus garvieae*, *Lactonifactor longoviformis*, *Leptotrichia
buccalis*, *Leptotrichia hofstadii*, *Leptotrichia
hongkongensis*, *Leptotrichia wadei*, *Leuconostoc
inhae*, *Levylla massiliensis*, *Loriellopsis cavemicola*, 30
Lutispora thermophila, *Marinilabilia salmonicolor*
JCM 21150, *Marvinbryanfia formatexigens*, *Meso-
plasma photuris*, *Methanobrevibacter smithii* ATCC
35061, *Methanomassiliicoccus luminyensis* BIO,
Methylobacterium extorquens, *Mitsuokellajalaludinii*, 35
Mobilitalea sibirica, *Mobiluncus curtisii*, *Mogibacte-
rium pumilum*, *Mogibacterium timidum*, *Moorella
glycerini*, *Moorella humiferrea*, *Moraxella nonliquefa-
ciens*, *Moraxella osloensis*, *Morganella morganii*,
Moryella indoligenes, *Muribaculum intestinale*, *Muri-
monas intestini*, *Natranaerovirga pectinivora*, *Neglecta
timonensis*, *Neisseria cinerea*, *Neisseria oxalis*, *Nocar-
dioides mesophilus*, *Novibacillus thermophilus*, *Ochro-
bactrum anthropi*, *Odoribacter splanchnicus*, 45
Olsenella profusa, *Olsenella uli*, *Oribacterium asac-
charolyticum* ACB7, *Oribacterium sinus*, *Oscillibacter
ruminantium* GH1, *Oscillibacter valericigenes*, *Oxo-
bacter pfennigii*, *Pantoea agglomerans*, *Papillibacter
cinnamivorans*, *Parabacteroides faecis*, *Parabacte-
roides goldsteinii*, *Parabacteroides gordonii*, 50
Parabacteroides merdae, *Parasporobacterium paucivorans*,
Parasutterella excrementihominis, *Parasutterella
secunda*, *Parvimonas micra*, *Peptococcus niger*, *Pepto-
niphilus duerdenii* ATCC BAA-1640, *Peptoniphilus
grossensis* ph5, *Peptoniphilus koenoeninae*, *Pepto-
niphilus senegalensis* JC140, *Peptostreptococcus
stomatis*, *Phascolarctobacterium succinatutens*, *Pho-
cea massiliensis*, *Porphibacter indicus*, *Porphyromonas
bennonis*, *Porphyromonas endodontalis*, *Porphyromonas
pasteri*, *Prevotella bergensis*, *Prevotella buccae* 60
ATCC 33574, *Prevotella denficola*, *Prevotella enoeca*,
Prevotella fusca JCM 17724, *Prevotella loescheii*,
Prevotella nigrescens, *Prevotella oris*, *Prevotella pollens*
ATCC 700821, *Prevotella stercorea* DSM 18206, *P rev
ote llamas silia timonensis*, *Propionispira arcuata*, 65
Proteus mirabilis, *Providencia rettgeri*, *Pseudobacte-
roides celluloso-solvens* ATCC 35603=DSM 2933,

Pseudobutyrvibrio ruminis, *Pseudoflavonifractor
capillosus* ATCC 29799, *Pseudomonas aeruginosa*,
Pseudomonas fluorescens, *Pseudomonas mandelii*,
Pseudomonas nitroreducens, *Pseudomonas pufida*,
Raoultella ornithinolytica, *Raoultella planficola*,
Raoultibacter massiliensis, *Robinsoniella peoriensis*,
Romboutsia timonensis, *Roseburia faecis*, *Roseburia
hominis* A2-183, *Roseburia intestinalis*, *Roseburia inu-
linivorans* DSM 16841, *Rothia dentocariosa* ATCC
17931, *Ruminiclostridium thermocellum*, *Ruminococ-
cus albus*, *Ruminococcus bromii*, *Ruminococcus calli-
dus*, *Ruminococcus champanellensis* 18P13=JCM
17042, *Ruminococcus faecis* JCM 15917, *Ruminococ-
cus flavefaciens*, *Ruminococcus gnavreuii*, *Rumi-
nococcus lactaris* ATCC 29176, *Rummeliibacillus pyc-
nus*, *Saccharofermentans acetigenes*, *Scardovia
wiggisiae*, *Schlegelella thermodepolymerans*, *Sedimen-
fibacter hongkongensis*, *Selenomonas sputigena* ATCC
35185, *Slackia exigua* ATCC 700122, *Slackia piri-
formis* YIT 12062, *Solitalea canadensis*, *Solobacterium
moorei*, *Sphingomonas aquafilis*, *Spiroplasma allegh-
enense*, *Spiroplasma chinense*, *Spiroplasma chryso-
picola*, *Spiroplasma culicicola*, *Spiroplasma lampro-
pyridicola*, *Sporobacter termifidis*, *Staphylococcus aureus*,
Stenotrophomonas maltophilia, *Stomatobaculum
longum*, *Streptococcus agalactiae* ATCC 13813, *Stre-
ptococcus cristatus*, *Streptococcus equinus*, *Streptococ-
cus gordonii*, *Streptococcus lactarius*, *Streptococcus
parauberis*, *Subdoligranulum variabile*, *Succinivibrio
dextrinosolvens*, *Sutterella stercoricanis*, *Sutterella
wadsworthensis*, *Syntrophococcus sucromutans*, *Syn-
trophomonas zehnderi* OL-4, *Terrisporobacter mayo-
mbei*, *Thermoleophilum album*, *Treponema denficola*,
Treponema socranskii, *Tyzzeraella nexilis* DSM 1787,
Vallitalea guaymasensis, *Vallitalea pronyensis*, *Vam-
pirovibrio chlorellavorus*, *Veillonella atypica*, *Veillo-
nella denficariosi*, *Veillonella dispar*, *Veillonella par-
vula*, *Victivallis vadensis*, *Vulcambacillus
modesticaldus* and *Weissella confusa*.

In an example, the transplant comprises or consists of
SER-109 or SER-262 (and optionally the condition is a *C
difficile* infection); VE202 or SER-287 (and optionally the
disease is ulcerative colitis); SER-301 (and optionally the
disease is IBD); SER-401 (and optionally the condition is a
cancer; eg, wherein the therapy further comprises adminis-
tration of an anti-PD-1 axis antibody, eg, an anti-PD-1
antibody); VE800 or SER-155 (and optionally the therapy
further comprises the administration of a transplant, eg, a
haematopoietic stem cell or solid organ transplant);
EDP1066 or EDP1815 (and optionally the disease is an
inflammatory condition, eg, colitis, Crohn's disease, asthma,
rheumatoid arthritis (RA), psoriasis, dermatitis (eg, atopic
dermatitis) or IBD); or EDP1503 (and the disease is a
cancer, eg, colorectal cancer, renal cell carcinoma, mela-
noma or a PD-1 relapsed cancer). In an example, the therapy
comprises the administration of SGM-1019, SG-2-0776 or
EB8018 (and optionally the disease or condition is NASH or
IBD or an inflammatory condition, eg, colitis, Crohn's
disease, asthma, rheumatoid arthritis (RA), psoriasis and
dermatitis (eg, atopic dermatitis)). Those starting "VE" are
developed by Vadanta Biosciences, SER are developed by
Seres Therapeutics, EDP are developed by Evelo Biosci-
ences, SG are developed by Second Genome and EB are
developed by Enterome.

In an example, the disease or condition herein is an inflammatory condition, eg, colitis, Crohn's disease, asthma, rheumatoid arthritis (RA), psoriasis, dermatitis (eg, atopic dermatitis) or IBD.

6. Other Agents

It is contemplated that other agents may be used in combination with certain aspects of the present embodiments to improve the therapeutic efficacy of treatment. These additional agents include agents that affect the upregulation of cell surface receptors and GAP junctions, cytostatic and differentiation agents, inhibitors of cell adhesion, agents that increase the sensitivity of the hyperproliferative cells to apoptotic inducers, or other biological agents. Increases in intercellular signalling by elevating the number of GAP junctions would increase the anti-hyperproliferative effects on the neighbouring hyperproliferative cell population. In other embodiments, cytostatic or differentiation agents can be used in combination with certain aspects of the present embodiments to improve the anti-hyperproliferative efficacy of the treatments. Inhibitors of cell adhesion are contemplated to improve the efficacy of the present embodiments. Examples of cell adhesion inhibitors are focal adhesion kinase (FAKs) inhibitors and Lovastatin. It is further contemplated that other agents that increase the sensitivity of a hyperproliferative cell to apoptosis, such as the antibody c225, could be used in combination with certain aspects of the present embodiments to improve the treatment efficacy.

Diseases and Conditions

- Optionally, the disease or condition is selected from
- (a) A neurodegenerative disease or condition;
 - (b) A brain disease or condition;
 - (c) A CNS disease or condition;
 - (d) Memory loss or impairment;
 - (e) A heart or cardiovascular disease or condition, eg, heart attack, stroke or atrial fibrillation;
 - (f) A liver disease or condition;
 - (g) A kidney disease or condition, eg, chronic kidney disease (CKD);
 - (h) A pancreas disease or condition;
 - (i) A lung disease or condition, eg, cystic fibrosis or COPD;
 - (j) A gastrointestinal disease or condition;
 - (k) A throat or oral cavity disease or condition;
 - (l) An ocular disease or condition;
 - (m) A genital disease or condition, eg, a vaginal, labial, penile or scrotal disease or condition;
 - (n) A sexually-transmissible disease or condition, eg, gonorrhoea, HIV infection, syphilis or *Chlamydia* infection;
 - (o) An ear disease or condition;
 - (p) A skin disease or condition;
 - (q) A heart disease or condition;
 - (r) A nasal disease or condition
 - (s) A haematological disease or condition, eg, anaemia, eg, anaemia of chronic disease or cancer;
 - (t) A viral infection;
 - (u) A pathogenic bacterial infection;
 - (v) A cancer;
 - (w) An autoimmune disease or condition, eg, SLE;
 - (x) An inflammatory disease or condition, eg, rheumatoid arthritis, psoriasis, eczema, asthma, ulcerative colitis, colitis, Crohn's disease or IBD;
 - (y) Autism;
 - (z) ADHD;
 - (aa) Bipolar disorder;
 - (bb) ALS [Amyotrophic Lateral Sclerosis];
 - (cc) Osteoarthritis;
 - (dd) A congenital or development defect or condition;
 - (ee) Miscarriage;

- (ff) A blood clotting condition;
 - (gg) Bronchitis;
 - (hh) Dry or wet AMD;
 - (ii) Neovascularisation (eg, of a tumour or in the eye);
 - 5 (jj) Common cold;
 - (kk) Epilepsy;
 - (ll) Fibrosis, eg, liver or lung fibrosis;
 - (mm) A fungal disease or condition, eg, thrush;
 - (nn) A metabolic disease or condition, eg, obesity, anorexia,
 - 10 (oo) Ulcer(s), eg, gastric ulceration or skin ulceration;
 - (pp) Dry skin;
 - (qq) Sjogren's syndrome;
 - (rr) Cytokine storm;
 - 15 (ss) Deafness, hearing loss or impairment;
 - (tt) Slow or fast metabolism (ie, slower or faster than average for the weight, sex and age of the subject);
 - (uu) Conception disorder, eg, infertility or low fertility;
 - (vv) Jaundice;
 - 20 (ww) Skin rash;
 - (xx) Kawasaki Disease;
 - (yy) Lyme Disease;
 - (zz) An allergy, eg, a nut, grass, pollen, dust mite, cat or dog fur or dander allergy;
 - 25 (aaa) Malaria, typhoid fever, tuberculosis or cholera;
 - (bbb) Depression;
 - (ccc) Mental retardation;
 - (ddd) Microcephaly;
 - (eee) Malnutrition;
 - 30 (fff) Conjunctivitis;
 - (ggg) Pneumonia;
 - (hhh) Pulmonary embolism;
 - (iii) Pulmonary hypertension;
 - (jjj) A bone disorder;
 - 35 (kkk) Sepsis or septic shock;
 - (lll) Sinusitis;
 - (mmm) Stress (eg, occupational stress);
 - (nnn) Thalassaemia, anaemia, von Willebrand Disease, or haemophilia;
 - (ooo) Shingles or cold sore;
 - (ppp) Menstruation;
 - (qqq) Low sperm count.
- Neurodegenerative or CNS Diseases or Conditions for Treatment or Prevention by the Method
- 45 In an example, the neurodegenerative or CNS disease or condition is selected from the group consisting of Alzheimer disease, geriatric psychosis, Down syndrome, Parkinson's disease, Creutzfeldt Jakob disease, diabetic neuropathy, Parkinson syndrome, Huntington's disease, Machado-Joseph disease, amyotrophic lateral sclerosis, diabetic neuropathy,
- 50 and Creutzfeldt Creutzfeldt-Jakob disease. For example, the disease is Alzheimer disease. For example, the disease is Parkinson syndrome.
- In an example, wherein the method of the invention is practiced on a human or animal subject for treating a CNS or neurodegenerative disease or condition, the method causes downregulation of Treg cells in the subject, thereby promoting entry of systemic monocyte-derived macrophages and/or Treg cells across the choroid plexus into the brain of the subject, whereby the disease or condition (eg, Alzheimer's disease) is treated, prevented or progression thereof is reduced. In an embodiment the method causes an increase of IFN-gamma in the CNS system (eg, in the brain and/or CSF) of the subject. In an example, the method
- 65 restores nerve fibre and/or reduces the progression of nerve fibre damage. In an example, the method restores nerve myelin and/or reduces the progression of nerve myelin

damage. In an example, the method of the invention treats or prevents a disease or condition disclosed in WO2015136541 and/or the method can be used with any method disclosed in WO2015136541 (the disclosure of this document is incorporated by reference herein in its entirety, eg, for providing disclosure of such methods, diseases, conditions and potential therapeutic agents that can be administered to the subject for effecting treatment and/or prevention of CNS and neurodegenerative diseases and conditions, eg, agents such as immune checkpoint inhibitors, eg, anti-PD-1, anti-PD-L1, anti-TIM3 or other antibodies disclosed therein).

Cancers for Treatment or Prevention by the Method

Cancers that may be treated include tumours that are not vascularized, or not substantially vascularized, as well as vascularized tumours. The cancers may comprise non-solid tumours (such as haematological tumours, for example, leukaemias and lymphomas) or may comprise solid tumours. Types of cancers to be treated with the invention include, but are not limited to, carcinoma, blastoma, and sarcoma, and certain leukaemia or lymphoid malignancies, benign and malignant tumours, and malignancies e.g., sarcomas, carcinomas, and melanomas. Adult tumours/cancers and paediatric tumours/cancers are also included.

Haematologic cancers are cancers of the blood or bone marrow. Examples of haematological (or haematogenous) cancers include leukaemias, including acute leukaemias (such as acute lymphocytic leukaemia, acute myelocytic leukaemia, acute myelogenous leukaemia and myeloblasts, promyeiocyctic, myelomonocytic, monocytic and erythroleukaemia), chronic leukaemias (such as chronic myelocytic (granulocytic) leukaemia, chronic myelogenous leukaemia, and chronic lymphocytic leukaemia), polycythemia vera, lymphoma, Hodgkin's disease, non-Hodgkin's lymphoma (indolent and high grade forms), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, myelodysplastic syndrome, hairy cell leukaemia and myelodysplasia.

Solid tumours are abnormal masses of tissue that usually do not contain cysts or liquid areas. Solid tumours can be benign or malignant. Different types of solid tumours are named for the type of cells that form them (such as sarcomas, carcinomas, and lymphomas). Examples of solid tumours, such as sarcomas and carcinomas, include fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteosarcoma, and other sarcomas, synovioma, mesothelioma, Ewing's tumour, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, lymphoid malignancy, pancreatic cancer, breast cancer, lung cancers, ovarian cancer, prostate cancer, hepatocellular carcinoma, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, medullary thyroid carcinoma, papillary thyroid carcinoma, pheochromocytomas sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, Wilms' tumour, cervical cancer, testicular tumour, seminoma, bladder carcinoma, melanoma, and CNS tumours (such as a glioma (such as brainstem glioma and mixed gliomas), glioblastoma (also known as glioblastoma multiforme) astrocytoma, CNS lymphoma, germinoma, medulloblastoma, Schwannoma craniopharyngioma, ependymoma, pineaioma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, neuroblastoma, retinoblastoma and brain metastases).

In an example, the cancer is a haematological cancer. In an example, the cancer is NSCLC. In an example, the cancer

is renal cell carcinoma. In an example, the cancer is urothelial carcinoma. In an example, the cancer is melanoma. Autoimmune Diseases for Treatment or Prevention by the Method

- 5 Acute Disseminated Encephalomyelitis (ADEM)
- Acute necrotizing hemorrhagic leukoencephalitis
- Addison's disease
- Agammaglobulinemia
- 10 Alopecia areata
- Amyloidosis
- Ankylosing spondylitis
- Anti-GBM/Anti-TBM nephritis
- Antiphospholipid syndrome (APS)
- 15 Autoimmune angioedema
- Autoimmune aplastic anemia
- Autoimmune dysautonomia
- Autoimmune hepatitis
- Autoimmune hyperlipidemia
- 20 Autoimmune immunodeficiency
- Autoimmune inner ear disease (AIED)
- Autoimmune myocarditis
- Autoimmune oophoritis
- Autoimmune pancreatitis
- 25 Autoimmune retinopathy
- Autoimmune thrombocytopenic purpura (ATP)
- Autoimmune thyroid disease
- Autoimmune urticaria
- Axonal & neuronal neuropathies
- 30 Balo disease
- Behcet's disease
- Bullous pemphigoid
- Cardiomyopathy
- Castleman disease
- 35 Celiac disease
- Chagas disease
- Chronic fatigue syndrome
- Chronic inflammatory demyelinating polyneuropathy (CIDP)
- 40 Chronic recurrent multifocal osteomyelitis (CRMO)
- Churg-Strauss syndrome
- Cicatricial pemphigoid/benign mucosal pemphigoid
- Crohn's disease
- Cogans syndrome
- 45 Cold agglutinin disease
- Congenital heart block
- Coxsackie myocarditis
- CREST disease
- Essential mixed cryoglobulinemia
- 50 Demyelinating neuropathies
- Dermatitis herpetiformis
- Dermatomyositis
- Devic's disease (neuromyelitis optica)
- Discoid lupus
- 55 Dressler's syndrome
- Endometriosis
- Eosinophilic esophagitis
- Eosinophilic fasciitis
- Erythema nodosum
- 60 Experimental allergic encephalomyelitis
- Evans syndrome
- Fibromyalgia
- Fibrosing alveolitis
- Giant cell arteritis (temporal arteritis)
- 65 Giant cell myocarditis
- Glomerulonephritis
- Goodpasture's syndrome

Granulomatosis with Polyangiitis (GPA) (formerly called Wegener's Granulomatosis)
 Graves' disease
 Guillain-Barre syndrome
 Hashimoto's encephalitis
 Hashimoto's thyroiditis
 Hemolytic anemia
 Henoch-Schonlein purpura
 Herpes gestationis
 Hypogammaglobulinemia
 Idiopathic thrombocytopenic purpura (ITP)
 IgA nephropathy
 IgG4-related sclerosing disease
 Immunoregulatory lipoproteins
 Inclusion body myositis
 Interstitial cystitis
 Juvenile arthritis
 Juvenile diabetes (Type 1 diabetes)
 Juvenile myositis
 Kawasaki syndrome
 Lambert-Eaton syndrome
 Leukocytoclastic vasculitis
 Lichen planus
 Lichen sclerosus
 Ligneous conjunctivitis
 Linear IgA disease (LAD)
 Lupus (SLE)
 Lyme disease, chronic
 Meniere's disease
 Microscopic polyangiitis
 Mixed connective tissue disease (MCTD)
 Mooren's ulcer
 Mucha-Habermann disease
 Multiple sclerosis
 Myasthenia gravis
 Myositis
 Narcolepsy
 Neuromyelitis optica (Devic's)
 Neutropenia
 Ocular cicatricial pemphigoid
 Optic neuritis
 Palindromic rheumatism
 PANDAS (Pediatric Autoimmune Neuropsychiatric Disorders Associated with *Streptococcus*)
 Paraneoplastic cerebellar degeneration
 Paroxysmal nocturnal hemoglobinuria (PNH)
 Parry Romberg syndrome
 Parsonnage-Turner syndrome
 Pars planitis (peripheral uveitis)
 Pemphigus
 Peripheral neuropathy
 Perivenous encephalomyelitis
 Pernicious anemia
 POEMS syndrome
 Polyarteritis nodosa
 Type I, II, & III autoimmune poly glandular syndromes
 Polymyalgia rheumatica
 Polymyositis
 Postmyocardial infarction syndrome
 Postpericardiotomy syndrome
 Progesterone dermatitis
 Primary biliary cirrhosis
 Primary sclerosing cholangitis
 Psoriasis
 Psoriatic arthritis
 Idiopathic pulmonary fibrosis
 Pyoderma gangrenosum

Pure red cell aplasia
 Raynauds phenomenon
 Reactive Arthritis
 Reflex sympathetic dystrophy
 Reiter's syndrome
 Relapsing polychondritis
 Restless legs syndrome
 Retroperitoneal fibrosis
 Rheumatic fever
 Rheumatoid arthritis
 Sarcoidosis
 Schmidt syndrome
 Scleritis
 Scleroderma
 Sjogren's syndrome
 Sperm & testicular autoimmunity
 Stiff person syndrome
 Subacute bacterial endocarditis (SBE)
 Susac's syndrome
 Sympathetic ophthalmia
 Takayasu's arteritis
 Temporal arteritis/Giant cell arteritis
 Thrombocytopenic purpura (TTP)
 Tolosa-Hunt syndrome
 Transverse myelitis
 Type 1 diabetes
 Ulcerative colitis
 Undifferentiated connective tissue disease (UCTD)
 Uveitis
 Vasculitis
 Vesiculobullous dermatosis
 Vitiligo
 Wegener's granulomatosis (now termed Granulomatosis with Polyangiitis (GPA)).
 Inflammatory Diseases for Treatment or Prevention by the Method
 Alzheimer's
 ankylosing spondylitis
 arthritis (osteoarthritis, rheumatoid arthritis (RA), psoriatic arthritis)
 asthma
 atherosclerosis
 Crohn's disease
 colitis
 dermatitis
 diverticulitis
 fibromyalgia
 hepatitis
 irritable bowel syndrome (IBS)
 systemic lupus erythematosus (SLE)
 nephritis
 Parkinson's disease
 ulcerative colitis.
 Concepts:—
 The invention provides the following Concepts.
 1. A programmable nuclease for use in a method of treating an acute microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes

- comprised by the subject are cut and acute microbial infection of the subject is treated.
2. A programmable nuclease for use in a method of durably treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are durably killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is durably treated.
 3. The nuclease of Concept 2, wherein the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1.
 4. The nuclease of any preceding Concept, wherein the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
 5. The nuclease of any preceding Concept, wherein the method comprises maintaining reduction of the infection by at least 100-fold for at least 60 minutes (eg, at least 120 minutes) after exposing the subject to the programmed nuclease.
 6. The nuclease of any preceding Concept, wherein the method comprises reducing the infection such that the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.
 7. The nuclease of any preceding Concept, wherein the method comprises administering to the subject a RNA or a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.
 8. The nuclease of Concept 7, wherein the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject.
 9. The nuclease of Concept 7, wherein the subject comprises the nuclease prior to administration of the RNA or nucleic acid to the subject.
 10. The nuclease of any one of Concepts 7 to 9, wherein a plurality of viruses (eg, phage) are administered to the subject, wherein each virus comprises a copy of the nucleic acid, wherein the viruses infect the microbes comprised by the subject to deliver thereto the nucleic acid.
 11. The nuclease of Concept 10, wherein the ratio of administered viruses:microbes comprised by the subject is from 10 to 150.
 12. The nuclease according to any preceding Concept, wherein the subject is a human or animal, optionally wherein the subject is a human over 65 years of age or is a paediatric patient.
 13. The nuclease according to Concept 12, wherein the infection is an infection of the lungs, abdomen or urinary tract; or wherein the subject has undergone surgery, is on an immunosuppressant medication and/or is suffering from a chronic disease.
 14. The nuclease according to any preceding Concept, wherein the infection is reduced by at least 90% for 1

- hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
15. The nuclease according to any preceding Concept, wherein the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.
 16. The nuclease according to any one of Concepts 12 to 15, wherein the method treats or prevents septicaemia and/or sepsis (eg, septic shock) in the subject.
 17. The nuclease of Concept 16, wherein at the start of the treatment, the subject (eg, a human) has a temperature of $<36^{\circ}\text{C}$. or $>38^{\circ}\text{C}$.; a heart rate of $>90/\text{min}$, a respiratory rate of >20 breaths/min or $\text{PaCO}_2 < 4.3\text{ kPa}$; and white blood cell count of $<4000/\text{mm}^3$ or $>12,000/\text{mm}^3$.
 18. The nuclease of Concept 16 or 17, wherein at the start of the treatment, the subject (eg, a human) has presence of two or more of the following: abnormal body temperature, abnormal heart rate, abnormal respiratory rate, abnormal blood gas and abnormal white blood cell count.
 19. The nuclease of any preceding Concept, wherein the subject is a human or animal and the microbes are bacteria (eg, *E coli* or *C difficile*), wherein blood infection of the subject by the bacteria is reduced at least 100- or 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
 20. The nuclease of any one of Concepts 12 to 19, wherein the blood of the subject is infected with from 10^7 to 10^{12} CFU/ml of the bacteria immediately before the treatment.
 21. The nuclease according to any one of Concepts 1 to 11, wherein the subject is a plant.
 22. The nuclease according to any preceding Concept, wherein the microbes are bacteria.
 23. The nuclease according to Concept 22, wherein the bacteria are gram positive bacteria.
 24. The nuclease according to Concept 22 or 23, wherein the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).
 25. The nuclease according to any preceding Concept, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.
 26. A plurality of viruses (eg, phage or phagemids for producing phage) for use with the nuclease of any preceding Concept in the method of treatment, wherein each virus comprises a copy of a nucleic acid as defined in any one of Concepts 7 to 9, wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.
 27. A composition comprising a plurality of nucleic acids for programming the nuclease of any one of Concepts 1 to 25 in the method of treatment, wherein each nucleic acid is a nucleic acid as defined in any one of Concepts 7 to 9.
 28. A CRISPR/Cas system comprising a nuclease according to any preceding Concept for use in the method of

- treatment, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of the microbes.
29. A guide RNA or a DNA encoding a guide RNA for use in the system of Concept 28 for use in the method of treating an acute microbial infection in the subject, eg, septicaemia or sepsis.
 30. A nucleic acid vector comprising the guide RNA or DNA recited in Concept 27 or 29.
 31. The vector of Concept 30 wherein the vector is a phage, phagemid, viriophage, virus, plasmid (eg, conjugative plasmid) or transposon.
 32. An anti-sepsis or anti-septicaemia composition for administration to a human or animal for treating sepsis or septicaemia, the composition comprising a plurality of vectors, wherein each vector is according to Concept 30 or 31.
 33. A method of treating an acute microbial infection of a subject, wherein the method is as defined by any preceding Concept.
 34. Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of any one of Concepts 1 to 26 and 28 to 30, in the manufacture of a composition for carrying out a method of treatment as defined by any preceding Concept, wherein the subject is an organism other than a human or animal.
 35. Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of any one of Concepts 1 to 26 and 28 to 30, in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.
 36. Use of a programmable nuclease in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.
 37. The use of Concept 34, 35 or 36, wherein the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject or substrate at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1.

38. The use of any one of Concepts 34 to 37, wherein the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
39. The use of any one of Concepts 34 to 38, wherein the reduction of the infection is maintained by at least 100-fold for at least 60 minutes (eg, at least 120 minutes) after exposing the subject to the programmed nuclease.
40. The use of any one of Concepts 34 to 39, wherein the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.
41. The use of any one of Concepts 34 to 40, wherein the method comprises administering to the subject or substrate a RNA or a nucleic acid that encodes an RNA for expression of the RNA in or on the subject or substrate, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject or substrate.
42. The use of Concept 41, wherein the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject or substrate.
43. The use of Concept 41, wherein the subject or substrate comprises the nuclease prior to administration of the RNA or nucleic acid.
44. The use of any one of Concepts 41 to 43, wherein a plurality of viruses (eg, phage) are administered to the subject or substrate, wherein each virus comprises a copy of the nucleic acid, wherein the viruses infect the microbes comprised by the subject or substrate to deliver thereto the nucleic acid.
45. The use of Concept 44, wherein the ratio of administered viruses:microbes is from 10 to 150.
46. The use of any one of Concepts 34 to 45, wherein the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
47. The use of any one of Concepts 34 to 46, wherein the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject or substrate to the programmed nuclease.
48. The use of any one of Concepts 34 to 47, wherein the subject is a plant; or wherein the substrate is a metallic, plastic, concrete, stone, wood, glass or ceramic substrate.
49. The use of any one of Concepts 34 to 48, wherein the microbes are bacteria.
50. The use according to Concept 49, wherein the bacteria are gram positive bacteria.
51. The use according to Concept 49 or 50, wherein the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Heamophilus*, *Ghonnorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).
52. The use of any one of Concepts 34 to 51, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.

Embodiments

1. A method for treating a pathogenic bacterial infection in a human or animal subject caused by bacteria (first

- bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a programmable nuclease that is programmed to cut the target site, wherein the subject is suffering from a further disease or condition other than the pathogenic bacterial infection and the method comprises administering a therapy to the subject for treating or preventing the further disease or condition, wherein the nuclease treats the infection and the therapy is efficacious in the presence of the programmed nuclease to treat or prevent the disease or condition.
2. The method of Embodiment 1, wherein the subject is a cancer patient and the therapy comprises administration of a haematopoietic stem cell transplant, chemotherapeutic agent, immune checkpoint inhibitor, immune checkpoint agonist or an immune cell enhancer; adoptive cell therapy; radiation or surgery.
 3. The method of Embodiment 2, wherein the therapy is an immune checkpoint inhibitor antibody, or an antibody selected from ipilimumab (or YERVOY®), tremelimumab, nivolumab (or OPDIVO®), pembrolizumab (or KEYTRUDA®), pidilizumab, BMS-936559, durvalumab (or IMFINZI®) and atezolizumab (or TECENTRIQ®).
 4. The method of Embodiment 1, wherein the therapy is a tissue, organ or cell transplant.
 5. The method of Embodiment 1, wherein the treatment of the bacterial infection is carried out simultaneously with the administration of the therapy to the subject.
 6. The method of Embodiment 1, wherein the treatment of the bacterial infection is carried out immediately before or after administering the therapy to the subject.
 7. The method of Embodiment 1, wherein the method comprises administering to the subject a or a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in first bacteria comprised by the subject, thereby killing the first bacteria.
 8. The method of Embodiment 1, comprising administering a nucleic acid vector to the subject, wherein the vector encodes the programmable nuclease.
 9. The method of Embodiment 1, wherein the programmable nuclease is an endogenous nuclease of the first cells.
 10. The method of Embodiment 1, wherein the efficacy of the therapy in the presence of the programmed nuclease is greater than the efficacy of the therapy in the presence of a broad-spectrum antibiotic.
 11. The method of Embodiment 1, wherein the efficacy of the therapy in the presence of the programmed nuclease is greater than the efficacy of the therapy in the presence of an antibiotic selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbapristin; teicoplanin; cephalosporin; carbapenem; fluoroquinolone; aminoglycoside; colistin; erythromycin; clindamycin; beta-lactam; macrolide; amoxicillin; azithromycin; penicillin; ceftriaxone; azithromycin; ciprofloxacin; isoniazid (INH); rifampicin (RMP); amikacin; kanamycin; capreomycin; trimethoprim; itrofurantoin; cefalexin; amoxicillin; metronidazole (MTZ); cefixime; tetracycline; and meropenem.
 12. The method of Embodiment 1, wherein the first bacteria is selected from (i) *Staphylococcus aureus* that is resistant to an antibiotic selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbapristin and teicoplanin; (ii) *Pseudomonas aeruginosa* that is resistant to an antibiotic selected from cephalosporins, carbapen-

- ems, fluoroquinolones, aminoglycosides and colistin; (iii) *Klebsiella* species that is resistant to carbapenem; (iv) *Streptococcus* species that is resistant to an antibiotic selected from erythromycin, clindamycin, beta-lactam, macrolide, amoxicillin, azithromycin and penicillin; (v) *Salmonella* species that is resistant to an antibiotic selected from ceftriaxone, azithromycin and ciprofloxacin; (vi) *Shigella* species that is resistant to ciprofloxacin or azithromycin; (vii) *Mycobacterium tuberculosis* that is resistant to an antibiotic selected from Resistance to isoniazid (INH), rifampicin (RMP), fluoroquinolone, amikacin, kanamycin, capreomycin and azithromycin; (viii) *Enterococcus* species that is resistant to vancomycin; (ix) *Enterobacteriaceae* species that is resistant to an antibiotic selected from cephalosporin and carbapenem; (x) *E coli* that is resistant to an antibiotic selected from trimethoprim, itrofurantoin, cefalexin and amoxicillin; (xi) *Clostridium* species that is resistant to metronidazole (MTZ), fluoroquinolone or carbapenem; (xii) *Neisseria gonorrhoea* that is resistant to an antibiotic selected from cefixime, ceftriaxone, azithromycin and tetracycline; (xiii) *Acinetobacter baumannii* that is resistant to an antibiotic selected from beta-lactam, meropenem and carbapenem; and (xiv) *Campylobacter* species that is resistant to ciprofloxacin or azithromycin.
13. The method of Embodiment 1, wherein the treatment of the infection treats or prevents in the subject a condition selected from vaginosis, meningitis, pneumonia, urinary tract infection, cystitis, nephritis, gastroenteritis, a skin infection, impetigo, erysipelas, cellulitis, septicaemia or sepsis in the subject.
 14. The method of Embodiment 1, wherein the further disease or condition is a cancer; autoimmune disease or condition; or GI tract disease or condition.
 15. The method of Embodiment 1, wherein the subject comprises bacteria (second bacteria) of one or more strains or species that are different to the first strain or species, wherein the genomes of the second bacteria do not comprise the target site, wherein the genomes of the second bacteria are not cut by the programmed nuclease in the subject, whereby second bacteria survive in the presence of the programmed nuclease in the patient; and wherein the therapy is efficacious in the presence of the second bacteria.
 16. The method of Embodiment 15, wherein reduction in the second bacteria in patients is associated with reduced efficacy of the therapy.
 17. The method of Embodiment 15, wherein the second bacteria are selected from the group consisting of *Akkermansia*, *Alistipes*, *Bacteroides*, *Barnesiella*, *Bifidobacterium*, *Clostridium*, *Collinsella*, *Enterococcus*, *Fusobacterium*, *Lactobacillus*, *Propionibacterium*, *Ruminococcus*, *Segmented filamentous* bacteria (SFB); *Veillonella*, *Prevotella*, *Escherichia* and *Streptococcus* bacteria.
 18. The method of Embodiment 1, wherein the first bacteria are selected from the group consisting of *E coli*, *C difficile*, *V cholera*, *Staphylococcus*, *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* bacteria.
 19. The method of Embodiment 1, wherein the nuclease is a Cas nuclease, a meganuclease, a Transcription activator-like effector nuclease (TALEN) or zinc finger nuclease.
 20. A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria,

wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer.

It will be understood that particular embodiments described herein are shown by way of illustration and not as limitations of the invention. The principal features of this invention can be employed in various embodiments without departing from the scope of the invention. Those skilled in the art will recognize, or be able to ascertain using no more than routine study, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of this invention and are covered by the claims. All publications and patent applications mentioned in the specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications and all US equivalent patent applications and patents are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference. The use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.” The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” Throughout this application, the term “about” is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

As used in this specification and claim(s), the words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, unrecited elements or method steps.

The term “or combinations thereof” or similar as used herein refers to all permutations and combinations of the listed items preceding the term. For example, “A, B, C, or combinations thereof” is intended to include at least one of: A, B, C, AB, AC, BC, or ABC, and if order is important in a particular context, also BA, CA, CB, CBA, BCA, ACB, BAC, or CAB. Continuing with this example, expressly included are combinations that contain repeats of one or more item or term, such as BB, AAA, MB, BBC, AAAB-CCCC, CBBAAA, CABABB, and so forth. The skilled artisan will understand that typically there is no limit on the number of items or terms in any combination, unless otherwise apparent from the context.

Any part of this disclosure may be read in combination with any other part of the disclosure, unless otherwise apparent from the context.

All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the

compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

The present invention is described in more detail in the following non limiting Examples.

EXAMPLES

Precision Fast Bacteria Killing with Programmable Nucleases

The examples provide a method for fast and precision killing of *Escherichia coli* and *Clostridium difficile* strains. As a model programmable nuclease system, we used a CRISPR guided vector (CGV™) system to specifically target enterohemorrhagic *E. coli* (EHEC) and probiotic *E. coli* Nissle.

Example 1. Precision Killing of Target Strain Enterohemorrhagic *E. coli* (EHEC)

1.1. Design, Construction and Delivery of CRISPR Guided Vector (CGV) System Targeting *E. coli* (EHEC) ATCC43888.

The invention provides a CGV system to specifically target enterohemorrhagic *E. coli* (EHEC) ATCC43888 (a human fecal isolate obtained from the American Type Culture Collection). The CGV system comprises two vectors: (a) a vector containing a tracrRNA and the Cas9 protein from *Streptococcus pyogenes* (SpCas); (b) a vector containing a guide RNA (gRNA) that comprises a nucleotide sequence capable of hybridizing to a target sequence in the host cells to guide SpCas9 to the target sequence. To enable specific killing of *E. coli* (EHEC) ATCC43888, a particular sequence from the genome of this strain was chosen to target. Specifically, the sequence contains 20 nucleotides from the 23S ribosomal RNA gene from *E. coli* (EHEC) ATCC43888. Additionally, the 5'-NGG protospacer adjacent motif (PAM) was located adjacent to the selected target sequence. The selected target sequence in the 23S rRNA gene can be found in Table 3.

1.2 Characterization of the CGV System Targeting *E. coli* (EHEC) ATCC43888.

To establish the CGV system functionality in mediating sequence-specific killing in *E. coli* (EHEC) ATCC43888, the system was transformed into *E. coli* (EHEC) ATCC43888 cells. Overnight cultures were diluted 1:100 in fresh lysogeny broth (LB) and grown to mid-exponential phase OD600~0.6. The CRISPR system was induced by adding theophylline and arabinose (2 mM theophylline and 1% arabinose), and survival of the strain was followed over time by plating the cultures in serial dilutions every 15 minutes, for 1 h (FIG. 1B). CRISPR induction in *E. coli* (EHEC) surprisingly triggered a rapid killing of the cells, achieving 99.98% killing within 30 minutes of induction (FIG. 1A).

Example 2. In Vivo CRISPR Killing of Target Strain Enterohemorrhagic *E. coli* (EHEC) In *Galleria mellonella* Larvae In Vivo Infection Model

2.1. CRISPR Efficacy Against *E. coli* (EHEC) ATCC43888 Infections in *Galleria mellonella*

CRISPR killing of target strain *E. coli* (EHEC) ATCC43888 was tested in *G. mellonella* in vivo infection model. To this aim, *G. mellonella* larvae were delivered injections of bacteria 10^8 CFU *E. coli* (EHEC) ATCC43888 behind the final left proleg. Approximately 1 h after the injection, CRISPR inducers (2 mM theophylline and 1% arabinose) were administered behind the final right proleg. Larvae were incubated at 37° C. and they were sacrificed after 1 and 2 h after induction. As shown in FIGS. 2 and 3, CRISPR induction killed 99% of the population after 2 h, as compared to the off-target control.

2.2. Survival Curves of *G. mellonella* Larvae Infected with Enterohemorrhagic *E. coli* (EHEC).

G. mellonella larvae were delivered injections of bacteria (8×10^4 CFU *E. coli* ATCC43888) behind the final left proleg. Approximately 1 h after the injection, CRISPR inducers (2 mM theophylline and 1% arabinose) were administered behind the final right proleg. Larvae were incubated at 37° C. and survival was monitored for 115 h, with death indicated by lack of movement and unresponsiveness to touch. CRISPR killing of target strain *E. coli* (EHEC) ATCC43888 in *G. mellonella* larvae significantly improved survival of the larvae compared to the off-target control (FIG. 4) (log-rank test, $P < 0.03$).

Example 3. Precision Killing of Target Strain Probiotic *E. coli* Nissle 1917

3.1. Design, Construction and Delivery of CRISPR Guided Vector (CGV) System Targeting *E. coli* Nissle 1917.

The invention provides a CGV system to specifically target *E. coli* Nissle 1917. The CGV system comprises two vectors: (a) a vector containing a tracrRNA and the Cas9 from *Streptococcus pyogenes* (SpCas9); (b) a vector containing a guide RNA (gRNA) that comprises a nucleotide sequence capable of hybridizing to a target sequence in the host cells to guide SpCas9 to the target sequence. To enable specific killing of *E. coli* Nissle 1917, a specific sequence from the genome of this strain was chosen to target. Specifically, the sequence contains 20 nucleotides from the pks gene from *E. coli* Nissle 1917. Additionally, the 5'-NGG protospacer adjacent motif (PAM) was located adjacent to the selected target sequence. The selected target sequence in the pks gene can be found in Table 3.

Furthermore, a different genome target was selected to specifically kill *E. coli* Nissle 1917. The sequence contains 20 nucleotides from the yapH gene. Additionally, the 5'-NGG protospacer adjacent motif (PAM) was located adjacent to the selected target sequence. The selected target sequence in yapH gene can be found in Table 3.

3.2. Construction and Delivery of CRISPR Guided Vectors (CGV) Targeting *E. coli* Nissle 1917

To establish CGVs functionality in mediating sequence-specific killing in *E. coli* Nissle 1917, the CGV system was transformed into *E. coli* Nissle 1917 cells. Overnight cultures were diluted 1:100 in fresh lysogeny broth (LB) and grown to mid-exponential phase OD₆₀₀~0.6. The CRISPR system was induced by adding theophylline and arabinose (2 mM theophylline and 1% arabinose), and survival of the strain was followed over time by plating the cultures in serial

dilutions every 15 minutes, for 3 h (FIGS. 5B and 6B). FIGS. 5B and 6A show CRISPR killing assay in *E. coli* Nissle 1917, targeting pks gene and yapH gene, respectively. In both cases, CRISPR induction triggers a rapid killing of *E. coli* Nissle 1917 cells, achieving 99.98% killing within only 15 minutes of induction.

Example 4. In Vitro CRISPR Killing of *Clostridium difficile* by Conjugative Plasmid Vectors and Cas3

This experiment involves the precision killing of *Clostridium difficile* using a gRNA-encoding CRISPR array that is delivered from a probiotic carrier bacterial species by conjugative plasmids as vectors (which we call CRISPR guided vectors (CGVTM)). A carrier bacterium (*E. coli* donor strain containing the CRISPR guided vector (CGVTM)) was mated with *Clostridium difficile* which was killed upon delivery of the CGVTM containing the designed array. This CGVTM harnessed the endogenous Cas3 machinery of *Clostridium difficile* 630Δerm. A 100% killing of *Clostridium difficile* cells was achieved.

Introduction

Clostridium difficile (*C. difficile*) is a spore-forming human opportunistic pathogen that can asymptotically colonize the intestine of healthy individuals. The two main risk factors for contracting *C. difficile*-associated diseases, such as nosocomial diarrhea, are age and antibiotic treatment and can have fatal consequences. *C. difficile* 630Δerm, the subject of our study, is a well-characterized strain and it is widely used for the generation of mutant specimens.

Study Objectives

Objective 1: Delivery of CGVs by Conjugation.

A CRISPR guided vector (CGV) containing an array to specifically target and kill *C. difficile* was designed and assembled. The same CGV lacking the array was assembled to use as a control for conjugation efficiency. Both CGVs were transformed into the carrier strain *Escherichia coli* CA434, which was used as a donor strain to conjugate the plasmid into our strain of interest *C. difficile* 630Δerm.

Objective 2: Harnessing *Clostridium difficile* Endogenous Cas3 Machinery.

Upon transcription of the delivered CRISPR array in the recipient target strain *C. difficile*, the endogenous Cas3 was guided to cut its own DNA, leading to bacterial death.

Objective 3: Eradication of *Clostridium difficile* 630Δerm.

Achievement of efficient killing of transconjugant *C. difficile* cells using designed CGVs.

Materials and Methods

Bacterial Strains and Growth Conditions

E. coli strain CA434 was acquired from Chain Biotech. It was cultured on nutrient-rich media (2xYT) and grown overnight at 37° C. and 250 rpm. Medium was supplemented with 12.5 μg/mL of thiamphenicol when required to maintain the CGVs.

Clostridium difficile 630Δerm was grown on BHI agar supplemented with 5 g/L of yeast extract, 0.03% L-cysteine, 250 μg/ml D-cycloserine and 8 μg/ml of cefoxitin (BHIS+CC). *C. difficile* was grown overnight in a Coy vinyl anaerobic cabinet in an atmosphere of 92% N₂, 6% CO₂ and 2% H₂ at 37° C. The mating of the donor CA434 and *C. difficile* was grown on plain BHI agar to allow for growth of the donor strain. Thiamphenicol was added to BHIS+CC plates to a final concentration of 12.5 μg/mL for selection of transconjugants after mating. All plates were dried for 1.5

hours and transferred, along with the broth version of this medium, to the anaerobic chamber at least 3 hours before use.

CGV Transfer Procedures

Carrier cells of *E. coli* CA434 were obtained by electroporation of either of our CGVs (control vector pMTL84151-FJ797649 and CRISPR vector pMTL84151-cdCRISPR1). In order to do that, overnight cultures of *E. coli* CA434 were diluted 1:100 in fresh 2x YT medium without selection and grown to OD600~0.5. Then, they were made electrocompetent by standard procedures (Sharan et al., 2009). Electrocompetent cells were transformed with either plasmid pMTL84151-FJ797649 or pMTL84151-cd-CRISPR1 and recovered in 2xYT for 1 h at 37° C. with shaking (250 rpm). Finally, they were plated on LB agar supplemented with 12.5 µg/mL thiamphenicol for selection of transformants. Transformants were grown in liquid 2xYT supplemented with 12.5 µg/mL thiamphenicol at 37° C. and 250 rpm for mating with *C. difficile*. 1 ml of donor cells was centrifuged at 4000×g for 2 minutes, supernatant removed and carefully washed with 400 µl of PBS. After a second centrifugation cycle the pellet was transferred to the anaerobic chamber for mating with *C. difficile* in BHI non-selective plates. *C. difficile* was prepared for mating following a modified protocol (Des Purdy et al., 2002). *C. difficile* 630Δerm was incubated overnight in selective BHIS+CC plates, from which, a scrape was inoculated overnight in 1 ml of non-selective BHI and incubated over night for mating. 200 µl of that culture was used to resuspend the pelleted donor cells and mixed culture was plated in 20 µl spots on top of non-selective BHI plates. The mating was incubated 24 h to allow for conjugation. After incubation, the whole plate was thoroughly scraped with a sterile inoculation loop, resuspended in BHI and serial dilutions were plated on BHI+CC plates to prevent growth of donor

E. coli and on BHI+CC supplemented with thiamphenicol for additional selection of transconjugants. Single colonies were counted after 48 hours.

Results

Replicates of BHI+CC+Thiamphenicol plates, selecting for *C. difficile* transconjugants carrying the control CGV, showed a consistent number of colonies resulting in about ~600-750 CFUs per mating experiment. For the mating of *C. difficile* with *E. coli* CA434 carrying the CGV with the CRISPR array the plates were empty, no colonies were observed. This translates into 100% killing of transconjugant *C. difficile* 630Δerm cells receiving the CRISPR array (see FIG. 7: Killing of transconjugant *C. difficile* 630Δerm).

DISCUSSION AND CONCLUSIONS

The results of this experiment show that we could successfully conjugate CGVs containing the desired CRISPR arrays into *C. difficile* 630Δerm from an *E. coli* carrier bacterium. We could also successfully harness *C. difficile* endogenous Cas3 machinery for very efficient CRISPR killing.

REFERENCES

- Purdy D, O'Keeffe T A, Elmore M, Herbert M, McLeod A, Bokori-Brown M, Ostrowski A, Minton N P. (2002) Conjugative transfer of clostridial shuttle vectors from *Escherichia coli* to *Clostridium difficile* through circumvention of the restriction barrier. *Molec. Microbiology* 46(2), 439-452
- Sharan, S. K., Thomason, L. C., Kuznetsov, S. G., and Court, D. L. (2009) Recombineering: a homologous recombination-based method of genetic engineering. *Nat. Protoc.* 4, 206-223

TABLE 1

Example Bacteria
Optionally, the bacteria are selected from this Table.

<i>Abiotrophia</i>	<i>Acidocella</i>	<i>Actinomyces</i>	<i>Alkalimimicola</i>	<i>Aquaspirillum</i>
<i>Abiotrophia defectiva</i>	<i>Acidocella aminolytica</i>	<i>Actinomyces bovis</i>	<i>Alkalimimicola ehrlichii</i>	<i>Aquaspirillum polymorphum</i>
<i>Acaricomes</i>	<i>Acidocella facilis</i>	<i>Actinomyces denticolens</i>	<i>Alkaliphilus</i>	<i>Aquaspirillum putridiconchylum</i>
<i>Acaricomes phytoseuii</i>	<i>Acidomonas</i>	<i>Actinomyces europaeus</i>	<i>Alkaliphilus oremlandii</i>	<i>Aquaspirillum serpens</i>
<i>Acetitomaculum</i>	<i>Acidomonas methanolica</i>	<i>Actinomyces georgiae</i>	<i>Alkaliphilus transvaalensis</i>	<i>Aquimarina</i>
<i>Acetitomaculum ruminis</i>	<i>Acidothermus</i>	<i>Actinomyces gerencseriae</i>	<i>Allochromatium</i>	<i>Aquimarina latercula</i>
<i>Acetivibrio</i>	<i>Acidothermus cellulolyticus</i>	<i>Actinomyces hordeovulneris</i>	<i>Allochromatium vinosum</i>	<i>Arcanobacterium haemolyticum</i>
<i>Acetivibrio cellulolyticus</i>	<i>Acidovorax</i>	<i>Actinomyces howellii</i>	<i>Alloiococcus</i>	<i>Arcanobacterium pyogenes</i>
<i>Acetivibrio ethanolognens</i>	<i>Acidovorax anthurii</i>	<i>Actinomyces hyovaginalis</i>	<i>Alloiococcus otitis</i>	<i>Archangium gephyra</i>
<i>Acetivibrio multivorans</i>	<i>Acidovorax caeni</i>	<i>Actinomyces israelii</i>	<i>Allokutzneria</i>	<i>Arcobacter</i>
<i>Acetoanaerobium</i>	<i>Acidovorax cattleayae</i>		<i>Allokutzneria alba</i>	<i>Arcobacter butzleri</i>
<i>Acetoanaerobium noterae</i>	<i>Acidovorax citrulli</i>	<i>Actinomyces johnsonii</i>	<i>Altererythrobacter</i>	<i>Arcobacter cryaerophilus</i>
<i>Acetobacter</i>	<i>Acidovorax defluvii</i>	<i>Actinomyces meyeri</i>	<i>Altererythrobacter ishigakiensis</i>	<i>Arcobacter halophilus</i>
<i>Acetobacter acetii</i>	<i>Acidovorax delafieldii</i>	<i>Actinomyces naeslundii</i>	<i>Altermonas</i>	<i>Arcobacter nitrofigilis</i>
<i>Acetobacter cerevisiae</i>	<i>Acidovorax facilis</i>	<i>Actinomyces neuii</i>	<i>Altermonas haloplanktis</i>	<i>Arcobacter skirrowii</i>
<i>Acetobacter cibirongensis</i>	<i>Acidovorax konjaci</i>	<i>Actinomyces odontolyticus</i>	<i>Altermonas macleodii</i>	<i>Arhodomonas</i>
<i>Acetobacter estunensis</i>	<i>Acidovorax temperans</i>	<i>Actinomyces oris</i>	<i>Alysiella</i>	<i>Arhodomonas aquaeolei</i>
<i>Acetobacter fabarum</i>	<i>Acidovorax valerianellae</i>	<i>Actinomyces radinae</i>	<i>Alysiella crassa</i>	<i>Arsenophonus</i>
<i>Acetobacter ghanensis</i>	<i>Acinetobacter</i>	<i>Actinomyces slackii</i>	<i>Alysiella filiformis</i>	<i>Arsenophonus nasoniae</i>
<i>Acetobacter indonesiensis</i>	<i>Acinetobacter baumannii</i>	<i>Actinomyces turicensis</i>	<i>Aminobacter</i>	<i>Arthrobacter</i>
<i>Acetobacter lovaniensis</i>	<i>Acinetobacter baylyi</i>	<i>Actinomyces viscosus</i>	<i>Aminobacter aganoensis</i>	<i>Arthrobacter agilis</i>
<i>Acetobacter malorum</i>	<i>Acinetobacter bouvetii</i>	<i>Actinoplanes</i>	<i>Aminobacter aminovorans</i>	<i>Arthrobacter albus</i>
<i>Acetobacter nitrogenifigens</i>	<i>Acinetobacter calcoaceticus</i>	<i>Actinoplanes auranticolor</i>	<i>Aminobacter nigataensis</i>	<i>Arthrobacter aurescens</i>
<i>Acetobacter oeni</i>	<i>Acinetobacter gernerii</i>	<i>Actinoplanes brasiliensis</i>	<i>Aminobacterium</i>	<i>Arthrobacter chlorophenolicus</i>
<i>Acetobacter orientalis</i>	<i>Acinetobacter haemolyticus</i>	<i>Actinoplanes consetensis</i>	<i>Aminobacterium mobile</i>	
<i>Acetobacter orleanensis</i>	<i>Acinetobacter johnsonii</i>	<i>Actinoplanes decanensis</i>	<i>Aminomonas</i>	
<i>Acetobacter pasteurianus</i>	<i>Acinetobacter junii</i>	<i>Actinoplanes derwentensis</i>	<i>Aminomonas paucivorans</i>	
<i>Acetobacter pomorum</i>	<i>Acinetobacter lwoffii</i>	<i>Actinoplanes digitatis</i>	<i>Ammophilus</i>	

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>Acetobacter senegalensis</i>	<i>Acinetobacter parvus</i>	<i>Actinoplanes durhamensis</i>	<i>Ammoniphilus oxalaticus</i>	<i>Arthrobacter citreus</i>
<i>Acetobacter xylinus</i>	<i>Acinetobacter radioresistens</i>	<i>Actinoplanes ferrugineus</i>	<i>Ammoniphilus oxalivorans</i>	<i>Arthrobacter crystallopoietes</i>
<i>Acetobacterium</i>	<i>Acinetobacter schindleri</i>	<i>Actinoplanes globisporus</i>	<i>Amphibacillus</i>	<i>Arthrobacter cumminsii</i>
<i>Acetobacterium bakii</i>	<i>Acinetobacter soli</i>	<i>Actinoplanes humidus</i>	<i>Amphibacillus xylanus</i>	<i>Arthrobacter globiformis</i>
<i>Acetobacterium carbinolicum</i>	<i>Acinetobacter tandoii</i>	<i>Actinoplanes italicus</i>	<i>Amphritea</i>	<i>Arthrobacter</i>
<i>Acetobacterium dehalogenans</i>	<i>Acinetobacter tjernbergiae</i>	<i>Actinoplanes liguriensis</i>	<i>Amphritea balanae</i>	<i>Arthrobacter histidinovorans</i>
<i>Acetobacterium fimetarium</i>	<i>Acinetobacter townneri</i>	<i>Actinoplanes lobatus</i>	<i>Amphritea japonica</i>	<i>Arthrobacter ilicis</i>
<i>Acetobacterium malicum</i>	<i>Acinetobacter ursingii</i>	<i>Actinoplanes missouriensis</i>	<i>Amycolatopsis</i>	<i>Arthrobacter luteus</i>
<i>Acetobacterium paludosum</i>	<i>Acinetobacter venetianus</i>	<i>Actinoplanes palleronii</i>	<i>Amycolatopsis alba</i>	<i>Arthrobacter methylophilus</i>
<i>Acetobacterium tundrae</i>	<i>Acrocarpospora</i>	<i>Actinoplanes philippinensis</i>	<i>Amycolatopsis albidoflavus</i>	<i>Arthrobacter mysorens</i>
<i>Acetobacterium wieringae</i>	<i>Acrocarpospora corrugata</i>	<i>Actinoplanes rectilineatus</i>	<i>Amycolatopsis azurea</i>	<i>Arthrobacter nicotianae</i>
<i>Acetobacterium woodii</i>	<i>Acrocarpospora</i>	<i>Actinoplanes regularis</i>	<i>Amycolatopsis coloradensis</i>	<i>Arthrobacter nicotinovorans</i>
<i>Acetofilamentum</i>	<i>macrocephala</i>	<i>Actinoplanes</i>	<i>Amycolatopsis lurida</i>	<i>Arthrobacter oxydans</i>
<i>Acetofilamentum rigidum</i>	<i>Acrocarpospora pleiomorpha</i>	<i>teichomyceticus</i>	<i>Amycolatopsis mediterranei</i>	<i>Arthrobacter pascens</i>
<i>Acetohalobium</i>	<i>Actibacter</i>	<i>Actinoplanes utahensis</i>	<i>Amycolatopsis rifamycinica</i>	<i>Arthrobacter</i>
<i>Acetohalobium arabaticum</i>	<i>Actibacter sediminis</i>	<i>Actinopolyspora</i>	<i>Amycolatopsis rubida</i>	<i>Arthrobacter phenanthrenivorans</i>
<i>Acetomicrobium</i>	<i>Actinoalloteichus</i>	<i>Actinopolyspora halophila</i>	<i>Amycolatopsis sulphurea</i>	<i>Arthrobacter</i>
<i>Acetomicrobium faecale</i>	<i>Actinoalloteichus</i>	<i>Actinopolyspora mortivallis</i>	<i>Amycolatopsis tolypomycinica</i>	<i>polychromogenes</i>
<i>Acetomicrobium flavidum</i>	<i>cyanogriseus</i>	<i>Actinosynnema</i>	<i>Anabaena</i>	<i>Arthrobacter protophormiae</i>
<i>Acetonema</i>	<i>Actinoalloteichus</i>	<i>Actinosynnema mirum</i>	<i>Anabaena cylindrica</i>	<i>Arthrobacter</i>
<i>Acetonema longum</i>	<i>hymeniacidonis</i>	<i>Actinotalea</i>	<i>Anabaena flos-aquae</i>	<i>psychrolactophilus</i>
<i>Acetothermus</i>	<i>Actinoalloteichus spitiensis</i>	<i>fermentans</i>	<i>Anabaena variabilis</i>	<i>Arthrobacter ramosus</i>
<i>Acetothermus paucivorans</i>	<i>Actinobacillus</i>	<i>Aerococcus</i>	<i>Anaerococcus</i>	<i>Arthrobacter sulfonivorans</i>
<i>Acholeplasma</i>	<i>Actinobacillus capsulatus</i>	<i>Aerococcus sanguinicola</i>	<i>Anaerococcus burkinensis</i>	<i>Arthrobacter sulfureus</i>
<i>Acholeplasma axanthum</i>	<i>Actinobacillus delphinicola</i>	<i>Aerococcus urinae</i>	<i>Anaerobaculum</i>	<i>Arthrobacter uratoxydans</i>
<i>Acholeplasma brassicae</i>	<i>Actinobacillus hominis</i>	<i>Aerococcus urinaequi</i>	<i>Anaerobaculum mobile</i>	<i>Arthrobacter ureafaciens</i>
<i>Acholeplasma cavigenitalium</i>	<i>Actinobacillus indolicus</i>	<i>Aerococcus urinaehomins</i>	<i>Anaerobiospirillum</i>	<i>Arthrobacter viscosus</i>
<i>Acholeplasma equifetale</i>	<i>Actinobacillus lignieresii</i>	<i>Aerococcus viridans</i>	<i>Anaerobiospirillum</i>	<i>Arthrobacter woluwensis</i>
<i>Acholeplasma granularum</i>	<i>Actinobacillus minor</i>	<i>Aeromicrobium</i>	<i>succiniciproducens</i>	<i>Asaia</i>
<i>Acholeplasma hippikon</i>	<i>Actinobacillus muris</i>	<i>Aeromicrobium erythreum</i>	<i>Anaerobiospirillum thomasi</i>	<i>Asaia bogorensis</i>
<i>Acholeplasma laidlawii</i>	<i>Actinobacillus</i>	<i>Aeromonas</i>	<i>Anaerococcus</i>	<i>Asanao</i>
<i>Acholeplasma modicum</i>	<i>pleuropneumoniae</i>	<i>Aeromonas</i>	<i>Anaerococcus hydrogenalis</i>	<i>Asanao ferruginea</i>
<i>Acholeplasma morum</i>	<i>Actinobacillus porcini</i>	<i>allosaccharophila</i>	<i>Anaerococcus lactolyticus</i>	<i>Asticcacaulis</i>
<i>Acholeplasma multilocale</i>	<i>Actinobacillus rossii</i>	<i>Aeromonas bestiarum</i>	<i>Anaerococcus prevotii</i>	<i>Asticcacaulis biprothecium</i>
<i>Acholeplasma oculi</i>	<i>Actinobacillus scotiae</i>	<i>Aeromonas caviae</i>	<i>Anaerococcus tetradius</i>	<i>Asticcacaulis excentricus</i>
<i>Acholeplasma palmae</i>	<i>Actinobacillus seminis</i>	<i>Aeromonas encheleia</i>	<i>Anaerococcus vaginalis</i>	<i>Atopobacter</i>
<i>Acholeplasma parvum</i>	<i>Actinobacillus succinogenes</i>	<i>Aeromonas</i>	<i>Anaerofustis</i>	<i>Atopobacter phocae</i>
<i>Acholeplasma plecicae</i>	<i>Actinobacillus suis</i>	<i>enteropelogenes</i>	<i>Anaerofustis stercorihominis</i>	<i>Atopobium</i>
<i>Acholeplasma vituli</i>	<i>Actinobacillus ureae</i>	<i>Aeromonas eucrenophila</i>	<i>Anaeromusa</i>	<i>Atopobium fossor</i>
<i>Achromobacter</i>	<i>Actinobaculum</i>	<i>Aeromonas ichtiosismia</i>	<i>Anaeromusa acidaminophila</i>	<i>Atopobium minutum</i>
<i>Achromobacter denitrificans</i>	<i>Actinobaculum massiliense</i>	<i>Aeromonas jandaei</i>	<i>Anaeromyxobacter</i>	<i>Atopobium parvulum</i>
<i>Achromobacter insolitus</i>	<i>Actinobaculum schaalii</i>	<i>Aeromonas media</i>	<i>Anaeromyxobacter</i>	<i>Atopobium rimae</i>
<i>Achromobacter piechaudii</i>	<i>Actinobaculum suis</i>	<i>Aeromonas popoffii</i>	<i>dehalogenans</i>	<i>Atopobium vaginae</i>
<i>Achromobacter ruhlandii</i>	<i>Actinomyces urinale</i>	<i>Aeromonas sobria</i>	<i>Anaerorhabdus</i>	<i>Aureobacterium</i>
<i>Achromobacter spanius</i>	<i>Actinocatenispora</i>	<i>Aeromonas veronii</i>	<i>Anaerorhabdus furcosa</i>	<i>Aureobacterium barkeri</i>
<i>Acidaminobacter</i>	<i>Actinocatenispora rupis</i>	<i>Agrobacterium</i>	<i>Anaerosinus</i>	<i>Aureobacterium</i>
<i>Acidaminobacter</i>	<i>Actinocatenispora</i>	<i>Agrobacterium</i>	<i>Anaerosinus glycerini</i>	<i>liquefaciens</i>
<i>hydrogenoformans</i>	<i>thailandica</i>	<i>gelatinovorum</i>	<i>Anaerovirgula</i>	<i>Avibacterium</i>
<i>Acidaminococcus</i>	<i>Actinocatenispora sera</i>	<i>Agrococcus</i>	<i>Anaerovirgula multivorans</i>	<i>Avibacterium avium</i>
<i>Acidaminococcus fermentans</i>	<i>Actinocorallia</i>	<i>Agrococcus citreus</i>	<i>Ancalomicrobium</i>	<i>Avibacterium gallinarum</i>
<i>Acidaminococcus intestini</i>	<i>Actinocorallia aurantiaca</i>	<i>Agrococcus jenensis</i>	<i>Ancalomicrobium adetum</i>	<i>Avibacterium paragallinarum</i>
<i>Acidicaldus</i>	<i>Actinocorallia aurea</i>	<i>Agromonas</i>	<i>Ancylobacter</i>	<i>Avibacterium volantium</i>
<i>Acidicaldus organivorans</i>	<i>Actinocorallia cavernae</i>	<i>Agromonas oligotrophica</i>	<i>Ancylobacter aquaticus</i>	<i>Azoarcus</i>
<i>Acidimicrobium</i>	<i>Actinocorallia glomerata</i>	<i>Agromyces</i>	<i>Aneurinibacillus</i>	<i>Azoarcus indigenus</i>
<i>Acidimicrobium ferrooxidans</i>	<i>Actinocorallia herbida</i>	<i>Agromyces fucosus</i>	<i>Aneurinibacillus aneurinilyticus</i>	<i>Azoarcus toluolyticus</i>
<i>Acidiphilium</i>	<i>Actinocorallia libanotica</i>	<i>Agromyces hippuratus</i>	<i>Aneurinibacillus migulanus</i>	<i>Azoarcus toluovorans</i>
<i>Acidiphilium acidophilum</i>	<i>Actinocorallia longicatena</i>	<i>Agromyces luteolus</i>	<i>Aneurinibacillus thermoaerophilus</i>	<i>Azohydromonas</i>
<i>Acidiphilium angustum</i>	<i>Actinomadura</i>	<i>Agromyces mediolanus</i>		<i>Azohydromonas australica</i>
<i>Acidiphilium cryptum</i>	<i>Actinomadura alba</i>	<i>Agromyces ramosus</i>	<i>Angiococcus</i>	<i>Azohydromonas lata</i>
<i>Acidiphilium multivorum</i>	<i>Actinomadura atramentaria</i>	<i>Agromyces rhizosphaerae</i>	<i>Angiococcus disciformis</i>	<i>Azomonas</i>
<i>Acidiphilium organovorum</i>	<i>Actinomadura</i>	<i>Akkermansia</i>	<i>Angulomicrobium</i>	<i>Azomonas agilis</i>
<i>Acidiphilium rubrum</i>	<i>bangladeshensis</i>	<i>Akkermansia muciniphila</i>	<i>Angulomicrobium tetraedrale</i>	<i>Azomonas insignis</i>
<i>Acidisoma</i>	<i>Actinomadura catellatispora</i>	<i>Albidiferax</i>	<i>Anoxybacillus</i>	<i>Azomonas macrocytogenes</i>

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>Acidisoma sibiricum</i>	<i>Actinomadura chibensis</i>	<i>Albidiferax ferrireducens</i>	<i>Anoxybacillus pushchinoensis</i>	<i>Azorhizobium</i>
<i>Acidisoma tundrae</i>	<i>Actinomadura chokoriensis</i>	<i>Albidovulum</i>	<i>Aquabacterium</i>	<i>Azorhizobium caulinodans</i>
<i>Acidisphaera</i>	<i>Actinomadura citrea</i>	<i>Albidovulum inexpectatum</i>	<i>Aquabacterium commune</i>	<i>Azorhizophilus paspali</i>
<i>Acidisphaera rubrifaciens</i>	<i>Actinomadura coerulea</i>	<i>Alcaligenes</i>	<i>Aquabacterium parvum</i>	<i>Azospirillum</i>
<i>Acidithiobacillus</i>	<i>Actinomadura echinospora</i>	<i>Alcaligenes denitrificans</i>		<i>Azospirillum brasilense</i>
<i>Acidithiobacillus albertensis</i>	<i>Actinomadura fibrosa</i>	<i>Alcaligenes faecalis</i>		<i>Azospirillum halopraeferens</i>
<i>Acidithiobacillus caldus</i>	<i>Actinomadura formosensis</i>	<i>Alcanivorax</i>		<i>Azospirillum irakense</i>
<i>Acidithiobacillus ferrooxidans</i>	<i>Actinomadura hibisca</i>	<i>Alcanivorax borkumensis</i>		<i>Azotobacter</i>
<i>Acidithiobacillus thiooxidans</i>	<i>Actinomadura kijaniata</i>	<i>Alcanivorax jadensis</i>		<i>Azotobacter beijerinckii</i>
<i>Acidobacterium</i>	<i>Actinomadura latina</i>	<i>Algicola</i>		<i>Azotobacter chroococcum</i>
<i>Acidobacterium capsulatum</i>	<i>Actinomadura livida</i>	<i>Algicola bacteriolytica</i>		<i>Azotobacter nigricans</i>
	<i>Actinomadura luteofluorescens</i>	<i>Alicyclobacillus</i>		<i>Azotobacter salinestrans</i>
	<i>Actinomadura macra</i>	<i>Alicyclobacillus disulfidooxidans</i>		<i>Azotobacter vinelandii</i>
	<i>Actinomadura madurae</i>	<i>Alicyclobacillus sendaiensis</i>		
	<i>Actinomadura oligospora</i>	<i>Alicyclobacillus vulcanalis</i>		
	<i>Actinomadura pelletieri</i>	<i>Alishewanella</i>		
	<i>Actinomadura rubrobrunea</i>	<i>Alishewanella fetalis</i>		
	<i>Actinomadura rugatobispora</i>	<i>Alkalibacillus</i>		
	<i>Actinomadura umbrina</i>	<i>Alkalibacillus haloalkaliphilus</i>		
	<i>Actinomadura verrucosospora</i>			
	<i>Actinomadura vinacea</i>			
	<i>Actinomadura viridilutea</i>			
	<i>Actinomadura viridis</i>			
	<i>Actinomadura yumaensis</i>			
<i>Bacillus</i>	<i>Bacteroides</i>	<i>Bibersteinia</i>	<i>Borrelia</i>	<i>Brevinema</i>
[see below]	<i>Bacteroides caccae</i>	<i>Bibersteinia trehalosi</i>	<i>Borrelia afzelii</i>	<i>Brevinema andersonii</i>
<i>Bacteriovorax</i>	<i>Bacteroides coagulans</i>	<i>Bifidobacterium</i>	<i>Borrelia americana</i>	<i>Brevundimonas</i>
<i>Bacteriovorax stolpii</i>	<i>Bacteroides eggerthii</i>	<i>Bifidobacterium adolescentis</i>	<i>Borrelia burgdorferi</i>	<i>Brevundimonas alba</i>
	<i>Bacteroides fragilis</i>	<i>Bifidobacterium angulatum</i>	<i>Borrelia carolinensis</i>	<i>Brevundimonas aurantiaca</i>
	<i>Bacteroides galacturonicus</i>	<i>Bifidobacterium animalis</i>	<i>Borrelia coriacea</i>	<i>Brevundimonas diminuta</i>
	<i>Bacteroides helcogenes</i>	<i>Bifidobacterium asteroides</i>	<i>Borrelia garinii</i>	<i>Brevundimonas intermedia</i>
	<i>Bacteroides ovatus</i>	<i>Bifidobacterium bifidum</i>	<i>Borrelia japonica</i>	<i>Brevundimonas subvibrioides</i>
	<i>Bacteroides pectinophilus</i>	<i>Bifidobacterium bourn</i>	<i>Bosea</i>	<i>Brevundimonas vancouverii</i>
	<i>Bacteroides pyogenes</i>	<i>Bifidobacterium breve</i>	<i>Bosea minaitlanensis</i>	<i>Brevundimonas variabilis</i>
	<i>Bacteroides salyersiae</i>	<i>Bifidobacterium catenulatum</i>	<i>Bosea thiooxidans</i>	<i>Brevundimonas vesicularis</i>
	<i>Bacteroides stercoris</i>	<i>Bifidobacterium choerinum</i>	<i>Brachybacterium</i>	<i>Brochothrix</i>
	<i>Bacteroides suis</i>	<i>Bifidobacterium coryneforme</i>	<i>Brachybacterium alimentarium</i>	<i>Brochothrix campestris</i>
	<i>Bacteroides tectus</i>	<i>Bifidobacterium cuculi</i>		<i>Brochothrix thermosphacta</i>
	<i>Bacteroides thetaiotaomicron</i>	<i>Bifidobacterium dentium</i>	<i>Brachybacterium faecium</i>	<i>Brucella</i>
	<i>Bacteroides uniformis</i>	<i>Bifidobacterium gallicum</i>	<i>Brachybacterium paraconglomeratum</i>	<i>Brucella canis</i>
	<i>Bacteroides ureolyticus</i>	<i>Bifidobacterium gallinarum</i>	<i>Brachybacterium rhamnosum</i>	<i>Brucella neotomae</i>
	<i>Bacteroides vulgatus</i>	<i>Bifidobacterium indicum</i>	<i>Brachybacterium tyrofermentans</i>	<i>Bryobacter</i>
	<i>Balnearium</i>	<i>Bifidobacterium longum</i>	<i>Brachybacterium magnum</i>	<i>Bryobacter aggregatus</i>
	<i>Balnearium lithotrophicum</i>	<i>Bifidobacterium merycicum</i>	<i>Brachyspira</i>	<i>Burkholderia</i>
	<i>Balnearix</i>		<i>Brachyspira alvinipulli</i>	<i>Burkholderia ambifaria</i>
	<i>Balnearix alpica</i>			<i>Burkholderia andropogonis</i>
	<i>Balneola</i>	<i>Bifidobacterium minimum</i>	<i>Brachyspira hyodysenteriae</i>	<i>Burkholderia anthina</i>
	<i>Balneola vulgaris</i>	<i>Bifidobacterium pseudocatenulatum</i>	<i>Brachyspira innocens</i>	<i>Burkholderia caledonica</i>
	<i>Barnesiella</i>	<i>Bifidobacterium</i>	<i>Brachyspira murdochii</i>	<i>Burkholderia caryophylli</i>
	<i>Barnesiella viscericola</i>		<i>Brachyspira pilosicoli</i>	<i>Burkholderia cenocepacia</i>
	<i>Bartonella</i>	<i>pseudolongum</i>	<i>Bradyrhizobium</i>	<i>Burkholderia cepacia</i>
	<i>Bartonella alsatica</i>	<i>Bifidobacterium pullorum</i>	<i>Bradyrhizobium canariense</i>	<i>Burkholderia cocovenenans</i>
	<i>Bartonella bacilliformis</i>	<i>Bifidobacterium ruminantium</i>	<i>Bradyrhizobium elkanii</i>	<i>Burkholderia dolosa</i>
	<i>Bartonella clarridgeiae</i>	<i>Bifidobacterium saeculare</i>	<i>Bradyrhizobium japonicum</i>	<i>Burkholderia fungorum</i>
	<i>Bartonella doshae</i>	<i>Bifidobacterium subtile</i>	<i>Bradyrhizobium liaoningense</i>	<i>Burkholderia glathei</i>
	<i>Bartonella elizabethae</i>	<i>Bifidobacterium thermophilum</i>	<i>Brenneria</i>	<i>Burkholderia glumae</i>
	<i>Bartonella grahamii</i>	<i>Bilophila</i>	<i>Brenneria alni</i>	<i>Burkholderia graminis</i>
	<i>Bartonella henselae</i>	<i>Bilophila wadsworthia</i>	<i>Brenneria nigrifluens</i>	<i>Burkholderia kururiensis</i>
	<i>Bartonella rochalimae</i>		<i>Brenneria quercina</i>	<i>Burkholderia multivorans</i>
	<i>Bartonella vinsonii</i>	<i>Biostraticola</i>	<i>Brenneria quercina</i>	<i>Burkholderia phenazinium</i>

TABLE 1-continued

Example Bacteria			
Optionally, the bacteria are selected from this Table.			
<i>Bavariococcus</i>	<i>Biostraticola tofi</i>	<i>Brenneria salicis</i>	<i>Burkholderia plantarii</i>
<i>Bavariococcus seileri</i>	<i>Bizionia</i>	<i>Brevibacillus</i>	<i>Burkholderia pyrrocinia</i>
<i>Bdellovibrio</i>	<i>Bizionia argentinensis</i>	<i>Brevibacillus agri</i>	<i>Burkholderia silvaticola</i>
<i>Bdellovibrio bacteriovorus</i>	<i>Blastobacter</i>	<i>Brevibacillus borstelensis</i>	<i>Burkholderia stabilis</i>
<i>Bdellovibrio exovorus</i>	<i>Blastobacter capsulatus</i>	<i>Brevibacillus brevis</i>	<i>Burkholderia thailandensis</i>
<i>Beggiatoa</i>	<i>Blastobacter denitrificans</i>	<i>Brevibacillus centrosporus</i>	<i>Burkholderia tropica</i>
<i>Beggiatoa alba</i>	<i>Blastococcus</i>	<i>Brevibacillus choshinensis</i>	<i>Burkholderia unamae</i>
<i>Beijerinckia</i>	<i>Blastococcus aggregatus</i>	<i>Brevibacillus invocatus</i>	<i>Burkholderia vietnamiensis</i>
<i>Beijerinckia dextrii</i>	<i>Blastococcus saxosidens</i>	<i>Brevibacillus laterosporus</i>	<i>Buttiauxella</i>
<i>Beijerinckia fluminensis</i>	<i>Blastochloris</i>	<i>Brevibacillus parabrevis</i>	<i>Buttiauxella agrestis</i>
<i>Beijerinckia indica</i>	<i>Blastochloris viridis</i>	<i>Brevibacillus reuszeri</i>	<i>Buttiauxella brennerae</i>
<i>Beijerinckia mobilis</i>	<i>Blastomonas</i>	<i>Brevibacterium</i>	<i>Buttiauxella ferruginea</i>
<i>Belliella</i>	<i>Blastomonas natatoria</i>	<i>Brevibacterium abidum</i>	<i>Buttiauxella gaviniae</i>
<i>Belliella baltica</i>	<i>Blastopirellula</i>	<i>Brevibacterium album</i>	<i>Buttiauxella izardii</i>
<i>Bellilinea</i>	<i>Blastopirellula marina</i>	<i>Brevibacterium aurantiacum</i>	<i>Buttiauxella noackiae</i>
<i>Bellilinea caldifistulae</i>	<i>Blautia</i>	<i>Brevibacterium celere</i>	<i>Buttiauxella warmboldiae</i>
<i>Belnapia</i>	<i>Blautia coccoides</i>	<i>Brevibacterium epidermidis</i>	<i>Butyrivibrio</i>
<i>Belnapia moabensis</i>	<i>Blautia hansenii</i>	<i>Brevibacterium</i>	<i>Butyrivibrio fibrisolvens</i>
<i>Bergeriella</i>	<i>Blautia producta</i>	<i>frigoritolerans</i>	<i>Butyrivibrio hungatei</i>
<i>Bergeriella denitrificans</i>	<i>Blautia wexlerae</i>	<i>Brevibacterium halotolerans</i>	<i>Butyrivibrio proteoclasticus</i>
<i>Beutenbergia</i>	<i>Bogoriella</i>	<i>Brevibacterium iodinum</i>	
<i>Beutenbergia cavernae</i>	<i>Bogoriella caseilytica</i>	<i>Brevibacterium linens</i>	
	<i>Bordetella</i>	<i>Brevibacterium lyticum</i>	
	<i>Bordetella avium</i>	<i>Brevibacterium mcbrellneri</i>	
	<i>Bordetella bronchiseptica</i>	<i>Brevibacterium oitidis</i>	
	<i>Bordetella hinzii</i>	<i>Brevibacterium oxydans</i>	
	<i>Bordetella holmesii</i>	<i>Brevibacterium paucivorans</i>	
	<i>Bordetella parapertussis</i>	<i>Brevibacterium stationis</i>	
	<i>Bordetella pertussis</i>		
	<i>Bordetella petrii</i>		
	<i>Bordetella trematum</i>		
<i>Bacillus</i>			
<i>B. acidicer</i>	<i>B. aminovorans</i>	<i>B. glucanolyticus</i>	<i>B. taeanensis</i>
<i>B. acidicola</i>	<i>B. amylolyticus</i>	<i>B. gordonae</i>	<i>B. tequilensis</i>
<i>B. acidiproducens</i>	<i>B. andresenii</i>	<i>B. gotheii</i>	<i>B. thermantarcticus</i>
<i>B. acidocaldarius</i>	<i>B. aneurinilyticus</i>	<i>B. graminis</i>	<i>B. thermoaerophilus</i>
<i>B. acidoterrestris</i>	<i>B. anthracis</i>	<i>B. halmapalus</i>	<i>B. thermoamylovorans</i>
<i>B. aeolius</i>	<i>B. aquimaris</i>	<i>B. haloalkaliphilus</i>	<i>B. thermocatenulatus</i>
<i>B. aerius</i>	<i>B. arenosi</i>	<i>B. halochares</i>	<i>B. thermocloacae</i>
<i>B. aerophilus</i>	<i>B. arseniciselenatis</i>	<i>B. halodenitrificans</i>	<i>B. thermocopriae</i>
<i>B. agaradhaerens</i>	<i>B. arsenicus</i>	<i>B. halodurans</i>	<i>B. thermodenitrificans</i>
<i>B. agri</i>	<i>B. aurantiacus</i>	<i>B. halophilus</i>	<i>B. thermoglucoisidarius</i>
<i>B. aitingensis</i>	<i>B. arvi</i>	<i>B. halosaccharovorans</i>	<i>B. thermolactis</i>
<i>B. akibai</i>	<i>B. aryabhatai</i>	<i>B. hemicellulosilyticus</i>	<i>B. thermoleovorans</i>
<i>B. alcalophilus</i>	<i>B. asahii</i>	<i>B. hemisphaericus</i>	<i>B. thermophilus</i>
<i>B. algicola</i>	<i>B. atrophaeus</i>	<i>B. herbersteinensis</i>	<i>B. thermoruber</i>
<i>B. alginolyticus</i>	<i>B. axarquiensis</i>	<i>B. horikoshii</i>	<i>B. thermosphaericus</i>
<i>B. alkalidiazotrophicus</i>	<i>B. azotofixans</i>	<i>B. horneckiae</i>	<i>B. thiaminolyticus</i>
<i>B. alkalinitrilicus</i>	<i>B. azotoformans</i>	<i>B. horti</i>	<i>B. thioparans</i>
<i>B. alkalisediminis</i>	<i>B. badius</i>	<i>B. huizhouensis</i>	<i>B. thuringiensis</i>
<i>B. alkalitelluris</i>	<i>B. barbaricus</i>	<i>B. humi</i>	<i>B. tianshenii</i>
<i>B. altitudinis</i>	<i>B. bataviensis</i>	<i>B. hwajinpoensis</i>	<i>B. trypoxylicola</i>
<i>B. alveayuensis</i>	<i>B. beijingensis</i>	<i>B. idriensis</i>	<i>B. tusciae</i>
<i>B. alvei</i>	<i>B. benzoovorans</i>	<i>B. indicus</i>	<i>B. validus</i>
<i>B. amyloliquefaciens</i>	<i>B. beringensis</i>	<i>B. infantis</i>	<i>B. vallismortis</i>
<i>B.</i>	<i>B. berkeleyi</i>	<i>B. infernus</i>	<i>B. vedderi</i>
<i>a. subsp. amyloliquefaciens</i>	<i>B. beveridgei</i>	<i>B. insolitus</i>	<i>B. velezensis</i>
<i>B. a. subsp. plantarum</i>	<i>B. bogoriensis</i>	<i>B. invictae</i>	<i>B. vietnamensis</i>
<i>B. dipososauri</i>	<i>B. boroniphilus</i>	<i>B. iranensis</i>	<i>B. vireti</i>
<i>B. drentensis</i>	<i>B. borstelensis</i>	<i>B. isabelliae</i>	<i>B. vulcani</i>
<i>B. edaphicus</i>	<i>B. brevis Migula</i>	<i>B. isronensis</i>	<i>B. wakoensis</i>
<i>B. ehimensis</i>	<i>B. butanolivorans</i>	<i>B. jeotgali</i>	<i>B. weihenstephanensis</i>
<i>B. eiseniae</i>	<i>B. canaveralius</i>	<i>B. kaustophilus</i>	<i>B. xiamenensis</i>
<i>B. enclensis</i>	<i>B. carboniphilus</i>	<i>B. kobensis</i>	<i>B. xiaoxiensis</i>
<i>B. endophyticus</i>	<i>B. cecembensis</i>	<i>B. kochii</i>	<i>B. zhanjiangensis</i>
<i>B. endoradicis</i>	<i>B. cellulosityticus</i>	<i>B. kokeshiformis</i>	<i>B. peoriae</i>
<i>B. farraginis</i>	<i>B. centrosporus</i>	<i>B. koreensis</i>	<i>B. persepolensis</i>
<i>B. fastidiosus</i>	<i>B. cereus</i>	<i>B. korlensis</i>	<i>B. persicus</i>
<i>B. fengqiensis</i>	<i>B. chagannorensis</i>	<i>B. kribbensis</i>	<i>B. pervagus</i>
<i>B. firmus</i>	<i>B. chitinolyticus</i>	<i>B. krulwichiae</i>	<i>B. plakortidis</i>
<i>B. flexus</i>	<i>B. chondroitinus</i>	<i>B. laevolacticus</i>	<i>B. pocheonensis</i>
<i>B. foraminis</i>	<i>B. choshinensis</i>	<i>B. larvae</i>	<i>B. polygona</i>

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>B. fordii</i>	<i>B. chungangensis</i>	<i>B. laterosporus</i>	<i>B. polymyxa</i>	<i>B. oceanisediminis</i>
<i>B. formosus</i>	<i>B. cibi</i>	<i>B. salexigens</i>	<i>B. popilliae</i>	<i>B. odysseyi</i>
<i>B. fortis</i>	<i>B. circulans</i>	<i>B. saliphilus</i>	<i>B. pseudalcalophilus</i>	<i>B. okhensis</i>
<i>B. fumarioli</i>	<i>B. clarkii</i>	<i>B. schlegelii</i>	<i>B. pseudofirmus</i>	<i>B. okuhidensis</i>
<i>B. funiculus</i>	<i>B. clausii</i>	<i>B. sediminis</i>	<i>B. pseudomycooides</i>	<i>B. oleronius</i>
<i>B. fusiiformis</i>	<i>B. coagulans</i>	<i>B. selenatarsenatis</i>	<i>B. psychrodurans</i>	<i>B. oryzaecorticis</i>
<i>B. galactophilus</i>	<i>B. coahuilensis</i>	<i>B. selenitireducens</i>	<i>B. psychrophilus</i>	<i>B. oshimensis</i>
<i>B. galactosidilyticus</i>	<i>B. cohnii</i>	<i>B. seohaeanensis</i>	<i>B. psychrosaccharolyticus</i>	<i>B. pabuli</i>
<i>B. galliciensis</i>	<i>B. composti</i>	<i>B. shacheensis</i>	<i>B. psychrotolerans</i>	<i>B. pakistanensis</i>
<i>B. gelatini</i>	<i>B. curdolanolyticus</i>	<i>B. shackletonii</i>	<i>B. pulvifaciens</i>	<i>B. pallidus</i>
<i>B. gibsonii</i>	<i>B. cycloheptanicus</i>	<i>B. siamensis</i>	<i>B. pumilus</i>	<i>B. pallidus</i>
<i>B. ginsengi</i>	<i>B. cytotoxicus</i>	<i>B. silvestris</i>	<i>B. purgationiresistens</i>	<i>B. panacisoli</i>
<i>B. ginsengihumi</i>	<i>B. daliensis</i>	<i>B. simplex</i>	<i>B. pycnus</i>	<i>B. panaciterrae</i>
<i>B. ginsengisoli</i>	<i>B. decisifrondis</i>	<i>B. siralis</i>	<i>B. qingdaonensis</i>	<i>B. pantothenicis</i>
<i>B. globisporus</i> (eg. <i>B.</i>	<i>B. decolorationis</i>	<i>B. smithii</i>	<i>B. qingshengii</i>	<i>B. parabravis</i>
g. subsp. <i>Globisporus</i> ; or <i>B.</i>	<i>B. deserti</i>	<i>B. soli</i>	<i>B. reuszeri</i>	<i>B. paraffexus</i>
g. subsp. <i>Marinus</i>)		<i>B. solimangrovi</i>	<i>B. rhizosphaerae</i>	<i>B. pasteurii</i>
		<i>B. solisalsi</i>	<i>B. rigui</i>	<i>B. patagoniensis</i>
		<i>B. songklensis</i>	<i>B. ruris</i>	
		<i>B. sonorensis</i>	<i>B. safensis</i>	
		<i>B. sphaericus</i>	<i>B. salarius</i>	
		<i>B. sporothermodurans</i>		
		<i>B. stearothermophilus</i>		
		<i>B. stratosphericus</i>		
		<i>B. subterraneus</i>		
		<i>B. subtilis</i> (eg. <i>B.</i>		
		s. subsp. <i>Inaquosorum</i> ; or <i>B.</i>		
		s. subsp. <i>Spizizenii</i> ; or <i>B.</i>		
		s. subsp. <i>Subtilis</i>)		
<i>Caenimonas</i>	<i>Campylobacter</i>	<i>Cardiobacterium</i>	<i>Catenuloplanes</i>	<i>Curtobacterium</i>
<i>Caenimonas koreensis</i>	<i>Campylobacter coli</i>	<i>Cardiobacterium hominis</i>	<i>Catenuloplanes atrovinosus</i>	<i>Curtobacterium albidum</i>
<i>Caldalkalibacillus</i>	<i>Campylobacter concisus</i>	<i>Camimonas</i>	<i>Catenuloplanes castaneus</i>	<i>Curtobacterium citreus</i>
<i>Caldalkalibacillus uzonensis</i>	<i>Campylobacter curvus</i>	<i>Camimonas nigrificans</i>	<i>Catenuloplanes crispus</i>	
<i>Caldanaerobacter</i>	<i>Campylobacter fetus</i>	<i>Carnobacterium</i>	<i>Catenuloplanes indicus</i>	
<i>Caldanaerobacter</i>	<i>Campylobacter gracilis</i>	<i>Carnobacterium alterfunditum</i>	<i>Catenuloplanes japonicus</i>	
<i>subterraneus</i>				
<i>Caldanaerobius</i>	<i>Campylobacter helveticus</i>	<i>Carnobacterium divergens</i>	<i>Catenuloplanes nepalensis</i>	
<i>Caldanaerobius fijiensis</i>	<i>Campylobacter hominis</i>	<i>Carnobacterium funditum</i>	<i>Catenuloplanes niger</i>	
<i>Caldanaerobius</i>	<i>Campylobacter hyointestinalis</i>	<i>Carnobacterium gallinarum</i>	<i>Chryseobacterium</i>	
<i>polysaccharolyticus</i>	<i>Campylobacter jejuni</i>	<i>Carnobacterium</i>	<i>Chryseobacterium</i>	
<i>Caldanaerobius zeae</i>	<i>Campylobacter lari</i>	<i>maltaromaticum</i>	<i>balustinum</i>	
<i>Caldanaerovirga</i>	<i>Campylobacter mucosalis</i>	<i>Carnobacterium mobile</i>	<i>Citrobacter</i>	
<i>Caldanaerovirga acetigignens</i>	<i>Campylobacter rectus</i>	<i>Carnobacterium viridans</i>	<i>C. amalonaticus</i>	
<i>Caldicellulosiruptor</i>	<i>Campylobacter showae</i>	<i>Caryophanon</i>	<i>C. braakii</i>	
<i>Caldicellulosiruptor bescii</i>	<i>Campylobacter sputorum</i>	<i>Caryophanon latum</i>	<i>C. diversus</i>	
<i>Caldicellulosiruptor</i>	<i>Campylobacter upsaliensis</i>	<i>Caryophanon tenue</i>	<i>C. farmeri</i>	
<i>kristjanssonii</i>				
<i>Caldicellulosiruptor</i>	<i>Capnocytophaga</i>	<i>Catellatospora</i>	<i>C. freundii</i>	
<i>owensensis</i>	<i>Capnocytophaga canimorsus</i>	<i>Catellatospora citrea</i>	<i>C. gilleanii</i>	
	<i>Capnocytophaga cynodegmi</i>	<i>Catellatospora</i>	<i>C. koseri</i>	
	<i>Capnocytophaga gingivalis</i>	<i>methionotrophica</i>	<i>C. murlinae</i>	
	<i>Capnocytophaga granulosa</i>	<i>Catenococcus</i>	<i>C. pasteurii</i> ^[1]	
	<i>Capnocytophaga haemolytica</i>	<i>Catenococcus thiocycli</i>	<i>C. rodentium</i>	
	<i>Capnocytophaga ochracea</i>		<i>C. sedlakii</i>	
	<i>Capnocytophaga sputigena</i>		<i>C. werkmanii</i>	
			<i>C. youngae</i>	
			<i>Clostridium</i>	
			(see below)	
			<i>Coccochloris</i>	
			<i>Coccochloris elabens</i>	
			<i>Corynebacterium</i>	
			<i>Corynebacterium flavescens</i>	
			<i>Corynebacterium variabile</i>	
<i>Clostridium</i>				
<i>Clostridium absonum</i> , <i>Clostridium aceticum</i> , <i>Clostridium acetireducens</i> , <i>Clostridium acetobutylicum</i> , <i>Clostridium acidisoli</i> , <i>Clostridium aciditolerans</i> ,				
<i>Clostridium acidurici</i> , <i>Clostridium aerotolerans</i> , <i>Clostridium aestuarii</i> , <i>Clostridium akagii</i> , <i>Clostridium aldenense</i> , <i>Clostridium aldrichii</i> , <i>Clostridium</i>				
<i>algidicarni</i> , <i>Clostridium algidixylanolyticum</i> , <i>Clostridium algifaecis</i> , <i>Clostridium algoriphilum</i> , <i>Clostridium alkalicellulosi</i> , <i>Clostridium aminophilum</i> ,				
<i>Clostridium aminovalericum</i> , <i>Clostridium amygdalinum</i> , <i>Clostridium amylolyticum</i> , <i>Clostridium arbusti</i> , <i>Clostridium arcticum</i> , <i>Clostridium</i>				
<i>argenfinense</i> , <i>Clostridium asparagiforme</i> , <i>Clostridium aurantibutyricum</i> , <i>Clostridium autoethanogenum</i> , <i>Clostridium baratii</i> , <i>Clostridium barkeri</i> ,				
<i>Clostridium bartlettii</i> , <i>Clostridium bejerinckii</i> , <i>Clostridium bifermentans</i> , <i>Clostridium bolteae</i> , <i>Clostridium bornimense</i> , <i>Clostridium botulinum</i> ,				
<i>Clostridium bowmanii</i> , <i>Clostridium bryantii</i> , <i>Clostridium butyricum</i> , <i>Clostridium cadaveris</i> , <i>Clostridium caenicola</i> , <i>Clostridium caminitermale</i> ,				
<i>Clostridium carboxidivorans</i> , <i>Clostridium carnis</i> , <i>Clostridium cavendishii</i> , <i>Clostridium celatum</i> , <i>Clostridium celerecrescens</i> , <i>Clostridium</i>				
<i>cellobioperum</i> , <i>Clostridium cellulofementans</i> , <i>Clostridium cellulolyticum</i> , <i>Clostridium cellulosi</i> , <i>Clostridium cellulovorans</i> , <i>Clostridium</i>				
<i>chartatabidum</i> , <i>Clostridium chauvoei</i> , <i>Clostridium chromiireducens</i> , <i>Clostridium citroniae</i> , <i>Clostridium clariflavum</i> , <i>Clostridium clostridioforme</i> ,				
<i>Clostridium coccoides</i> , <i>Clostridium cochlearium</i> , <i>Clostridium colletant</i> , <i>Clostridium colicanis</i> , <i>Clostridium colinum</i> , <i>Clostridium collagenovorans</i> ,				
<i>Clostridium cylindrosporum</i> , <i>Clostridium difficile</i> , <i>Clostridium diolis</i> , <i>Clostridium disporicum</i> , <i>Clostridium drakei</i> , <i>Clostridium durum</i> , <i>Clostridium</i>				
<i>estertheticum</i> , <i>Clostridium estertheticum estertheticum</i> , <i>Clostridium estertheticum laramiense</i> , <i>Clostridium fallax</i> , <i>Clostridium felsineum</i> ,				

TABLE 1-continued

Example Bacteria
Optionally, the bacteria are selected from this Table.

<i>Clostridium fervidum, Clostridium fimetarium, Clostridium formicaceticum, Clostridium frigidicarnis, Clostridium frigris, Clostridium ganghwense, Clostridium gasigenes, Clostridium ghonii, Clostridium glycolicum, Clostridium glycyrrhizinilyticum, Clostridium grantii, Clostridium haemolyticum, Clostridium halophilum, Clostridium hastiforme, Clostridium hathewayi, Clostridium herbivorans, Clostridium hiranonis, Clostridium histolyticum, Clostridium homopropionicum, Clostridium huakuii, Clostridium hungatei, Clostridium hydrogeniformans, Clostridium hydroxybenzoicum, Clostridium hylemonae, Clostridium jejuense, Clostridium indolis, Clostridium innocuum, Clostridium intestinale, Clostridium irregulare, Clostridium isatidis, Clostridium josui, Clostridium kluyveri, Clostridium lactatifermentans, Clostridium lacusfryxellense, Clostridium laramiense, Clostridium lavalense, Clostridium lentocellum, Clostridium lentoputrescens, Clostridium leptum, Clostridium limosum, Clostridium litorale, Clostridium lituseburens, Clostridium ljungdahlii, Clostridium lortetii, Clostridium lundense, Clostridium magnum, Clostridium malenominatum, Clostridium manganotii, Clostridium mayombeii, Clostridium methoxybenzovorans, Clostridium methylpentosum, Clostridium neopropionicum, Clostridium nexile, Clostridium nitrophenolicum, Clostridium oceanicum, Clostridium orbiscindens, Clostridium oroticum, Clostridium oxalicum, Clostridium papyrosolvans, Clostridium paradoxum, Clostridium paraperfringens (Alias: C. welchii), Clostridium paraputrificum, Clostridium pascui, Clostridium pasteurianum, Clostridium peptidivorans, Clostridium perenne, Clostridium perfringens, Clostridium pfennigii, Clostridium phytofermentans, Clostridium piliforme, Clostridium polysaccharolyticum, Clostridium populeti, Clostridium propionicum, Clostridium proteoclasticum, Clostridium proteolyticum, Clostridium psychrophilum, Clostridium puniceum, Clostridium purinilyticum, Clostridium putrefaciens, Clostridium putrificum, Clostridium quercicolum, Clostridium quinii, Clostridium ramosum, Clostridium rectum, Clostridium roseum, Clostridium saccharobutylicum, Clostridium saccharogumia, Clostridium saccharolyticum, Clostridium saccharoperbutylacetonicum, Clostridium sardiniense, Clostridium sartagoforme, Clostridium scatologenes, Clostridium schirmacherense, Clostridium scindens, Clostridium septicum, Clostridium sordellii, Clostridium sphenoides, Clostridium spiroforme, Clostridium sporogenes, Clostridium sporosphaeroides, Clostridium stercorarium, Clostridium stercorarium leptospartum, Clostridium stercorarium stercorarium, Clostridium stercorarium thermolacticum, Clostridium sticklandii, Clostridium straminisolvans, Clostridium subterminale, Clostridium sufflavum, Clostridium sulfidigenes, Clostridium symbiosum, Clostridium tagluense, Clostridium tepidiprofundii, Clostridium termitidis, Clostridium tertium, Clostridium tetani, Clostridium tetanomorphum, Clostridium thermaceticum, Clostridium thermautotrophicum, Clostridium thermoalcaliphilum, Clostridium thermobutyricum, Clostridium thermoceillum, Clostridium thermocopriae, Clostridium thermohydrosulfuricum, Clostridium thermolacticum, Clostridium thermopalmarium, Clostridium thermopapyrolyticum, Clostridium thermosaccharolyticum, Clostridium thermosuccinogenes, Clostridium thermosulfurigenes, Clostridium thiosulfatireducens, Clostridium tyrobutyricum, Clostridium uliginosum, Clostridium ultunense, Clostridium villosum, Clostridium vincentii, Clostridium viride, Clostridium xylanolyticum, Clostridium xylanovorans</i>						
<i>Dactyloporangium</i>	<i>Deinococcus</i>	<i>Delftia</i>	<i>Echinicola</i>			
<i>Dactyloporangium aurantiacum</i>	<i>Deinococcus aerius</i>	<i>Delftia acidovorans</i>	<i>Echinicola pacifica</i>			
<i>Dactyloporangium fulvum</i>	<i>Deinococcus apachensis</i>	<i>Desulfovibrio</i>	<i>Echinicola vietnamensis</i>			
<i>Dactyloporangium matsuzakiense</i>	<i>Deinococcus aquaticus</i>	<i>Desulfovibrio desulfuricans</i>				
<i>Dactyloporangium roseum</i>	<i>Deinococcus aquatilis</i>	<i>Diplococcus</i>				
<i>Dactyloporangium thailandense</i>	<i>Deinococcus caeni</i>	<i>Diplococcus pneumoniae</i>				
<i>Dactyloporangium vinaceum</i>	<i>Deinococcus radiodurans</i>					
	<i>Deinococcus radiophilus</i>					
<i>Enterobacter</i>	<i>Enterobacter kobei</i>	<i>Faecalibacterium</i>	<i>Flavobacterium</i>			
<i>E. aerogenes</i>	<i>E. ludwigii</i>	<i>Faecalibacterium prausnitzii</i>	<i>Flavobacterium antarcticum</i>			
<i>E. amnigenus</i>	<i>E. mori</i>	<i>Fangia</i>	<i>Flavobacterium aquatile</i>			
<i>E. agglomerans</i>	<i>E. nimipressuralis</i>	<i>Fangia hongkongensis</i>	<i>Flavobacterium aquidurensis</i>			
<i>E. arachidis</i>	<i>E. oryzae</i>	<i>Fastidiosipila</i>	<i>Flavobacterium balustinum</i>			
<i>E. asburiae</i>	<i>E. pulveris</i>	<i>Fastidiosipila sanguinis</i>	<i>Flavobacterium croceum</i>			
<i>E. cancerogenus</i>	<i>E. pyrinus</i>	<i>Fusobacterium</i>	<i>Flavobacterium cucumis</i>			
<i>E. cloacae</i>	<i>E. radiccinitans</i>	<i>Fusobacterium nucleatum</i>	<i>Flavobacterium daejeonense</i>			
<i>E. cowanii</i>	<i>E. taylorae</i>		<i>Flavobacterium defluvii</i>			
<i>E. dissolvans</i>	<i>E. turicensis</i>		<i>Flavobacterium degerlachei</i>			
<i>E. gergoviae</i>	<i>E. sakazakii</i>	<i>Enterobacter soli</i>	<i>Flavobacterium denitrificans</i>			
<i>E. helveticus</i>	<i>Enterococcus</i>		<i>Flavobacterium filum</i>			
<i>E. hormaechei</i>	<i>Enterococcus durans</i>		<i>Flavobacterium flevense</i>			
<i>E. intermedius</i>	<i>Enterococcus faecalis</i>		<i>Flavobacterium frigidarium</i>			
	<i>Enterococcus faecium</i>		<i>Flavobacterium mizutaii</i>			
	<i>Erwinia</i>		<i>Flavobacterium okeanokoites</i>			
	<i>Erwinia hapontici</i>					
	<i>Escherichia</i>					
	<i>Escherichia coli</i>					
<i>Gaebulibacter</i>	<i>Haemophilus</i>	<i>Ideonella</i>	<i>Janibacter</i>			
<i>Gaebulibacter saemankumensis</i>	<i>Haemophilus aegyptius</i>	<i>Ideonella azotifigens</i>	<i>Janibacter anophelis</i>			
<i>Gallibacterium</i>	<i>Haemophilus aphrophilus</i>	<i>Idiomarina</i>	<i>Janibacter corallicola</i>			
<i>Gallibacterium anatis</i>	<i>Haemophilus felis</i>	<i>Idiomarina abyssalis</i>	<i>Janibacter limosus</i>			
<i>Gallicola</i>	<i>Haemophilus gallinarum</i>	<i>Idiomarina baltica</i>	<i>Janibacter melonis</i>			
<i>Gallicola barnesae</i>	<i>Haemophilus haemolyticus</i>	<i>Idiomarina fontislapidosi</i>	<i>Janibacter terrae</i>			
<i>Garciella</i>	<i>Haemophilus influenzae</i>	<i>Idiomarina loihensis</i>	<i>Jamaschia</i>			
<i>Garciella nitratireducens</i>	<i>Haemophilus paracuniculus</i>	<i>Idiomarina ramblicola</i>	<i>Jamaschia cystaugens</i>			
<i>Geobacillus</i>	<i>Haemophilus parahaemolyticus</i>	<i>Idiomarina seosinensis</i>	<i>Jamaschia helgolandensis</i>			
	<i>Haemophilus parainfluenzae</i>	<i>Idiomarina zobellii</i>	<i>Jamaschia pohangensis</i>			
<i>Geobacillus thermoglucosidasius</i>						
<i>Geobacillus stearothermophilus</i>	<i>Haemophilus</i>	<i>Ignatzschineria</i>	<i>Jamaschia rubra</i>			
<i>Geobacter</i>	<i>paraphrohaemolyticus</i>	<i>Ignatzschineria larvae</i>	<i>Janthinobacterium</i>			
<i>Geobacter bemidjensis</i>	<i>Haemophilus parasuis</i>	<i>Ignavigranum</i>	<i>Janthinobacterium agaricidamnosum</i>			
<i>Geobacter bremensis</i>	<i>Haemophilus pittmaniae</i>	<i>Ignavigranum ruoffiae</i>	<i>Janthinobacterium lividum</i>			
<i>Geobacter chapellei</i>	<i>Hafnia</i>	<i>Ilumatobacter</i>	<i>Jejuia</i>			
<i>Geobacter grbiciae</i>	<i>Hafnia alvei</i>	<i>Ilumatobacter fluminis</i>	<i>Jejuia pallidilutea</i>			
<i>Geobacter hydrogenophilus</i>	<i>Hahella</i>	<i>Ilyobacter</i>				

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>Geobacter lovleyi</i>	<i>Hahella ganghwensis</i>	<i>Ilyobacter delafieldii</i>	<i>Jeotgalibacillus</i>	
<i>Geobacter metallireducens</i>	<i>Halalkalibacillus</i>	<i>Ilyobacter insuetus</i>	<i>Jeotgalibacillus</i>	
<i>Geobacter pelophilus</i>	<i>Halalkalibacillus halophilus</i>	<i>Ilyobacter polytropus</i>	<i>alimentarius</i>	
<i>Geobacter pickeringii</i>	<i>Helicobacter</i>	<i>Ilyobacter tartaricus</i>	<i>Jeotgalicoccus</i>	
<i>Geobacter sulfurreducens</i>	<i>Helicobacter pylori</i>		<i>Jeotgalicoccus halotolerans</i>	
<i>Geodermatophilus</i>				
<i>Geodermatophilus obscurus</i>				
<i>Gluconacetobacter</i>				
<i>Gluconacetobacter xylinus</i>				
<i>Gordonia</i>				
<i>Gordonia rubripertincta</i>				
<i>Kaistia</i>	<i>Labedella</i>	<i>Listeria ivanovii</i>	<i>Micrococcus</i>	<i>Nesterenkonia</i>
<i>Kaistia adipata</i>	<i>Labedella gwakjiensis</i>	<i>L. marthii</i>	<i>Micrococcus luteus</i>	<i>Nesterenkonia holobia</i>
<i>Kaistia soli</i>	<i>Labrenzia</i>	<i>L. monocytogenes</i>	<i>Micrococcus lylae</i>	<i>Nocardia</i>
<i>Kangiella</i>	<i>Labrenzia aggregata</i>	<i>L. newyorkensis</i>	<i>Moraxella</i>	<i>Nocardia argentinensis</i>
<i>Kangiella aquimarina</i>	<i>Labrenzia alba</i>	<i>L. riparia</i>	<i>Moraxella bovis</i>	<i>Nocardia corallina</i>
<i>Kangiella koreensis</i>	<i>Labrenzia alexandrii</i>	<i>L. rocourtiae</i>	<i>Moraxella nonliquefaciens</i>	<i>Nocardia</i>
<i>Kersteria</i>	<i>Labrenzia marina</i>	<i>L. seeligeri</i>	<i>Moraxella osloensis</i>	<i>otitidiscaeviarum</i>
<i>Kersteria gyiorum</i>	<i>Labrys</i>	<i>L. weihenstephanensis</i>	<i>Nakamurella</i>	
<i>Kiloniella</i>	<i>Labrys methylaminiphilus</i>	<i>L. welschimeri</i>	<i>Nakamurella multipartita</i>	
<i>Kiloniella laminariae</i>	<i>Labrys miyagiensis</i>	<i>Listonella</i>	<i>Nannocystis</i>	
<i>Klebsiella</i>	<i>Labrys monachus</i>	<i>Listonella anguillarum</i>	<i>Nannocystis pusilla</i>	
<i>K. granulomatis</i>	<i>Labrys okinawensis</i>	<i>Macrococcus</i>	<i>Natranaerobius</i>	
<i>K. oxytoca</i>	<i>Labrys portucalensis</i>	<i>Macrococcus bovicus</i>	<i>Natranaerobius</i>	
<i>K. pneumoniae</i>	<i>Lactobacillus</i>	<i>Marinobacter</i>	<i>thermophilus</i>	
<i>K. terrigena</i>	[see below]	<i>Marinobacter algicola</i>	<i>Natranaerobius trueperi</i>	
<i>K. varicola</i>	<i>Laceyella</i>	<i>Marinobacter bryozorum</i>	<i>Naxibacter</i>	
<i>Kluyvera</i>	<i>Laceyella putida</i>	<i>Marinobacter flavimaris</i>	<i>Naxibacter alkalitolerans</i>	
<i>Kluyvera ascorbata</i>	<i>Lechevalieria</i>	<i>Meiothermus</i>	<i>Neisseria</i>	
<i>Kocuria</i>	<i>Lechevalieria</i>	<i>Meiothermus ruber</i>	<i>Neisseria cinerea</i>	
	<i>aerocolonigenes</i>			
<i>Kocuria roasea</i>	<i>Legionella</i>	<i>Methylophilus</i>	<i>Neisseria denitrificans</i>	
<i>Kocuria varians</i>	[see below]	<i>Methylophilus methylotrophus</i>	<i>Neisseria gonorrhoeae</i>	
<i>Kurthia</i>	<i>Listeria</i>	<i>Microbacterium</i>	<i>Neisseria lactamica</i>	
<i>Kurthia zopfii</i>	<i>L. aquatica</i>	<i>Microbacterium</i>	<i>Neisseria mucosa</i>	
	<i>L. booriae</i>	<i>ammoniphilum</i>	<i>Neisseria sicca</i>	
	<i>L. cornellensis</i>	<i>Microbacterium arborescens</i>	<i>Neisseria subflava</i>	
	<i>L. fleischmannii</i>	<i>Microbacterium liquefaciens</i>	<i>Neptunomonas</i>	
	<i>L. floridensis</i>	<i>Microbacterium oxydans</i>	<i>Neptunomonas japonica</i>	
	<i>L. grandensis</i>			
	<i>L. grayi</i>			
	<i>L. imocua</i>			
<i>Lactobacillus</i>				
<i>L. acetotolerans</i>	<i>L. cateniformis</i>	<i>L. mali</i>	<i>L. parakefiri</i>	<i>L. sakei</i>
<i>L. acidifarinae</i>	<i>L. ceti</i>	<i>L. manihotivorans</i>	<i>L. paralimentarius</i>	<i>L. salivarius</i>
<i>L. acidipiscis</i>	<i>L. coleohominis</i>	<i>L. mindensis</i>	<i>L. paraplantarum</i>	<i>L. sanfranciscensis</i>
<i>L. acidophilus</i>	<i>L. collinoides</i>	<i>L. mucosae</i>	<i>L. pentosus</i>	<i>L. satsumensis</i>
<i>Lactobacillus agilis</i>	<i>L. composti</i>	<i>L. murinus</i>	<i>L. perolens</i>	<i>L. secaliphilus</i>
<i>L. algidus</i>	<i>L. concavus</i>	<i>L. nagelii</i>	<i>L. plantarum</i>	<i>L. sharpeae</i>
<i>L. alimentarius</i>	<i>L. coryniformis</i>	<i>L. namuensis</i>	<i>L. pontis</i>	<i>L. siliginis</i>
<i>L. amylolyticus</i>	<i>L. crispatus</i>	<i>L. nantensis</i>	<i>L. protectus</i>	<i>L. spicheri</i>
<i>L. amylophilus</i>	<i>L. crustorum</i>	<i>L. oligofermentans</i>	<i>L. psittaci</i>	<i>L. suebicus</i>
<i>L. amylophobicus</i>	<i>L. curvatus</i>	<i>L. oris</i>	<i>L. rennini</i>	<i>L. thailandensis</i>
<i>L. amylovorus</i>	<i>L. delbrueckii</i> subsp. <i>bulgaricus</i>	<i>L. panis</i>	<i>L. reuteri</i>	<i>L. ultunensis</i>
<i>L. animalis</i>	<i>L. delbrueckii</i> subsp. <i>delbrueckii</i>	<i>L. pantheris</i>	<i>L. rhamnosus</i>	<i>L. vaccinostercus</i>
<i>L. antri</i>	<i>L. delbrueckii</i> subsp. <i>lactis</i>	<i>L. parabrevis</i>	<i>L. rimae</i>	<i>L. vaginalis</i>
<i>L. apodemi</i>	<i>L. dextrinicus</i>	<i>L. parabuchneri</i>	<i>L. rogosae</i>	<i>L. versmoldensis</i>
<i>L. aviarius</i>	<i>L. diolivorans</i>	<i>L. paracasei</i>	<i>L. rossiae</i>	<i>L. vini</i>
<i>L. bifermantans</i>	<i>L. equi</i>	<i>L. paracollinoides</i>	<i>L. ruminis</i>	<i>L. vitulinus</i>
<i>L. brevis</i>	<i>L. equigenerosi</i>	<i>L. parafarraginis</i>	<i>L. saerimmeri</i>	<i>L. zeae</i>
<i>L. buchneri</i>	<i>L. farraginis</i>	<i>L. homiohiochii</i>	<i>L. jensenii</i>	<i>L. zymae</i>
<i>L. camelliae</i>	<i>L. farciminis</i>	<i>L. iners</i>	<i>L. johnsonii</i>	<i>L. gastricus</i>
<i>L. casei</i>	<i>L. fermentum</i>	<i>L. ingluviei</i>	<i>L. kalixensis</i>	<i>L. ghanensis</i>
<i>L. kitasatonis</i>	<i>L. fornicalis</i>	<i>L. intestinalis</i>	<i>L. kefiranofermentans</i>	<i>L. graminis</i>
<i>L. kunkei</i>	<i>L. fructivorans</i>	<i>L. fuchuiensis</i>	<i>L. kefirii</i>	<i>L. hammesii</i>
<i>L. leichmannii</i>	<i>L. frumentii</i>	<i>L. gallinarum</i>	<i>L. kimchii</i>	<i>L. hamsteri</i>
<i>L. lindneri</i>		<i>L. gasserii</i>	<i>L. helveticus</i>	<i>L. harbinensis</i>
<i>L. malefermentans</i>			<i>L. hilgardii</i>	<i>L. hayakitensis</i>
<i>Legionella</i>				
<i>Legionella adelaidensis</i>	<i>Legionella drancourtii</i>	<i>Candidatus Legionella jeonii</i>	<i>Legionella quinlivanii</i>	
<i>Legionella anisa</i>	<i>Legionella dresdenensis</i>	<i>Legionella jordanis</i>	<i>Legionella rowbothamii</i>	
<i>Legionella beliardensis</i>	<i>Legionella drozanskii</i>	<i>Legionella lansingensis</i>	<i>Legionella rubrilucens</i>	
<i>Legionella birminghamensis</i>	<i>Legionella dumoffii</i>	<i>Legionella londiniensis</i>	<i>Legionella sainthelensi</i>	
<i>Legionella bozemanae</i>	<i>Legionella erythra</i>	<i>Legionella longbeachae</i>	<i>Legionella santacrucis</i>	
<i>Legionella brunensis</i>	<i>Legionella fairfieldensis</i>	<i>Legionella lytica</i>	<i>Legionella shakespearei</i>	
<i>Legionella busanensis</i>	<i>Legionella fallonii</i>	<i>Legionella maceachernii</i>	<i>Legionella spiritensis</i>	

TABLE 1-continued

Example Bacteria			
Optionally, the bacteria are selected from this Table.			
<i>Legionella cardiaca</i>	<i>Legionella feelei</i>	<i>Legionella massiliensis</i>	<i>Legionella steelei</i>
<i>Legionella cherrii</i>	<i>Legionella geestiana</i>	<i>Legionella micdadei</i>	<i>Legionella steigerwaltii</i>
<i>Legionella cincinnatiensis</i>	<i>Legionella genomospecies</i>	<i>Legionella monrovia</i>	<i>Legionella taurinensis</i>
<i>Legionella clemsonensis</i>	<i>Legionella gormanii</i>	<i>Legionella moravica</i>	<i>Legionella tucsonensis</i>
<i>Legionella donaldsonii</i>	<i>Legionella gratiana</i>	<i>Legionella nagasakiensis</i>	<i>Legionella tunisiensis</i>
	<i>Legionella gresilensis</i>	<i>Legionella nautarum</i>	<i>Legionella wadsworthii</i>
	<i>Legionella hackeliae</i>	<i>Legionella norrlandica</i>	<i>Legionella waltersii</i>
	<i>Legionella impletisoli</i>	<i>Legionella oakridgensis</i>	<i>Legionella worsteiensis</i>
	<i>Legionella israelensis</i>	<i>Legionella parisiensis</i>	<i>Legionella yabuuchiae</i>
	<i>Legionella jamestowniensis</i>	<i>Legionella pittsburghensis</i>	
		<i>Legionella pneumophila</i>	
		<i>Legionella quateirensis</i>	
<i>Oceanibulbus</i>	<i>Paenibacillus</i>	<i>Prevotella</i>	<i>Quadriflora</i>
<i>Oceanibulbus indolifex</i>	<i>Paenibacillus thiaminolyticus</i>	<i>Prevotella albensis</i>	<i>Quadriflora granulorum</i>
<i>Oceanicaulis</i>	<i>Pantoea</i>	<i>Prevotella amnii</i>	<i>Quatrimonococcus</i>
<i>Oceanicaulis alexandrii</i>	<i>Pantoea agglomerans</i>	<i>Prevotella bergensis</i>	<i>Quatrimonococcus</i>
<i>Oceanicola</i>	<i>Paracoccus</i>	<i>Prevotella bivia</i>	<i>australiensis</i>
<i>Oceanicola batsensis</i>	<i>Paracoccus alcaliphilus</i>	<i>Prevotella brevis</i>	<i>Quinella</i>
<i>Oceanicola granulosis</i>	<i>Paucimonas</i>	<i>Prevotella bryantii</i>	<i>Quinella ovalis</i>
<i>Oceanicola nanhaiensis</i>	<i>Paucimonas lemoignei</i>	<i>Prevotella buccae</i>	<i>Ralstonia</i>
<i>Oceanimonas</i>	<i>Pectobacterium</i>	<i>Prevotella buccalis</i>	<i>Ralstonia eutropha</i>
<i>Oceanimonas baumannii</i>	<i>Pectobacterium aroidearum</i>	<i>Prevotella copri</i>	<i>Ralstonia insidiosa</i>
<i>Oceaniserpentilla</i>	<i>Pectobacterium atrosepticum</i>	<i>Prevotella dentalis</i>	<i>Ralstonia mannitolilytica</i>
<i>Oceaniserpentilla haliotis</i>	<i>Pectobacterium betavasculorum</i>	<i>Prevotella denticola</i>	<i>Ralstonia pickettii</i>
<i>Oceanisphaera</i>	<i>Pectobacterium cacticida</i>	<i>Prevotella distens</i>	<i>Ralstonia</i>
<i>Oceanisphaera donghaensis</i>	<i>Pectobacterium carnegieana</i>	<i>Prevotella histicola</i>	<i>pseudosolanacearum</i>
<i>Oceanisphaera litoralis</i>	<i>Pectobacterium carotovorum</i>	<i>Prevotella intermedia</i>	<i>Ralstonia syzygii</i>
<i>Oceanithermus</i>	<i>Pectobacterium chrysanthemi</i>	<i>Prevotella maculosa</i>	<i>Ralstonia solanacearum</i>
<i>Oceanithermus desulfurans</i>	<i>Pectobacterium cypripedii</i>	<i>Prevotella marshii</i>	<i>Ramlibacter</i>
<i>Oceanithermus profundus</i>	<i>Pectobacterium rhapontici</i>	<i>Prevotella melaninogenica</i>	<i>Ramlibacter henchirensis</i>
<i>Oceanobacillus</i>	<i>Pectobacterium wasabiae</i>	<i>Prevotella micans</i>	<i>Ramlibacter tataouinensis</i>
<i>Oceanobacillus caeni</i>	<i>Planococcus</i>	<i>Prevotella multiformis</i>	<i>Raoultella</i>
<i>Oceanospirillum</i>	<i>Planococcus citreus</i>	<i>Prevotella nigrescens</i>	<i>Raoultella ornithinolytica</i>
<i>Oceanospirillum linum</i>	<i>Planomicrobium</i>	<i>Prevotella oralis</i>	<i>Raoultella planticola</i>
	<i>Planomicrobium okeanoikoites</i>	<i>Prevotella oris</i>	<i>Raoultella terrigena</i>
	<i>Plesiomonas</i>	<i>Prevotella oulorum</i>	<i>Rathayibacter</i>
	<i>Plesiomonas shigelloides</i>	<i>Prevotella pallens</i>	<i>Rathayibacter caricis</i>
	<i>Proteus</i>	<i>Prevotella salivae</i>	<i>Rathayibacter festucae</i>
	<i>Proteus vulgaris</i>	<i>Prevotella stercorea</i>	<i>Rathayibacter iranicus</i>
		<i>Prevotella tammarae</i>	<i>Rathayibacter rathayi</i>
		<i>Prevotella timonensis</i>	<i>Rathayibacter toxicus</i>
		<i>Prevotella veroralis</i>	<i>Rathayibacter tritici</i>
		<i>Providencia</i>	<i>Rhodobacter</i>
		<i>Providencia stuartii</i>	<i>Rhodobacter sphaeroides</i>
		<i>Pseudomonas</i>	<i>Ruegeria</i>
		<i>Pseudomonas aeruginosa</i>	<i>Ruegeria gelatinovorans</i>
		<i>Pseudomonas alcaligenes</i>	
		<i>Pseudomonas anguilliseptica</i>	
		<i>Pseudomonas fluorescens</i>	
		<i>Pseudoalteromonas haloplanktis</i>	
		<i>Pseudomonas mendocina</i>	
		<i>Pseudomonas pseudoalcaligenes</i>	
		<i>Pseudomonas putida</i>	
		<i>Pseudomonas tutzeri</i>	
		<i>Pseudomonas syringae</i>	
		<i>Psychrobacter</i>	
		<i>Psychrobacter faecalis</i>	
		<i>Psychrobacter phenylpyruvicus</i>	
		<i>Sanguibacter</i>	
<i>Saccharococcus</i>	<i>Sagittula</i>	<i>Sanguibacter keddietii</i>	<i>Stenotrophomonas maltophilia</i>
<i>Saccharococcus thermophilus</i>	<i>Sagittula stellata</i>	<i>Sanguibacter suarezi</i>	<i>Streptococcus</i>
<i>Saccharomonospora</i>	<i>Salagentibacter</i>	<i>Saprospira</i>	[also see below]
<i>Saccharomonospora azurea</i>	<i>Salagentibacter salegens</i>	<i>Saprospira grandis</i>	<i>Streptomyces</i>
<i>Saccharomonospora cyanea</i>	<i>Salimicrobium</i>	<i>Sarcina</i>	<i>Streptomyces cescaepitosus</i>
<i>Saccharomonospora viridis</i>	<i>Salimicrobium album</i>	<i>Sarcina maxima</i>	<i>Streptomyces cesdiastaticus</i>
<i>Saccharophagus</i>	<i>Salinibacter</i>	<i>Sarcina ventriculi</i>	<i>Streptomyces achromogenes</i>
<i>Saccharophagus degradans</i>	<i>Salinibacter ruber</i>	<i>Sebaldella</i>	<i>Streptomyces cesalbus</i>
<i>Saccharopolyspora</i>	<i>Salinicoccus</i>	<i>Sebaldella termitidis</i>	<i>Streptomyces cessexfoliatus</i>
<i>Saccharopolyspora erythraea</i>	<i>Salinicoccus alkaliphilus</i>	<i>Serratia</i>	<i>Streptomyces fimbriatus</i>
<i>Saccharopolyspora gregortii</i>	<i>Salinicoccus hispanicus</i>	<i>Serratia fonticola</i>	<i>Streptomyces fradiae</i>
<i>Saccharopolyspora hirsuta</i>	<i>Salinicoccus roseus</i>	<i>Serratia marcescens</i>	
<i>Saccharopolyspora hordei</i>	<i>Salinispora</i>	<i>Sphaerotilus</i>	
<i>Saccharopolyspora rectivirgula</i>	<i>Salinispora arenicola</i>		
			<i>Tatlockia</i>
			<i>Tatlockia maceachernii</i>
			<i>Tatlockia micdadei</i>
			<i>Tenacibaculum amylolyticum</i>
			<i>Tenacibaculum discolor</i>
			<i>Tenacibaculum lutimaris</i>
			<i>Tenacibaculum gallaicum</i>
			<i>Tenacibaculum mesophilum</i>
			<i>Tenacibaculum</i>

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>Saccharopolyspora spinosa</i>	<i>Salinispora tropica</i>	<i>Sphaerotilus natans</i>	<i>Streptomyces fulvisimus</i>	<i>skagerrakense</i>
<i>Saccharopolyspora taberi</i>	<i>Salinivibrio</i>	<i>Sphingobacterium</i>	<i>Streptomyces griseoruber</i>	<i>Tepidanaerobacter</i>
<i>Saccharothrix</i>	<i>Salinivibrio costicola</i>	<i>Sphingobacterium multivorum</i>	<i>Streptomyces griseus</i>	<i>Tepidanaerobacter</i>
<i>Saccharothrix australiensis</i>	<i>Salmonella</i>	<i>Staphylococcus</i>	<i>Streptomyces lavendulae</i>	<i>syntrophicus</i>
<i>Saccharothrix coeruleofusca</i>	<i>Salmonella bongori</i>	[see below]	<i>Streptomyces</i>	<i>Tepidibacter</i>
<i>Saccharothrix espanaensis</i>	<i>Salmonella enterica</i>		<i>phaeochromogenes</i>	<i>Tepidibacter</i>
<i>Saccharothrix longispora</i>	<i>Salmonella subterranea</i>		<i>Streptomyces</i>	<i>formicigenes</i>
<i>Saccharothrix mutabilis</i>	<i>Salmonella typhi</i>		<i>thermodiastaticus</i>	<i>Tepidibacter thalassicus</i>
<i>Saccharothrix syringae</i>			<i>Streptomyces tubercidicus</i>	<i>Thermus</i>
<i>Saccharothrix tangerinus</i>				<i>Thermus aquaticus</i>
<i>Saccharothrix texasensis</i>				<i>Thermus filiformis</i>
				<i>Thermus thermophilus</i>
<i>Staphylococcus</i>				
<i>S. arlettae</i>	<i>S. equorum</i>	<i>S. microti</i>	<i>S. schleiferi</i>	
<i>S. agnetis</i>	<i>S. felis</i>	<i>S. muscae</i>	<i>S. sciuri</i>	
<i>S. aureus</i>	<i>S. fleurettii</i>	<i>S. nepalensis</i>	<i>S. simiae</i>	
<i>S. auricularis</i>	<i>S. gallinarum</i>	<i>S. pasteurii</i>	<i>S. simulans</i>	
<i>S. capitis</i>	<i>S. haemolyticus</i>	<i>S. petrasii</i>	<i>S. stepanovicii</i>	
<i>S. caprae</i>	<i>S. hominis</i>	<i>S. pettenkoferi</i>	<i>S. succinus</i>	
<i>S. carnosus</i>	<i>S. hyicus</i>	<i>S. piscifermentans</i>	<i>S. vitulinus</i>	
<i>S. caseolyticus</i>	<i>S. intermedius</i>	<i>S. pseudintermedius</i>	<i>S. warneri</i>	
<i>S. chromogenes</i>	<i>S. kloosii</i>	<i>S. pseudolugdunensis</i>	<i>S. xylosus</i>	
<i>S. cohnii</i>	<i>S. leei</i>	<i>S. pulvereri</i>		
<i>S. condimenti</i>	<i>S. lentus</i>	<i>S. rostri</i>		
<i>S. delphini</i>	<i>S. lugdunensis</i>	<i>S. saccharolyticus</i>		
<i>S. devriesei</i>	<i>S. lutrae</i>	<i>S. saprophyticus</i>		
<i>S. epidermidis</i>	<i>S. lyticans</i>			
	<i>S. massiliensis</i>			
<i>Streptococcus</i>				
<i>Streptococcus agalactiae</i>	<i>Streptococcus infantarius</i>	<i>Streptococcus orisratti</i>	<i>Streptococcus thermophilus</i>	
<i>Streptococcus anginosus</i>	<i>Streptococcus iniae</i>	<i>Streptococcus parasanguinis</i>	<i>Streptococcus sanguinis</i>	
<i>Streptococcus bovis</i>	<i>Streptococcus intermedius</i>	<i>Streptococcus peroris</i>	<i>Streptococcus sobrinus</i>	
<i>Streptococcus canis</i>	<i>Streptococcus lactarius</i>	<i>Streptococcus pneumoniae</i>	<i>Streptococcus suis</i>	
<i>Streptococcus constellatus</i>	<i>Streptococcus milleri</i>	<i>Streptococcus</i>	<i>Streptococcus uberis</i>	
<i>Streptococcus downei</i>	<i>Streptococcus mitis</i>	<i>pseudopneumoniae</i>	<i>Streptococcus vestibularis</i>	
<i>Streptococcus dysgalactiae</i>	<i>Streptococcus mutans</i>	<i>Streptococcus pyogenes</i>	<i>Streptococcus viridans</i>	
<i>Streptococcus equines</i>	<i>Streptococcus oralis</i>	<i>Streptococcus rattii</i>	<i>Streptococcus</i>	
<i>Streptococcus faecalis</i>	<i>Streptococcus tigurinus</i>	<i>Streptococcus salivariu</i>	<i>zooepidemicus</i>	
<i>Streptococcus ferus</i>				
<i>Uliginosibacterium</i>	<i>Vagococcus</i>	<i>Vibrio</i>	<i>Virgibacillus</i>	<i>Xanthobacter</i>
<i>Uliginosibacterium</i>	<i>Vagococcus carniphilus</i>	<i>Vibrio aerogenes</i>	<i>Virgibacillus</i>	<i>Xanthobacter agilis</i>
<i>gangwonense</i>				
<i>Ulvibacter</i>	<i>Vagococcus elongatus</i>	<i>Vibrio aestuarianus</i>	<i>halodenitrificans</i>	<i>Xanthobacter</i>
<i>Ulvibacter litoralis</i>	<i>Vagococcus fessus</i>	<i>Vibrio albensis</i>	<i>Virgibacillus</i>	<i>aminoxidans</i>
<i>Umezawaea</i>	<i>Vagococcus fluvialis</i>	<i>Vibrio alginolyticus</i>	<i>pantothenicus</i>	<i>Xanthobacter</i>
<i>Umezawaea tangerina</i>	<i>Vagococcus lutrae</i>	<i>Vibrio campbellii</i>	<i>Weissella</i>	<i>autotrophicus</i>
<i>Undibacterium</i>	<i>Vagococcus salmoninarum</i>	<i>Vibrio cholerae</i>	<i>Weissella cibaria</i>	<i>Xanthobacter flavus</i>
<i>Undibacterium pigrum</i>	<i>Variovorax</i>	<i>Vibrio cincinnatiensis</i>	<i>Weissella confusa</i>	<i>Xanthobacter tagetidis</i>
<i>Ureaplasma</i>	<i>Variovorax boronicumulans</i>	<i>Vibrio corallilyticus</i>	<i>Weissella halotolerans</i>	<i>Xanthobacter viscosus</i>
<i>Ureaplasma urealyticum</i>	<i>Variovorax dokdonensis</i>	<i>Vibrio cyclitrophicus</i>	<i>Weissella hellenica</i>	<i>Xanthomonas</i>
<i>Ureibacillus</i>	<i>Variovorax paradoxus</i>	<i>Vibrio diazotrophicus</i>	<i>Weissella kandleri</i>	<i>Xanthomonas</i>
<i>Ureibacillus composti</i>	<i>Variovorax soli</i>	<i>Vibrio fluvialis</i>	<i>Weissella koreensis</i>	<i>albilineans</i>
<i>Ureibacillus suwonensis</i>	<i>Veillonella</i>	<i>Vibrio furnissii</i>	<i>Weissella minor</i>	<i>Xanthomonas alfalfae</i>
<i>Ureibacillus terrenus</i>	<i>Veillonella atypica</i>	<i>Vibrio gazogenes</i>	<i>Weissella</i>	<i>Xanthomonas</i>
<i>Ureibacillus thermophilus</i>	<i>Veillonella caviae</i>	<i>Vibrio haliotocoli</i>	<i>paramesenteroides</i>	<i>arboricola</i>
<i>Ureibacillus</i>	<i>Veillonella criceti</i>	<i>Vibrio harveyi</i>	<i>Weissella soli</i>	<i>Xanthomonas</i>
<i>thermosphaericus</i>	<i>Veillonella dispar</i>	<i>Vibrio ichthyenteri</i>	<i>Weissella thailandensis</i>	<i>axonopodis</i>
	<i>Veillonella montpellierensis</i>	<i>Vibrio mediterranei</i>	<i>Weissella viridescens</i>	<i>Xanthomonas</i>
	<i>Veillonella parvula</i>	<i>Vibrio metschnikovii</i>	<i>Williamsia</i>	<i>campestris</i>
	<i>Veillonella ratti</i>	<i>Vibrio mytili</i>	<i>Williamsia marianensis</i>	<i>Xanthomonas citri</i>
	<i>Veillonella rodentium</i>	<i>Vibrio natriegens</i>	<i>Williamsia maris</i>	<i>Xanthomonas codiaei</i>
	<i>Venenivibrio</i>	<i>Vibrio navarrensensis</i>	<i>Williamsia serinedens</i>	<i>Xanthomonas</i>
	<i>Venenivibrio stagnispumantis</i>	<i>Vibrio nereis</i>	<i>Winogradskyella</i>	<i>cucurbitae</i>
	<i>Verminephrobacter</i>	<i>Vibrio nigripulchritudo</i>	<i>Winogradskyella</i>	<i>Xanthomonas</i>
	<i>Verminephrobacter eiseniae</i>	<i>Vibrio ordalii</i>	<i>thalassocola</i>	<i>euvasicatoria</i>
	<i>Verrucomicrobium</i>	<i>Vibrio orientalis</i>	<i>Wolbachia</i>	<i>Xanthomonas fragariae</i>
	<i>Verrucomicrobium spinosum</i>	<i>Vibrio parahaemolyticus</i>	<i>Wolbachia persica</i>	<i>Xanthomonas fuscans</i>
		<i>Vibrio pectenicida</i>	<i>Wolinella</i>	<i>Xanthomonas gardneri</i>
		<i>Vibrio penaeicida</i>	<i>Wolinella succinogenes</i>	<i>Xanthomonas hortorum</i>
		<i>Vibrio proteolyticus</i>	<i>Zobellia</i>	<i>Xanthomonas hyacinthi</i>
		<i>Vibrio shilonii</i>	<i>Zobellia galactamivorans</i>	<i>Xanthomonas perforans</i>
		<i>Vibrio splendidus</i>	<i>Zobellia uliginosa</i>	<i>Xanthomonas phaseoli</i>
		<i>Vibrio tubiashii</i>	<i>Zoogloea</i>	<i>Xanthomonas pisi</i>
		<i>Vibrio vulnificus</i>	<i>Zoogloea ramigera</i>	<i>Xanthomonas populi</i>
			<i>Zoogloea resiniphila</i>	<i>Xanthomonas theicola</i>
				<i>Xanthomonas</i>
				<i>translucens</i>

TABLE 1-continued

Example Bacteria
Optionally, the bacteria are selected from this Table.

<i>Xenophilus</i>	<i>Yangia</i>	<i>Yersinia mollaretii</i>	<i>Zooshikella</i>	<i>Xanthomonas vesicatoria</i>
<i>Xenophilus azovorans</i>	<i>Yangia pacifica</i>	<i>Yersinia philomiragia</i>	<i>Zooshikella ganghwensis</i>	<i>Xylella</i>
<i>Xenorhabdus</i>	<i>Yaniella</i>	<i>Yersinia pestis</i>	<i>Zunongwangia</i>	<i>Xylella fastidiosa</i>
<i>Xenorhabdus beddingii</i>	<i>Yaniella flava</i>	<i>Yersinia pseudotuberculosis</i>	<i>Zunongwangia profunda</i>	<i>Xylophilus</i>
<i>Xenorhabdus bovienii</i>	<i>Yaniella halotolerans</i>	<i>Yersinia rohdei</i>	<i>Zymobacter</i>	<i>Xylophilus ampelinus</i>
<i>Xenorhabdus cabanillasii</i>	<i>Yeosuana</i>	<i>Yersinia ruckeri</i>	<i>Zymobacter palmae</i>	<i>Zobellella</i>
<i>Xenorhabdus doucetiae</i>	<i>Yeosuana aromativorans</i>	<i>Yokenella</i>	<i>Zymomonas</i>	<i>Zobellella denitrificans</i>
<i>Xenorhabdus griffinae</i>	<i>Yersinia</i>	<i>Yokenella regensburgei</i>	<i>Zymomonas mobilis</i>	<i>Zobellella taiwanensis</i>
<i>Xenorhabdus hominickii</i>	<i>Yersinia aldovae</i>	<i>Yonghaparkia</i>	<i>Zymophilus</i>	<i>Zeaxanthinibacter</i>
<i>Xenorhabdus koppenhoferi</i>	<i>Yersinia bercovieri</i>	<i>Yonghaparkia alkaliphila</i>	<i>Zymophilus paucivorans</i>	<i>enoshimensis</i>
<i>Xenorhabdus nematophila</i>	<i>Yersinia enterocolitica</i>	<i>Zavarzinia</i>	<i>Zymophilus raffinosivorans</i>	<i>Zihengliuella</i>
<i>Xenorhabdus poinarii</i>	<i>Yersinia entomophaga</i>	<i>Zavarzinia compransoris</i>		<i>halotolerans</i>
<i>Xylanibacter</i>	<i>Yersinia frederiksenii</i>			<i>Xylanibacterium</i>
<i>Xylanibacter oryzae</i>	<i>Yersinia intermedia</i>			<i>ulmi</i>
	<i>Yersinia kristensenii</i>			

TABLE 2

Systemic inflammatory response syndrome	
Finding	Value
Temperature	<36° C. (96.8° F.) or >38° C. (100.4° F.)
Heart rate	>90/min
Respiratory rate	>20/min or PaCO ₂ <32 mmHg (4.3 kPa)
WBC	<4 × 10 ⁹ /L (<4000/mm ³), >12 × 10 ⁹ /L (>12,000/mm ³), or 10% bands

TABLE 3

Sequences of genomic targets to selectively CRISPR-kill *E. coli* strains. The 3-bp PAM are shown in bold type (ie, tgg, agg and cgg). Expected SpCas9-gRNA cleavage sites are indicated by vertical arrows (3 bp upstream of the PAM).

Target strain	Target gene	Genomic target sequence
<i>E. coli</i> (EHEC) ATCC43888	23S ribosomal RNA	taggtgaagtcctcgcgga tcgg ↓ (SEQ ID NO: 1)

TABLE 3-continued

Sequences of genomic targets to selectively CRISPR-kill *E. coli* strains. The 3-bp PAM are shown in bold type (ie, tgg, agg and cgg). Expected SpCas9-gRNA cleavage sites are indicated by vertical arrows (3 bp upstream of the PAM).

Target strain	Target gene	Genomic target sequence
<i>E. coli</i> Nissle 1917	yapH	caccgcattactttatg cagg ↓ (SEQ ID NO: 2)
<i>E. coli</i> Nissle 1917	pks	gcgcgggtgtggtt gtcgg ↓ (SEQ ID NO: 3)

TABLE 4

Sequences of genomic targets & Cas to selectively CRISPR-kill *C. difficile*

(a) Overview of design spacers in pMTL84151 - cdCRISPR1 targeting *C. difficile*

Gene	Name	PAM	target
tcdA	TA2	CCA	ACACCTTAACCCAGCCATAGAGTCTGATAATAACTTC (SEQ ID NO: 4)
tcdB	TB2	CCA	GACTTATTTGAGTCTATAGAGAAACCTAGTTCAGTAA (SEQ ID NO: 5)
rRNA-1	TRR1	CCT	TCGACGACTTCTTCCAAAAGGTTAGATAATCGGCTTC (SEQ ID NO: 6)
rRNA-2	TRR2	CCA	GTACAGGATGGACCCGCGTCTGATTAGCTAGTTGGTA (SEQ ID NO: 7)

TABLE 4-continued

Sequences of genomic targets & Cas to selectively CRISPR-kill *C. difficile*

gyrA TARI CCA TCCTCATGGAGATACTGCTGTTTATTATGCTATGGTA (SEQ ID NO: 8)

(b) Cas3 sequence (SEQ ID NO: 9)

Sequence

TCAAATAAATGGTCTATTTCACTTATAAGCACACCTTTACCAAATATCTGTT
 TTGTATGCTTATTTTCGTATATATCATATTTATATAAAAGTATTTTAAATCTTCAA
 GACCTTTCACCTGTATGCAATTACATTTTGTAGCTTTATATATTGGTAAATTTA
 CAGTTTTTTAAATATCTCTCTTCTTGCTTTTTTCTCTTAGACTTTATTTTATTTATC
 AATTGTCTATCATCACTTACTATATGCATTTATAAGCTCTTGTCCTCAACTCTTCATAA
 TCTTTGATTAAGTTTCTCAATTTTATATATATATCTCTGGAATTAAGTATAT
 CCTTCAATATTTCTAAGTATATTTTGTGCATCTTTAGAACCTAAACTATATGGCGTT
 ATAGTATCTAAAATATTTAAAGCACTGGTAAATCTCTTTTCAAAGCTGTACCTTC
 TAGACTTTCTTAGAGTATAGTATCTTAAACATTTCTACCTTATATTTTCTTTTCAT
 CTTCACACATCTTTTCCATTTATAAAAGTTTTCAGTAGCTCAAGTCTCTTTTCTAC
 AATACTCTTACATAAAATACTACCTATTCCTGTAGCTTTTCTGTATATATAAATAT
 ATTTGGGGTATTTTCTTACATCTACGACTTCTATAACATCTACCAAATCGTTGAA
 ATAGACTGTCAAGTGTGAATTTCTGTATGAAGCTCATCAAATCAATATCAAG
 GGATGCTTCCACTAATTTGTAGTAAATCCAATACCATTACTATCACTATCTGCAA
 ATCTTTTATATATTTTCTAATTTTGTCTATCTTCTGTATATACATAGAAATGTA
 AAAGATTAAGTTGACATCTATCTCTTTGGCTTAACTATTTCTTCTATTAACATCAT
 ATTTTCTACAGCACTTTTAACTGTATTTACTATTTACAAGAACTTTTTTATTCATTCT
 CACTTTGTATTTTACCTAAGTTTTCATCTATTGAATTTCTACAATAGACACAC
 AATGCTTATTTTCTGTATTACATGTCAATTCAGCTAAGTTACTATTTCATTACAC
 CTCTTTTTTAAATCACTATATATATGGTTGGCATAGTAGCTGTCAATATCATAA
 ATCTGCCACCTATCTTATGTATCATTTCTATACCTTTTACCAATACAGCTGCTATTT
 CTGGTGAATATGCTTGTATCTCATCTATTACTACTTTTGAATATGCTAATGTTGAGT
 ACACTTTTTCAATACCCTCTATACAAAAAAGGAAATTTAAATATTTGGTCTATTGTA
 GAAAATGTCAGTTTGAAGATAATAACTTTGCTAAATCTACAATCTCACTTGAAT
 TTCTTGATTACTTTCTCTAGATAATCTATTGCTGTTGAATGTAAACAAACCTAAGA
 ATGTATCATTGATTTCTCCAACCTCAACTATATTTTGGTCTATCAAAATATGCAAT
 TTATACTTACTCTAATGGAAGTGTAAAAATGCCTTGTCTTTATCTATCCAAAT
 AAGGCAGTTTCTGTTTTTCCCATCTCTGTAGGTGCAATCAGTATTATATCTTATTT
 CTATTAGATTTAGCAAAATGATGAGCCCTCTCAAACTACCAAACTCTTTTCAATTA
 ATAATTTTCTGCTGTTCTCTATATTTATAACATTTATGCAATCCACAACCTCATG
 AGCAGAAGCACTGTGGTCTAATCTATGTAGTATTTCTTTTAAACATAATATATAAAT
 TATAGTACTTTGATTTTTTATCTATTTCTTTTCTACACTTTGTAGATATACTTTACT
 CAATTTTTCTGTTTTTATGGATATCTAATTTAAATTCATGCTGTAGTTCATAAAT
 TTTATTTATTAATCTTCACTAATATTTTGTATTAATTTTAAATCTTTATCT
 ATAAAGATATCTCTTTCATGATGATATAACAATACTTGATTTAATAGTCTCAAG
 TTCTTTATTTTTTCTGTCTATATACTATAATCAATAAATGCAGGAGAAAGAT
 AATTATGGCCTACATTATTTCTAAATGAGTTACTATTTAGGTTCAATTTATTTAG
 TATCCATTCTGCTTTTTTATTAACCTTTGAAACGGTGAAAATGCTTTTCAAATATCAT
 GAAATCTATAACAAGTCAAGTAAATGCCAAAATATCTCTCTTCTAGAAAATCT
 AAGCTATTTATATTTTTTCCATAACTTTCTCTAATACATTCATTTGTTTTAAAGT
 TCATCAGTATGTTCTCAAGTGTTCCTCACTGGATTAGATTTAGCATATAACAT

Sequences of Cas9 used to selectively kill *E. coli*.

SEQ ID NO: 10 (Cas9 nucleotide sequence)

ATGGATAAGAAATACTCAATAGGCTTAGATATCGGCACAAATAGCGTGGATGGGCGGT
 GATCACTGATGAATATAAGGTTCCGCTCAAAAAGTTCAAGGTTCTGGGAAATACAGACC
 GCCACAGTATCAAAAAAATCTTATAGGGGCTTTTTATTGACAGTGGAGAGACAGCG
 GAAGCGACTCGTCTCAAACGGACAGCTCGTAGAAGGTATACAGCTCGGAAGAAATCGTAT
 TTGTATCTACAGGAGATTTTTCAAATGAGATGGCGAAAGTAGATGATAGTTCTTTCA
 TCGACTTGAAGAGTCTTTTTTGGTGAAGAAGACAGAAGCATGAACGCTCATCCTATTTT
 TGGAAAATATAGTAGATGAAGTTGCTTATCATGAGAAAATCACTATCTATCATCTGCG
 AAAAAATTTGGTAGATTCTACTGATAAAGCGGATTTGCGCTTAACTATTTTGGCCTTAGC
 GCATATGATTAAGTTTCTGTTGTCATTTTTGATTGAGGGAGATTTAAATCCTGATAATAGT
 GATGTGGACAACTATTTATCCAGTTGGTACAAACCTACAATCAATTTTGAAGAAAAC
 CCTATTAACGCAAGTGGAGTAGATGCTAAAGCGATTCTTTCTGCACGATTGAGTAAATCA
 AGACGATTAGAAAATCTCATTTGCTCAGCTCCCGGTGAGAAGAAAATGGCTTATTTGG
 GAATCTCATTTGCTTTTGTATTTGGGTTGACCCCTAATTTTAAATCAAATTTGATTTGGCA
 GAAGATGCTAAATACAGCTTTCAAAGATACTTACGATGATGATTTAGATAATTTATTTG
 GCGAAAATGGAGATCAATATGCTGATTTGTTTTGGCAGCTAAGAAATTTATCAGATGCT
 ATTTTACTTTTCAATATCTAAGAGTAAATACTGAAATAACTAAGGCTCCCTATCAGCT
 TCAATGATTAACGCTACGATGAACATCATCAAGACTTGACTTTTTAAAAGCTTTAGTT
 CGACAACAACTTTCCAGAAAAGTATAAAGAAATCTTTTTGATCAATCAAAAAACGGATA
 TGCAGGTTATATGATGGGGAGCTAGCCAAGAAGAAATTTATAAATTTATCAAACCAAT
 TTTAGAAAATTTGATGGTACTGAGGAATTTGTTGAAACTAAATCGTGAAGATTGCT
 GCGCAAGCAACGGACCTTTGACAACGGCTCTATTTCCCATCAAATTTCACTTTGGGTGAGCT
 GCATGCTATTTGAGAAGACAAGAAGACTTTTATCCATTTTTAAAAGCAATCGTGAGAA
 GATTGAAAATCTTGACTTTTCAATCTCTTATATGTTGGTCCATTGGCGCGTGGCAAT
 AGTCTTTTTGATGATGACTCGGAAGTCTGAAGAAAATAATCCCATGGAATTTTGA
 GAAGTTGTCGATAAAGGTTGCTCAGCTCAATCATTTATGAAACGATGACAACCTTTGAT
 AAAATCTTCAAATGAAAAGTACTACCAAACATAGTTTGTCTTTATGAGTATTTTACG
 GTTTATAACGAATTCAGAAAAGTCAAATATGTTACTGAAGGAATGCGAAAACCGACAT
 TCTTTCAAGGTGACAGAAAGCCATTTGTTGATTTACTCTTCAAACAAATCGAAAAGT

TABLE 4-continued

Sequences of genomic targets & Cas to selectively CRISPR-kill *C. difficile*

AACCGTTAAGCAATAAAAGAAGATTATTTCAAAAAATAGAATGTTTTGATAGTGTGA
AATTTTCAGGAGTTGAAGATAGATTTAATGCCTCATTAGGTACCTACCATGATTTGCCTAAA
AATTATTAAGATAAAGATTTTTTGGATAATGAAGAAAATGAAGATATCTTAGAGGATA
TTGTTTTAACATTGACCTTATTTGAAGATAGGGAGATGATTGAGGAAAAGACTTAAACAT
ATGCTCACCTCTTTGATGATAAGGTGATGAAACAGCTTAAACGTCGCCGTTATACTGGTT
GGGGACGTTTGTCTCGAAAATGATTAATGGTATTAGGGATAAGCAATCTGGCAAAAACA
ATATTAGATTTTTTGAATCAGATGGTTTTGCCAATCGCAATTTTATGCAGCTGATCCATG
ATGATAGTTTGACATTTAAAGAAGACATTTCAAAAAGCACAAGTGTCTGGACAAGCGAT
AGTTTACATGAACATATTGCAAAATTTAGCTGGTAGCCCTGCTATTTAAAAAGGTATTTTA
CAGACTGTAAAAGTTGTTGATGAATTTGGTCAAAGTAATGGGGCCGCAATAAGCCAGAAAA
TATCGTTATTGAATGGCACGTGAAAATCAGACAACCTCAAAGGGCCAGAAAAATTCGC
GAGAGCGTATGAAACGAATCGAAGAAGGTATCAAAGAATTAGGAAGTCAAGTTCTTAAA
GAGCATCTGTGAAAATACTCAATGCAAAATGAAAAGCTCTATCTATATCTCCAA
AATGGAAGAGACATGATGTTGGACCAAGAAATAGATATTAATCGTTTAAAGTATTATGAT
GTCGATCACATTTGTTCCACAAAGTTTCTTAAAGACGATCAATAGACAATAGGTCTTA
ACGCGTTCTGATAAAAACTCGTGGTAAATCGGATAACGTTTCCAAAGTGAAGAGTAGTCAA
AAAGATGAAAACTATTGGAGACAACCTCTAAACGCCAAGTTAATCACTCAACGTAAGT
TTGATAATTTAACGAAAGCTGAACGTTGGAGTTTGGTGAAGTGAATGATAAAGCTGGTTTTA
TCAAACGCCAATTTGTTGAAACTCGCCAAATCACTAAGCATGTGGCACAATTTTGATA
GTCGCATGAATACTAAATACGATGAAAATGATAAACTTATTCGAGAGGTTAAAGTGAAT
ACCTTAAATCTAAATTAGTTTCTGACTCCGAAAAGATTTCCAATCTATAAAGTACGT
GAGATTAACAATTAACATCATGCCCATGATGCGTATCTAAATGCCGTCGTTGGAAGTCT
TTGATTAAGAAATATCCAAAACCTGAATCGGAGTTTGTCTATGGTATTATAAAGTTTAT
GATGTTCTGATAAATGATTGCTAAGTCTGAGCAAGAAATAGGCAAGCAACCCGCAAAAA
TTTCTTTTACTCTAATATCATGAACCTCTTCAAACAGAAATACACTTGCAAAATGGAGA
GATTCGCAAAACGCCCTTAATCGAAACTAATGGGGAAACTGGAGAAATGTTCTGGGATA
AAGGGCGAGATTTTGGCCACAGTGGCCAAAGTATTGTCATGCCCAAGTCAATATTGTCA
AGAAAACAGAAGTACAGACAGGCGGATTTCTCAAGGAGTCAATTTACCAAAAAGAAAT
TCGGACAAGCTTATTTGCTCGTAAAAAAGACTGGGATCCAAAAAATAATGGTGGTTTTGAT
AGTCCAACGGTAGCTTATTAGTCCTAGTGGTGTCTAAGGTGAAAAAGGGAAATCGAA
GAAGTTAAATCCGTTAAAGAGTTACTAGGGATCACAATTAATGGAAGAAAGTTCTTTG
AAAAAATCCGATTGACTTTTTAGAAAGCTAAAGGATATAAGGAAGTTAAAAAAGACTTA
ATCATAAACTACCTAAATATAGTCTTTTTGAGTTAGAAAACGGTTCGTAACCGGATGCTG
GCTAGTGCAGGAGAATTACAAAAAGAAATGAGCTGGCTCTGCCAAGCAAAATATGTGAA
TTTTTTATATTTAGTCTAGTATTATGAAAAGTTGAAGGGTAGTCCAGAAGATAACGAACA
AAAACAATTTGTTTGGAGCAGCATAAGCATTATTAGATGAGATTATTGAGCAAAATCAG
TGAATTTCTAAGCGTGTATTTTAGCAGATGCCAATTTAGATAAAGTTCTTAGTGCATAT
ACAACAATAGAGACAAAACCAATACGTTGAAACAAGCAGAAAAATATTATTCATTATTAC
GTTGACGAATCTGGAGCTCCCGTGTCTTTTAAATATTTGATACAACAATTGATCGTAA
ACGATATACGCTACAAAAGAAGTTTTAGATGCCACTTTATCCATCAATCCATCACTGG
TCTTTATGAAACACGATTGATTTGAGTCAGCTAGGAGGTGACTGA

SEQ ID NO: 11 (Cas9 amino acid sequence)
MDKKS IGLDIGTNSVGVAVI TDEYKVPSSKPKV LGNTDRHS I KKNLIGALLFDSGETAEAT
RLKR TARRRYTRRNRI CYLQEI PFSNEMAKVDDSPFHRLEESFLVEEDKKHERHP I FGNIVDE
VAYHEKYPTI YHLRKLKLDSTDKADLR LIYLALAHMIKFRGHFLI EGDLPDNDSDVDKLF IQL
VQTYNQLFEENP INASGVDAKAI L SARLSKSRLENLI AQLPGEKKNLFGNLIALSLGLTPNF
KSNFDLAEDAKLQSKDYDDLDNLLAQIGDQYADLFLAANKLSDAILLSDI LRVNTEITKA
PLSASMIKRYDEHHQDL TLLKALVRQQLPEKYKE IFFDQSKNGYAGYIDGGASQEEFYKFIKP
ILEKMDGT EELLVKLNREDLLRQR TFDNGS I PHQIHLGELHAILRRQEDFPFLKDNREKIEK
ILTFRI PYYVGPLARGNSRFAMTRKSEETITPWNFEVVDKGASQSFIERMTNFDKNLPNE
KVLPHKSLLYEYPTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLPKTNRKVTVKQLKE
DYFKI ECFDSVEISGVEDRFNASLGT YHDLKIKDKDFLDNEENEDI LEDIVLTLTFEDRE
MIEERLKYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTI LDFLKSDFANRN
FMQLIHDDSLTFKEDIQKAQVSGQDLSHEHIANLAGSPA I KKGILQTVKVVDELVKVMGRH
KPENIV IEMARENQTTQKQKNSRERMKRI BEGIKELGSQILKEHPVENTQLQNEKLYLYLQ
NGRDMYVDQELINRLSDYVDHIVPQSF LKDDSDNKVLRSDKNRGSFNDVPEEVEVKK
MKNYWRQLLNAKLI TQRKFDNLTKAERGGSELDKAGFIKQRLVETRQITKHVAQILD SRM
NTKYDENDKLIREVKVITLKS KLVSDFRKDFQFYKVI INNYHHAHDAYLNAVGTALIKKY
PKLESEFVYGDYKVDVRKMI AKSQEIGKATAKYFFYSNIMNPFKTEITLANG EIRKRPLIET
NGETGEIVWDKGRDFATVRKVL SMPQVNI VKKTEVQTGGFSKESILPKRNSDKLIARKKDW
DPKYYGGFDSPTVAYSVLVVAKEVKGSKKLSVKELLGITIMERSSFEKNP I DFL EAKGYE
VKKDLI IKLPKYSLEFENGRKMLASAGELQKGNELALPSKYVNFYLASHYEKLGKSPED
NEQKQLFVEQHKHYLDEIIEQISEFSKRVI LADANLDKVL SAYNKHDKPIREQAENI IHLFTL
TNLGAPAAFKYFDTTIDRKRYTSTKEVLDTLIHQSIITGLYETRIDLSQLGGD

115

TABLE 5

Publication	Gram-neg. bacteria as causative infectious pathogen in cancer patients	<i>E. coli/K. pneumoniae/ P. aeruginosa</i> as causative pathogens in bacteraemia in cancer patients [fraction of Gram-neg. cases]
Samonis et al	65%	54% [85%]
Velasco et al	45%	33% [74%]

116

TABLE 5-continued

Publication	Gram-neg. bacteria as causative infectious pathogen in cancer patients	<i>E. coli/K. pneumoniae/ P. aeruginosa</i> as causative pathogens in bacteraemia in cancer patients [fraction of Gram-neg. cases]
Marín et al	55%	51% [92%]
Anatoliotaki et al	47%	34% [73%]

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 1

taggtgaagt ccctcgcgga tgg

23

<210> SEQ ID NO 2

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 2

cacccgcatt acctttatgc agg

23

<210> SEQ ID NO 3

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 3

gcgcggtgt ggttgtgctt cgg

23

<210> SEQ ID NO 4

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 4

acaccttaac ccagccatag agtctgataa taacttc

37

<210> SEQ ID NO 5

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 5

gacttatattg agtctataga gaaacctagt tcagtaa

37

<210> SEQ ID NO 6

<211> LENGTH: 37

-continued

<212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 6

 tcgacgactt cttccaaaag gttagataat cggtctc 37

<210> SEQ ID NO 7
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 7

 gtacaggatg gaccgcgctc tgattagcta gttggta 37

<210> SEQ ID NO 8
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 8

 tctctatgga gatactgctg tttattatgc tatggta 37

<210> SEQ ID NO 9
 <211> LENGTH: 2322
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 9

 tcaataaat tggctatatt cattaactat aagcacacct ttaccaata tctgttttgt 60
 atgcttattt tcgtatatat catatttata taaaagtatt tttaaatctt caagaccttt 120
 cacctgtatg tcaattacat tttgtttagc tttatatatt ggtaaattha cagttttttt 180
 aattatctct cttcttgctt tttttctctt agactttatt ttattttatca attgtctatc 240
 atcattacta tatgcattta taagctcttg tccaactctc tcataatctt tgattaaagt 300
 ttcttcaatt ttattatata taccctctgg aattactgta tatccttcaa tattttetaag 360
 tatattttgt gcactcttag aaacctaaact atatggcggt atagtatcta aaatatttaa 420
 agcactggta aatctctttt caaaagctgt accttctaga ctttctttag agtatagtat 480
 cttaaccatt tctaccttat atttttcttt catcttcaca cattcttttc catttataaa 540
 agttttcagt agctcaagtc cttttctctc aatatactta tcataaatac taacctattcc 600
 tgtagctttt tctgtatata taaatatatt tgggctattt tcttcatact cactgacttct 660
 ataacatcta ccaaatcggt gaaatagact gtcaagtggt gaattttctg tatgaagctc 720
 atcaaaatca atatcaaggg atgcttccac taattgtgta gtaatccaaa taccattact 780
 atcaactatc gcaaatctt ttatatattt ttctaatttt gctctatctt cttgtatata 840
 catagaatgt aaaagattta agttgacatc tattctcttt ggcttaacta tttcttctat 900
 taactcatat tttctacag cacttttaac tgtatttact attacaagaa cttttttatt 960
 cattccactt tgtattattt tacctaagtt ttcactatatt gaattttcta caatagacac 1020
 acaatgtctt atttttctg tattacatgt caattcagct aagttaactat tcattacacc 1080

-continued

tctttttttt aattcatcta tatatatggg tggcatagta gctgtcatta tcataaatct	1140
gccacctatc ttatgtatca tttctatacc ttttaccat acagctgcta tttctgggta	1200
atatgcttgt atctcatcta ttactacttt tgaatatgct aatggtgagt acactttttc	1260
ataccctcta tacaaaaaag gaaatttaa tatttggctc attgtagaaa atgtcagttt	1320
gcaagataat aactttgcta aatctacaat ctcaactgaa ttttcttgat tactttcttc	1380
tagataatct attgctgttg aatgtaacaa acctaagaat gtatcatttg attctccaac	1440
tccaactata ttttttggtc tatcaataa tgcatttata cttactctta atggaagtgt	1500
aaaaaatgcc ttgtctttat ctatccaaat taaggcagtt tctgtttttc ccattctctg	1560
aggtgcaatc agtattatat tcttatttct attagattta gcaaatgatt gagcctctct	1620
caaaactacca aactctttca ttaaataatt ttctgtctgt tctcctatat ttataacatt	1680
attgcattcc acaacttcat gaggcagaagc actgtggctc aatctatgta gtattccttt	1740
taacataata tataaattat agtacttggg atttttatct attctttttt ctacactttg	1800
tagatatact ttactcaatt tttctgtttt tattggatat ctaactttaa attcatgctg	1860
tagttcataa actttattta ttaaactctc atctaataatt ttttgtatta aatttttaaa	1920
atctttatct ataagatat ctctttcatg atgatataca ataacttgat ttaatactgc	1980
tctaagttct ttattttttt tctctgtctat ataactataa tcaataaatg caggagaaa	2040
ataattatgg cctacattat tttctaaatg agttactatt ttaggttcat ttatttttagt	2100
atccattctg ctttttatta actcttggaa cggtgaaaa gcttttccaa tatcatgaaa	2160
ttctataaca aagcaagta attgccaaaa tatctctctc tctagaaaat ctaagctatt	2220
tatatttttt ccataacttt ctcttaatac attcatttgt tttaaaagtt catcagtatg	2280
ttctctaagt gtttccactg gatttagattt agcatataac at	2322

<210> SEQ ID NO 10
 <211> LENGTH: 4107
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 10

atggataaga aataactcaat aggccttagat atcggcacia atagcgtcgg atgggcgggtg	60
atcactgatg aatataaggt tccgtctaaa aagttcaagg ttctgggaaa tacagaccgc	120
cacagtatca aaaaaaatct tataggggct cttttatttg acagtggaga gacagcggaa	180
gcgactcgtc tcaaacggac agctcgtaga aggtatacac gtcggaagaa tcgtatttgt	240
tatctacagg agattttttc aaatgagatg gcgaaagtag atgatagttt ctttcatcga	300
cttgaagagt cttttttggg ggaagaagac aagaagcatg aacgtcatcc tatttttggg	360
aatatagtag atgaagtggc ttatcatgag aaatatccaa ctatctatca tctgcgaaaa	420
aaattggtag attctactga taaagcggat ttgcgcttaa tctatttggc cttagcgcct	480
atgattaagt ttcgtgggca ttttttgatt gagggagatt taaatcctga taatagtgat	540
gtggacaaac tatttatcca gttgttacia acctacaatc aattatttga agaaaaccct	600
attaacgcaa gtggagtaga tgcataaagc attctttctg cacgattgag taaatcaaga	660
cgattagaaa atctcattgc tcagctcccc ggtgagaaga aaaatggctt atttgggaat	720
ctcattgctt tgtcattggg tttgacccct aattttaaat caaattttga tttggcagaa	780
gatgctaact tacagcttcc aaaagatact tacgatgatg atttagataa tttattggcg	840

-continued

caaattggag atcaatatgc tgatttggtt ttggcagcta agaatttatac agatgctatt	900
ttactttcag atacctaag agtaaatact gaaataacta aggcctccct atcagettca	960
atgattaaac gctacgatga acatcatcaa gacttgactc ttttaaaagc tttagttcga	1020
caacaacttc cagaaaagta taaagaaatc ttttttgatc aatcaaaaaa cggatatgca	1080
ggttatatg atgggggagc tagccaagaa gaattttata aatttatcaa accaatttta	1140
gaaaaatgg atggctactga ggaattattg gtgaaactaa atcgtgaaga tttgctgcgc	1200
aagcaacgga cctttgacaa cggctctatt ccccatcaaa ttcacttggg tgagctgcat	1260
gctattttga gaagacaaga agacttttat ccatttttaa aagacaatcg tgagaagatt	1320
gaaaaatct tgacttttcg aattccttat tatgttggtc cattggcgcg tggcaatagt	1380
cgttttgcat ggtgactcg gaagtctgaa gaaacaatta ccccatggaa ttttgaagaa	1440
gttgcgata aaggtgcttc agctcaatca tttattgaac gcatgacaaa ctttgataaa	1500
aatcttccaa atgaaaaagt actacaaaaa catagtttgc tttatgagta ttttaccggt	1560
tataacgaat tgacaaaagt caaatatgtt actgaaggaa tgcgaaaaac agcatttctt	1620
tcaggtgaac agaagaaagc cattgtgat ttactctca aaacaaatcg aaaagtaacc	1680
gttaagcaat taaagaaga ttatttcaaa aaaatagaat gttttgatag tgttgaatt	1740
tcaggagttg aagatagatt taatgcttca ttaggtacct accatgattt gctaaaaatt	1800
attaagata aagatttttt ggataatgaa gaaaatgaag atatcttaga ggatattgtt	1860
ttaacattga ccttatttga agatagggag atgattgagg aaagacttaa aacatagct	1920
cacctcttg atgataaagt gatgaaacag cttaaacgtc gccgttatac tggttgggga	1980
cgtttgctc gaaaattgat taatggtatt agggataagc aatctggcaa aacaatatta	2040
gattttttga aatcagatgg ttttgccaat cgcaatttta tgcagctgat ccatgatgat	2100
agtttgacat ttaaagaaga cattcaaaaa gcacaagtgt ctggacaagg cgatagtta	2160
catgaacata ttgcaaatat agctggtagc cctgctatta aaaaaggat tttacagact	2220
gtaaaagtgt ttgatgaatt ggtcaaagta atggggcggc ataagccaga aaatctggt	2280
attgaaatgg cacgtgaaaa tcagacaact caaaagggcc agaaaaatc gcgagagcgt	2340
atgaaacgaa tcgaagaagg tatcaagaa ttaggaagtc agattcttaa agagcatcct	2400
gttgaataa ctcaattgca aaatgaaaag ctctatctct attatctcca aaatggaaga	2460
gacatgatg tggaccaaga attagatatt aatcgtttta gtgattatga tgtcgatcac	2520
attgttccac aaagtctct taaagacgat tcaatagaca ataaggctct aacgcgtct	2580
gataaaaaatc gtgtaaatc ggataacgtt ccaagtgaag aagtagtcaa aaagatgaaa	2640
aactattgga gacaacttct aaacgccaaag ttaatcactc aacgtaagtt tgataattta	2700
acgaaagctg aacgtggagg tttgagtga cttgataaag ctggttttat caaacgcca	2760
ttggttgaat ctcgccaaat cactaagcat gtggcacaaa ttttggatag tcgcatgaa	2820
actaaatag atgaaaatga taaacttatt cgagaggtta aagtgattac cttaaaaatc	2880
aaattagtt ctgacttccg aaaagatttc caattctata aagtacgtga gattaacaat	2940
taccatcatg cccatgatgc gtatctaaat gccgtcgttg gaactgcttt gattaagaaa	3000
tatccaaaac ttgaatcgga gtttgtctat ggtgattata aagtttatga tgttcgtaaa	3060
atgattgcta agtctgagca agaaatagc aaagcaaccg caaaatattt cttttactct	3120
aatatcatga acttcttcaa aacagaaatt acacttgcaa atggagagat tcgcaaacgc	3180

-continued

```

cctctaactcg aaactaatgg ggaactgga gaaattgtct gggataaagg gcgagatttt 3240
gccacagtgc gcaagtatt gtccatgcc caagtcaata ttgtcaagaa aacagaagta 3300
cagacaggcg gattctccaa ggagtcaatt ttaccaaaaa gaaattcggg caagcttatt 3360
gctcgtaaaa aagactggga tccaaaaaaa tatggtggtt ttgatagtcc aacggtagct 3420
tattcagtcc tagtggttgc taaggtggaa aaagggaaat cgaagaagtt aaaatccggt 3480
aaagagttac tagggatcac aattatggaa agaagttcct ttgaaaaaaa tccgattgac 3540
tttttagaag ctaaaggata taaggaagtt aaaaaagact taatcattaa actacctaaa 3600
tatagtcttt ttgagttaga aaacggtcgt aaacggatgc tggctagtgc cggagaatta 3660
caaaaaggaa atgagctggc tctgccaaagc aaatatgtga attttttata tttagctagt 3720
cattatgaaa agttgaaggg tagtccagaa gataacgaac aaaaacaatt gtttgtggag 3780
cagcataagc attatttaga tgagattatt gagcaaatca gtgaattttc taagcgtggt 3840
attttagcag atgccaatth agataaagtt cttagtgcac ataacaaca tagagacaaa 3900
ccaatacgtg aacaagcaga aaatattatt catttattta cgttgacgaa tcttgagct 3960
cccgtgctt ttaaatattt tgatacaaca attgatcgta aacgatatac gtctacaaaa 4020
gaagttttag atgccactct tatccatcaa tccatcaactg gtctttatga aacacgcatt 4080
gatttgagtc agctaggagg tgactga 4107
    
```

```

<210> SEQ ID NO 11
<211> LENGTH: 1368
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
    
```

<400> SEQUENCE: 11

```

Met Asp Lys Lys Tyr Ser Ile Gly Leu Asp Ile Gly Thr Asn Ser Val
1           5           10           15
Gly Trp Ala Val Ile Thr Asp Glu Tyr Lys Val Pro Ser Lys Lys Phe
20           25           30
Lys Val Leu Gly Asn Thr Asp Arg His Ser Ile Lys Lys Asn Leu Ile
35           40           45
Gly Ala Leu Leu Phe Asp Ser Gly Glu Thr Ala Glu Ala Thr Arg Leu
50           55           60
Lys Arg Thr Ala Arg Arg Arg Tyr Thr Arg Arg Lys Asn Arg Ile Cys
65           70           75           80
Tyr Leu Gln Glu Ile Phe Ser Asn Glu Met Ala Lys Val Asp Asp Ser
85           90           95
Phe Phe His Arg Leu Glu Glu Ser Phe Leu Val Glu Glu Asp Lys Lys
100          105          110
His Glu Arg His Pro Ile Phe Gly Asn Ile Val Asp Glu Val Ala Tyr
115          120          125
His Glu Lys Tyr Pro Thr Ile Tyr His Leu Arg Lys Lys Leu Val Asp
130          135          140
Ser Thr Asp Lys Ala Asp Leu Arg Leu Ile Tyr Leu Ala Leu Ala His
145          150          155          160
Met Ile Lys Phe Arg Gly His Phe Leu Ile Glu Gly Asp Leu Asn Pro
165          170          175
Asp Asn Ser Asp Val Asp Lys Leu Phe Ile Gln Leu Val Gln Thr Tyr
180          185          190
Asn Gln Leu Phe Glu Glu Asn Pro Ile Asn Ala Ser Gly Val Asp Ala
    
```

-continued

195				200				205							
Lys	Ala	Ile	Leu	Ser	Ala	Arg	Leu	Ser	Lys	Ser	Arg	Arg	Leu	Glu	Asn
210						215						220			
Leu	Ile	Ala	Gln	Leu	Pro	Gly	Glu	Lys	Lys	Asn	Gly	Leu	Phe	Gly	Asn
225					230					235					240
Leu	Ile	Ala	Leu	Ser	Leu	Gly	Leu	Thr	Pro	Asn	Phe	Lys	Ser	Asn	Phe
				245					250					255	
Asp	Leu	Ala	Glu	Asp	Ala	Lys	Leu	Gln	Leu	Ser	Lys	Asp	Thr	Tyr	Asp
			260					265					270		
Asp	Asp	Leu	Asp	Asn	Leu	Leu	Ala	Gln	Ile	Gly	Asp	Gln	Tyr	Ala	Asp
		275				280						285			
Leu	Phe	Leu	Ala	Ala	Lys	Asn	Leu	Ser	Asp	Ala	Ile	Leu	Leu	Ser	Asp
290						295					300				
Ile	Leu	Arg	Val	Asn	Thr	Glu	Ile	Thr	Lys	Ala	Pro	Leu	Ser	Ala	Ser
305					310					315					320
Met	Ile	Lys	Arg	Tyr	Asp	Glu	His	His	Gln	Asp	Leu	Thr	Leu	Leu	Lys
				325					330						335
Ala	Leu	Val	Arg	Gln	Gln	Leu	Pro	Glu	Lys	Tyr	Lys	Glu	Ile	Phe	Phe
			340					345					350		
Asp	Gln	Ser	Lys	Asn	Gly	Tyr	Ala	Gly	Tyr	Ile	Asp	Gly	Gly	Ala	Ser
		355					360					365			
Gln	Glu	Glu	Phe	Tyr	Lys	Phe	Ile	Lys	Pro	Ile	Leu	Glu	Lys	Met	Asp
370						375					380				
Gly	Thr	Glu	Glu	Leu	Leu	Val	Lys	Leu	Asn	Arg	Glu	Asp	Leu	Leu	Arg
385					390					395					400
Lys	Gln	Arg	Thr	Phe	Asp	Asn	Gly	Ser	Ile	Pro	His	Gln	Ile	His	Leu
				405					410					415	
Gly	Glu	Leu	His	Ala	Ile	Leu	Arg	Arg	Gln	Glu	Asp	Phe	Tyr	Pro	Phe
			420					425					430		
Leu	Lys	Asp	Asn	Arg	Glu	Lys	Ile	Glu	Lys	Ile	Leu	Thr	Phe	Arg	Ile
		435				440						445			
Pro	Tyr	Tyr	Val	Gly	Pro	Leu	Ala	Arg	Gly	Asn	Ser	Arg	Phe	Ala	Trp
450						455					460				
Met	Thr	Arg	Lys	Ser	Glu	Glu	Thr	Ile	Thr	Pro	Trp	Asn	Phe	Glu	Glu
465					470					475				480	
Val	Val	Asp	Lys	Gly	Ala	Ser	Ala	Gln	Ser	Phe	Ile	Glu	Arg	Met	Thr
				485					490					495	
Asn	Phe	Asp	Lys	Asn	Leu	Pro	Asn	Glu	Lys	Val	Leu	Pro	Lys	His	Ser
		500						505					510		
Leu	Leu	Tyr	Glu	Tyr	Phe	Thr	Val	Tyr	Asn	Glu	Leu	Thr	Lys	Val	Lys
		515					520					525			
Tyr	Val	Thr	Glu	Gly	Met	Arg	Lys	Pro	Ala	Phe	Leu	Ser	Gly	Glu	Gln
530						535					540				
Lys	Lys	Ala	Ile	Val	Asp	Leu	Leu	Phe	Lys	Thr	Asn	Arg	Lys	Val	Thr
545					550					555				560	
Val	Lys	Gln	Leu	Lys	Glu	Asp	Tyr	Phe	Lys	Lys	Ile	Glu	Cys	Phe	Asp
				565					570					575	
Ser	Val	Glu	Ile	Ser	Gly	Val	Glu	Asp	Arg	Phe	Asn	Ala	Ser	Leu	Gly
			580					585					590		
Thr	Tyr	His	Asp	Leu	Leu	Lys	Ile	Ile	Lys	Asp	Lys	Asp	Phe	Leu	Asp
				595			600					605			
Asn	Glu	Glu	Asn	Glu	Asp	Ile	Leu	Glu	Asp	Ile	Val	Leu	Thr	Leu	Thr
610						615					620				

-continued

Leu Phe Glu Asp Arg Glu Met Ile Glu Glu Arg Leu Lys Thr Tyr Ala
 625 630 635 640
 His Leu Phe Asp Asp Lys Val Met Lys Gln Leu Lys Arg Arg Arg Tyr
 645 650 655
 Thr Gly Trp Gly Arg Leu Ser Arg Lys Leu Ile Asn Gly Ile Arg Asp
 660 665 670
 Lys Gln Ser Gly Lys Thr Ile Leu Asp Phe Leu Lys Ser Asp Gly Phe
 675 680 685
 Ala Asn Arg Asn Phe Met Gln Leu Ile His Asp Asp Ser Leu Thr Phe
 690 695 700
 Lys Glu Asp Ile Gln Lys Ala Gln Val Ser Gly Gln Gly Asp Ser Leu
 705 710 715 720
 His Glu His Ile Ala Asn Leu Ala Gly Ser Pro Ala Ile Lys Lys Gly
 725 730 735
 Ile Leu Gln Thr Val Lys Val Val Asp Glu Leu Val Lys Val Met Gly
 740 745 750
 Arg His Lys Pro Glu Asn Ile Val Ile Glu Met Ala Arg Glu Asn Gln
 755 760 765
 Thr Thr Gln Lys Gly Gln Lys Asn Ser Arg Glu Arg Met Lys Arg Ile
 770 775 780
 Glu Glu Gly Ile Lys Glu Leu Gly Ser Gln Ile Leu Lys Glu His Pro
 785 790 795 800
 Val Glu Asn Thr Gln Leu Gln Asn Glu Lys Leu Tyr Leu Tyr Tyr Leu
 805 810 815
 Gln Asn Gly Arg Asp Met Tyr Val Asp Gln Glu Leu Asp Ile Asn Arg
 820 825 830
 Leu Ser Asp Tyr Asp Val Asp His Ile Val Pro Gln Ser Phe Leu Lys
 835 840 845
 Asp Asp Ser Ile Asp Asn Lys Val Leu Thr Arg Ser Asp Lys Asn Arg
 850 855 860
 Gly Lys Ser Asp Asn Val Pro Ser Glu Glu Val Val Lys Lys Met Lys
 865 870 875 880
 Asn Tyr Trp Arg Gln Leu Leu Asn Ala Lys Leu Ile Thr Gln Arg Lys
 885 890 895
 Phe Asp Asn Leu Thr Lys Ala Glu Arg Gly Gly Leu Ser Glu Leu Asp
 900 905 910
 Lys Ala Gly Phe Ile Lys Arg Gln Leu Val Glu Thr Arg Gln Ile Thr
 915 920 925
 Lys His Val Ala Gln Ile Leu Asp Ser Arg Met Asn Thr Lys Tyr Asp
 930 935 940
 Glu Asn Asp Lys Leu Ile Arg Glu Val Lys Val Ile Thr Leu Lys Ser
 945 950 955 960
 Lys Leu Val Ser Asp Phe Arg Lys Asp Phe Gln Phe Tyr Lys Val Arg
 965 970 975
 Glu Ile Asn Asn Tyr His His Ala His Asp Ala Tyr Leu Asn Ala Val
 980 985 990
 Val Gly Thr Ala Leu Ile Lys Lys Tyr Pro Lys Leu Glu Ser Glu Phe
 995 1000 1005
 Val Tyr Gly Asp Tyr Lys Val Tyr Asp Val Arg Lys Met Ile Ala Lys
 1010 1015 1020
 Ser Glu Gln Glu Ile Gly Lys Ala Thr Ala Lys Tyr Phe Phe Tyr Ser
 1025 1030 1035 1040

-continued

Asn Ile Met Asn Phe Phe Lys Thr Glu Ile Thr Leu Ala Asn Gly Glu
 1045 1050 1055

Ile Arg Lys Arg Pro Leu Ile Glu Thr Asn Gly Glu Thr Gly Glu Ile
 1060 1065 1070

Val Trp Asp Lys Gly Arg Asp Phe Ala Thr Val Arg Lys Val Leu Ser
 1075 1080 1085

Met Pro Gln Val Asn Ile Val Lys Lys Thr Glu Val Gln Thr Gly Gly
 1090 1095 1100

Phe Ser Lys Glu Ser Ile Leu Pro Lys Arg Asn Ser Asp Lys Leu Ile
 1105 1110 1115 1120

Ala Arg Lys Lys Asp Trp Asp Pro Lys Lys Tyr Gly Gly Phe Asp Ser
 1125 1130 1135

Pro Thr Val Ala Tyr Ser Val Leu Val Val Ala Lys Val Glu Lys Gly
 1140 1145 1150

Lys Ser Lys Lys Leu Lys Ser Val Lys Glu Leu Leu Gly Ile Thr Ile
 1155 1160 1165

Met Glu Arg Ser Ser Phe Glu Lys Asn Pro Ile Asp Phe Leu Glu Ala
 1170 1175 1180

Lys Gly Tyr Lys Glu Val Lys Lys Asp Leu Ile Ile Lys Leu Pro Lys
 1185 1190 1195 1200

Tyr Ser Leu Phe Glu Leu Glu Asn Gly Arg Lys Arg Met Leu Ala Ser
 1205 1210 1215

Ala Gly Glu Leu Gln Lys Gly Asn Glu Leu Ala Leu Pro Ser Lys Tyr
 1220 1225 1230

Val Asn Phe Leu Tyr Leu Ala Ser His Tyr Glu Lys Leu Lys Gly Ser
 1235 1240 1245

Pro Glu Asp Asn Glu Gln Lys Gln Leu Phe Val Glu Gln His Lys His
 1250 1255 1260

Tyr Leu Asp Glu Ile Ile Glu Gln Ile Ser Glu Phe Ser Lys Arg Val
 1265 1270 1275 1280

Ile Leu Ala Asp Ala Asn Leu Asp Lys Val Leu Ser Ala Tyr Asn Lys
 1285 1290 1295

His Arg Asp Lys Pro Ile Arg Glu Gln Ala Glu Asn Ile Ile His Leu
 1300 1305 1310

Phe Thr Leu Thr Asn Leu Gly Ala Pro Ala Ala Phe Lys Tyr Phe Asp
 1315 1320 1325

Thr Thr Ile Asp Arg Lys Arg Tyr Thr Ser Thr Lys Glu Val Leu Asp
 1330 1335 1340

Ala Thr Leu Ile His Gln Ser Ile Thr Gly Leu Tyr Glu Thr Arg Ile
 1345 1350 1355 1360

Asp Leu Ser Gln Leu Gly Gly Asp
 1365

The invention claimed is:

1. A method for treating an acute pathogenic bacterial infection in a subject caused by first bacteria that are *E. coli*, *Pseudomonas aeruginosa* or *Klebsiella* bacteria, the method comprising selectively killing the first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by a guide RNA to cut the target site, wherein the subject has a cancer, wherein the method is carried out no more than 7 days before or after administration of an immunotherapy to the subject, wherein the programmed Cas nuclease treats the infection and the immunotherapy treats the cancer;

55 wherein:

(a) the immunotherapy comprises administering to the subject an anti-PD-1/PD-L1 axis antibody; and

(b) the cancer is selected from melanoma; renal cell carcinoma; bladder cancer; a solid tumour; non-small cell lung cancer (NSCLC); forehead and neck squamous cell carcinoma (HNSCC); Hodgkin's lymphoma; a cancer that overexpresses PD-L1 and has no mutations in EGFR or in ALK; colorectal cancer and hepatocellular carcinoma.

2. The method of claim 1, wherein the first bacteria are *E. coli*.

131

3. The method of claim 2, wherein the first bacteria are enterohemorrhagic *E. coli* (EHEC).

4. The method of claim 1, wherein the anti-PD-1/PD-L1 axis antibody is selected from the group consisting of pembrolizumab, nivolumab, atezolimumab, avelumab and durvalumab.

5. The method of claim 1, wherein the treatment of the bacterial infection is carried out simultaneously with the administration of the immunotherapy to the subject.

6. The method of claim 1, wherein the treatment of the bacterial infection is carried out immediately before or after administering the immunotherapy to the subject.

7. The method of claim 1, wherein the method comprises administering to the subject a nucleic acid vector comprising the guide RNA or a DNA encoding the guide RNA.

8. The method of claim 7, comprising administering a second nucleic acid vector to the subject, wherein the second vector encodes the Cas nuclease.

9. The method of claim 7, wherein the Cas nuclease is an endogenous Cas nuclease of the first bacteria.

10. The method of claim 7, wherein the vector is a phage, phagemid, viriophage, virus, plasmid, or transposon.

11. The method of claim 7, wherein the vector is a conjugative plasmid.

12. The method of claim 11, wherein the conjugative plasmid is delivered from carrier bacteria.

13. The method of claim 12, wherein the carrier bacteria are *Lactobacillus* bacteria.

132

14. The method of claim 1, wherein the Cas nuclease is Cas9.

15. The method of claim 1, wherein the Cas nuclease is Cas3.

16. The method of claim 8, wherein the Cas nuclease is Cas9.

17. The method of claim 16, wherein the second vector further comprises a tracrRNA or a DNA encoding a tracrRNA.

18. The method of claim 1, wherein the treatment of the bacterial infection treats in the subject a disease or condition selected from the group consisting of vaginosis, meningitis, pneumonia, urinary tract infection, cystitis, nephritis, gastroenteritis, a skin infection, impetigo, erysipelas, cellulitis, a dental infection, septicaemia and sepsis, or reduces the risk of the disease or condition in the subject, wherein the disease or condition is mediated by the first bacteria.

19. The method of claim 1, wherein the guide RNA does not program a Cas nuclease to cut the genome of second bacteria in the subject, wherein the second bacteria are selected from the group consisting of *Akkermansia*, *Alistipes*, *Bacteroides*, *Barnesiella*, *Bifidobacterium*, *Clostridium*, *Collinsella*, *Enterococcus*, *Fusobacterium*, *Lactobacillus*, *Propionibacterium*, *Ruminococcus*, *Segmented filamentous bacteria* (SFB); *Veillonella*, *Prevotella*, *Escherichia* and *Streptococcus* bacteria.

* * * * *