



US010920222B2

(12) **United States Patent**
Sommer et al.

(10) **Patent No.:** **US 10,920,222 B2**

(45) **Date of Patent:** **Feb. 16, 2021**

(54) **TREATING AND PREVENTING MICROBIAL INFECTIONS**

(71) Applicant: **SNIPR Biome ApS**, Copenhagen Ø (DK)

(72) Inventors: **Morten Sommer**, Copenhagen Ø (DK); **Virginia Martinez**, Copenhagen Ø (DK); **Eric Van Der Helm**, Copenhagen Ø (DK); **Jakob Krause Haaber**, Copenhagen Ø (DK); **Ana De Santiago Torio**, Copenhagen Ø (DK); **Christian Grøndahl**, Copenhagen Ø (DK); **Jasper Clube**, London (GB)

(73) Assignee: **SNIPR Biome ApS**, Copenhagen Ø (DK)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/700,856**

(22) Filed: **Dec. 2, 2019**

(65) **Prior Publication Data**

US 2020/0087660 A1 Mar. 19, 2020

Related U.S. Application Data

(63) Continuation of application No. 15/967,484, filed on Apr. 30, 2018, now Pat. No. 10,760,075.

(51) **Int. Cl.**

A61K 39/00 (2006.01)
A61K 39/38 (2006.01)
A61K 39/02 (2006.01)
A61K 39/09 (2006.01)
A61K 39/085 (2006.01)
A61K 39/10 (2006.01)
A61K 39/08 (2006.01)
A61K 39/102 (2006.01)
A61K 39/118 (2006.01)
C12N 15/11 (2006.01)
A61P 31/04 (2006.01)
A61K 39/395 (2006.01)
A61K 38/46 (2006.01)
C07K 16/28 (2006.01)

(52) **U.S. Cl.**

CPC **C12N 15/11** (2013.01); **A61K 38/465** (2013.01); **A61K 39/3955** (2013.01); **A61K 39/39558** (2013.01); **A61P 31/04** (2018.01); **C07K 16/2818** (2013.01); **C07K 16/2827** (2013.01); **C12N 2310/20** (2017.05); **C12N 2320/30** (2013.01); **C12N 2320/31** (2013.01); **Y02A 50/30** (2018.01)

(58) **Field of Classification Search**

CPC **C12N 15/11**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,626,504 A 12/1986 Puhler
4,870,287 A 9/1989 Cole
5,633,154 A 5/1997 Schaefer
5,760,395 A 6/1998 Johnstone
5,844,905 A 12/1998 Mckay
5,885,796 A 3/1999 Linsley
6,207,156 B1 3/2001 Kuchroo
7,459,272 B2 12/2008 Morris
8,003,323 B2 8/2011 Morris
8,008,449 B2 8/2011 Korman
8,017,114 B2 9/2011 Korman
8,119,129 B2 2/2012 Jure-kunkel
8,241,498 B2 8/2012 Summer
8,252,576 B2 8/2012 Campbell
8,329,867 B2 12/2012 Lazar
8,354,509 B2 1/2013 Carven
8,735,553 B1 5/2014 Li
8,906,682 B2 12/2014 June
8,911,993 B2 12/2014 June
8,916,381 B1 12/2014 June
8,975,071 B1 3/2015 June
9,101,584 B2 8/2015 June

(Continued)

FOREIGN PATENT DOCUMENTS

EP 2320940 B1 5/2011
EP 2325332 A1 5/2011

(Continued)

OTHER PUBLICATIONS

Jiang et al., Nature Biotechnology, 2013, 31(3): 233-241 (Year: 2013).*

Barrango, R. et al. (Mar. 2007). "CRISPR Provides Acquired Resistance Against Viruses in Prokaryotes," Science, 315:1709-1712.

Bryksin, A. V. et al. (Oct. 8, 2010). "Rational Design of a Plasmid Origin That Replicates Efficiently in Both Gram-Positive and Gram-Negative Bacteria," PloS One 5(10):e13244, 9 pages.

(Continued)

Primary Examiner — Gary B Nickol

Assistant Examiner — Lokia J Jackson-Tongue

(74) Attorney, Agent, or Firm — Morrison & Foerster LLP

(57) **ABSTRACT**

The invention provides methods for treating or preventing microbial (eg, bacterial) infections and means for performing these methods. In particular, treatment of infections requiring rapid and durable therapy is made possible, such as for treating acute conditions such as septicemia, sepsis, SIRS or septic shock. The invention is particularly useful, for example, for treatment of microbes such as for environmental, food and beverage use. The invention relates inter alia to methods of controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate or fluid in an industrial or domestic system. The invention also useful for the treatment of pathogenic bacterial infections in subjects receiving a treatment for a disease or condition, such as a transplant or a treatment for cancer, a viral infection or an autoimmune disease.

28 Claims, 9 Drawing Sheets

Specification includes a Sequence Listing.

(56)	References Cited			2016/0024510	A1	1/2016	Bikard	
	U.S. PATENT DOCUMENTS			2016/0081314	A1	3/2016	Thurston	
	9,102,760	B2	8/2015 June	2016/0115488	A1	4/2016	Zhang	
	9,102,761	B2	8/2015 June	2016/0115489	A1	4/2016	Zhang	
	9,113,616	B2	8/2015 Stevens	2016/0130355	A1	5/2016	June	
	9,328,156	B2	5/2016 June	2016/0159905	A1	6/2016	Abdiche	
	9,464,140	B2	10/2016 June	2016/0159907	A1	6/2016	June	
	9,481,728	B2	11/2016 June	2016/0160186	A1	6/2016	Parsley	
	9,499,629	B2	11/2016 June	2016/0194404	A1	7/2016	June	
	9,518,123	B2	12/2016 June	2016/0208012	A1	7/2016	June	
	9,540,445	B2	1/2017 June	2016/0237455	A1	8/2016	Glucksmann	
	9,701,964	B2	7/2017 Clube	2016/0324938	A1	11/2016	Bikard	
	9,758,583	B2	9/2017 Wang	2016/0333348	A1	11/2016	Clube	
	9,822,372	B2	11/2017 Zhang	2016/0345578	A1	12/2016	Barrangou	
	9,879,269	B2	1/2018 Barrangou	2016/0347836	A1	12/2016	Grosso	
	10,066,233	B2	9/2018 Barrangou	2016/0354416	A1	12/2016	Gajewski	
	10,136,639	B2	11/2018 Wuest	2017/0022499	A1	1/2017	Lu	
	10,136,649	B2	11/2018 Barrangou	2017/0037416	A1	2/2017	Barrangou	
	10,195,273	B2	2/2019 Clube	2017/0106026	A1	4/2017	Kovarik	
	10,300,138	B2	5/2019 Clube	2017/0143772	A1	5/2017	Mulder	
	10,300,139	B2	5/2019 Clube	2017/0173085	A1	6/2017	Kovarik	
	10,363,308	B2	7/2019 Clube	2017/0174713	A1	6/2017	Du	
	10,463,049	B2	11/2019 Clube	2017/0175142	A1	6/2017	Zhang	
	10,506,812	B2	12/2019 Clube	2017/0196225	A1	7/2017	Clube	
	10,524,477	B2	1/2020 Clube	2017/0246221	A1	8/2017	Clube	
	10,561,148	B2	2/2020 Clube	2017/0304443	A1	10/2017	Lebwohl	
	10,582,712	B2	3/2020 Clube	2017/0327582	A1	11/2017	Bissonnette	
	10,596,255	B2	3/2020 Clube	2017/0340733	A1	11/2017	Cao	
	10,603,379	B2	3/2020 Clube	2018/0015131	A1	1/2018	Gajewski	
	10,624,349	B2	4/2020 Clube	2018/0055852	A1	3/2018	Kutok	
	10,765,740	B2	9/2020 Clube et al.	2018/0064114	A1	3/2018	Clube	
	2003/0049841	A1	3/2003 Short	2018/0064115	A1	3/2018	Clube	
	2004/0096974	A1	5/2004 Herron	2018/0070594	A1	3/2018	Clube	
	2005/0118719	A1	6/2005 Schmidt	2018/0084785	A1	3/2018	Clube	
	2009/0155768	A1	6/2009 Scholl	2018/0084786	A1	3/2018	Clube	
	2010/0076057	A1	3/2010 Sontheimer	2018/0140698	A1	5/2018	Clube	
	2010/0093617	A1	4/2010 Barrangou	2018/0146681	A1	5/2018	Clube	
	2010/0172874	A1	7/2010 Turnbaugh	2018/0155721	A1	6/2018	Lu	
	2011/0002889	A1	1/2011 Barrangou	2018/0155729	A1	6/2018	Beisel	
	2011/0008369	A1	1/2011 Finnefrock	2018/0179547	A1	6/2018	Zhang	
	2011/0136688	A1	6/2011 Scholl	2018/0200342	A1	7/2018	Bikard	
	2012/0177645	A1	7/2012 Langermann	2018/0273940	A1*	9/2018	Sommer A61K 38/465
	2012/0269859	A1	10/2012 Minato	2018/0303934	A1	10/2018	Clube	
	2012/0294796	A1	11/2012 Johnson	2018/0326057	A1	11/2018	Clube	
	2013/0011828	A1	1/2013 Barrangou	2018/0371405	A1	12/2018	Barrangou	
	2013/0109053	A1	5/2013 Macdonald	2019/0015441	A1	1/2019	Shachar	
	2013/0121968	A1	5/2013 Quay	2019/0021343	A1	1/2019	Barrangou	
	2013/0287748	A1	10/2013 June	2019/0070233	A1	3/2019	Yeung	
	2013/0288368	A1	10/2013 June	2019/0117709	A1	4/2019	Kovarik	
	2013/0309258	A1	11/2013 June	2019/0133135	A1	5/2019	Clube	
	2014/0022021	A1	1/2014 Kusachi	2019/0134194	A1	5/2019	Clube	
	2014/0068797	A1	3/2014 Doudna	2019/0160120	A1	5/2019	Haaber	
	2014/0105912	A1	4/2014 Noelle	2019/0230936	A1	8/2019	Clube	
	2014/0106449	A1	4/2014 June	2019/0240325	A1*	8/2019	Clube A61P 35/00
	2014/0107092	A1	4/2014 Meyerson	2019/0240326	A1	8/2019	Clube	
	2014/0179726	A1	6/2014 Bajaj	2019/0255084	A1	8/2019	Schentag	
	2014/0199767	A1	7/2014 Barrangou	2019/0256900	A1	8/2019	Zhang	
	2014/0234972	A1	8/2014 Zhang	2019/0298779	A1	10/2019	Falb	
	2014/0294898	A1	10/2014 Miller	2019/0321468	A1	10/2019	Clube et al.	
	2014/0341920	A1	11/2014 Noelle	2019/0321469	A1	10/2019	Clube et al.	
	2014/0370017	A1	12/2014 June	2019/0321470	A1	10/2019	Clube	
	2015/0031134	A1	1/2015 Zhang	2019/0359933	A1	11/2019	Swee	
	2015/0050699	A1	2/2015 Siksnys	2020/0030444	A1	1/2020	Clube	
	2015/0050729	A1	2/2015 June	2020/0046773	A1	2/2020	Borody	
	2015/0064138	A1	3/2015 Lu	2020/0068901	A1	3/2020	Clube	
	2015/0093822	A1	4/2015 June	2020/0077663	A1	3/2020	Clube	
	2015/0099299	A1	4/2015 June	2020/0085066	A1	3/2020	Clube	
	2015/0118202	A1	4/2015 June	2020/0102551	A1	4/2020	Barrangou	
	2015/0125463	A1	5/2015 Cogswell	2020/0115716	A1	4/2020	Martinez	
	2015/0132419	A1	5/2015 Arvik	2020/0121787	A1	4/2020	Clube	
	2015/0139943	A1	5/2015 Campana	2020/0128832	A1	4/2020	Clube	
	2015/0140001	A1	5/2015 Lee	2020/0157237	A1	5/2020	Regev	
	2015/0184139	A1	7/2015 Zhang	2020/0164070	A1	5/2020	Clube	
	2015/0290244	A1	10/2015 June	2020/0179460	A1	6/2020	Kovarik	
	2015/0353905	A1	12/2015 Weiss					
	2016/0009805	A1	1/2016 Kowanetz					
	2016/0009813	A1	1/2016 Themeli					

(56)

References Cited

U.S. PATENT DOCUMENTS

2020/0199570 A1 6/2020 Novick
2020/0267992 A1 8/2020 Clube

FOREIGN PATENT DOCUMENTS

EP 2840140 A1 2/2015
EP 3132035 B1 1/2020
EP 3132036 B1 1/2020
EP 3630975 A1 4/2020
EP 3633032 A2 4/2020
EP 3634442 A1 4/2020
EP 3634473 A1 4/2020
RU 2531343 C2 10/2014
WO WO1995001994 A1 1/1995
WO WO1998042752 A1 10/1998
WO WO2000037504 A2 6/2000
WO WO2000037504 A3 6/2000
WO WO2001014424 A2 3/2001
WO WO2001014424 A3 3/2001
WO WO2005003168 A2 1/2005
WO WO2005009465 A1 2/2005
WO WO2005003168 A3 5/2005
WO WO2005046579 A2 5/2005
WO WO2005046579 A3 8/2005
WO WO2006003179 A2 1/2006
WO WO2006003179 A3 5/2006
WO WO2006072625 A2 7/2006
WO WO2006072626 A1 7/2006
WO WO2006121168 A1 11/2006
WO WO2006072625 A3 12/2006
WO WO2007025097 A2 3/2007
WO WO2007042573 A2 4/2007
WO WO2007025097 A3 7/2007
WO WO2007042573 A3 7/2007
WO WO2008084106 A1 7/2008
WO WO2008108989 A2 9/2008
WO WO2008132601 A1 11/2008
WO WO2008108989 A3 3/2009
WO WO2009044273 A2 4/2009
WO WO2009101611 A1 8/2009
WO WO2008084106 A9 9/2009
WO WO2009044273 A3 9/2009
WO WO2009114335 A2 9/2009
WO WO2010011961 A2 1/2010
WO WO2010027827 A2 3/2010
WO WO2010027827 A3 5/2010
WO WO2009114335 A3 6/2010
WO WO2010011961 A3 6/2010
WO WO2010065939 A1 6/2010
WO WO2010075424 A2 7/2010
WO WO2010075424 A3 9/2010
WO WO2011014438 A1 2/2011
WO WO2011066342 A2 6/2011
WO WO2011066342 A3 7/2011
WO WO2012071411 A2 5/2012
WO 2012079000 A1 6/2012
WO 2012079000 A4 8/2012
WO WO2012071411 A3 8/2012
WO WO2012160448 A2 11/2012
WO WO2012164565 A1 12/2012
WO WO2013006490 A2 1/2013
WO WO2013025779 A1 2/2013
WO 2013063361 A1 5/2013
WO WO2012160448 A3 5/2013
WO WO2013067492 A1 5/2013
WO WO2013176772 A1 11/2013
WO 2014012001 A2 1/2014
WO WO2014015252 A1 1/2014
WO WO2014018423 A2 1/2014
WO WO2014018423 A3 1/2014
WO WO2013006490 A3 5/2014
WO WO2014093595 A1 6/2014
WO WO2014093661 A2 6/2014
WO WO2014093661 A3 8/2014
WO WO2014124226 A1 8/2014

WO WO2014093661 A9 10/2014
WO WO2015016718 A1 2/2015
WO WO2015034872 A2 3/2015
WO 2014012001 A3 4/2015
WO 2015058018 A1 4/2015
WO WO2015034872 A3 4/2015
WO 2015070083 A1 5/2015
WO WO2015069682 A2 5/2015
WO WO2015071474 A2 5/2015
WO WO2015075688 A1 5/2015
WO 2015089351 A1 6/2015
WO WO2015088643 A1 6/2015
WO WO2015069682 A3 7/2015
WO WO2015071474 A3 8/2015
WO WO2015136541 A2 9/2015
WO WO2015148680 A1 10/2015
WO WO2015153940 A1 10/2015
WO WO2015155686 A2 10/2015
WO WO2015159068 A1 10/2015
WO WO2015159086 A1 10/2015
WO WO2015159087 A1 10/2015
WO WO2015136541 A3 11/2015
WO WO2015155686 A3 12/2015
WO 2016044745 A1 3/2016
WO WO2016063263 A2 4/2016
WO WO2016063263 A3 6/2016
WO WO2016084088 A1 6/2016
WO WO2016177682 A1 11/2016
WO WO2016196361 A1 12/2016
WO WO2016196605 A1 12/2016
WO WO2016205276 A1 12/2016
WO WO2017009399 A1 1/2017
WO WO2017042347 A1 3/2017
WO WO2017058751 A1 4/2017
WO WO2017112620 A1 6/2017
WO WO2017211753 A1 12/2017
WO WO2018064165 A2 4/2018
WO WO2018081502 A1 5/2018
WO WO2018115519 A1 6/2018
WO WO2018217351 A1 11/2018
WO WO2018217981 A1 11/2018
WO WO2018222969 A1 12/2018
WO WO2018226853 A1 12/2018
WO WO2018064165 A3 6/2019
WO WO2020072248 A1 4/2020
WO WO2020072250 A1 4/2020
WO WO2020072253 A1 4/2020
WO WO2020072254 A1 4/2020

OTHER PUBLICATIONS

Chan, C.T.Y. et al. (Dec. 2015). "'Deadman' and 'Passcode' Microbial Kill Switches for Bacterial Containment," *Nat Chem Biol* 12(2):82-86, 19 pages.
Cheadle, E.J. et al. (2012). "Chimeric Antigen Receptors for T-Cell Based Therapy," *Methods Mol. Biol.* 907:645-666, 36 pages.
Edgar et al. (Dec. 2010). "The *Escherichia coli* CRISPR System Protects From λ Lysogenization, Lysogens, and Prophage Induction," *Journal of Bacteriology* 192(23):6291-6294, Supplemental Material, 2 pages.
Garrett W.S. et al. (Oct. 5, 2007). "Communicable Ulcerative Colitis Induced by T-Bet Deficiency in the Innate Immune System," *Cell* 131(1):33-45, 23 pages.
Goodall, E.C.A. et al. (Feb. 20, 2018). "The Essential Genome of *Escherichia coli* K-12," *Am. Society for Microbiology—mBio* 9(1):e02096-17, 18 pages.
Harrington, L.E. (Nov. 2005, e-pub. Oct. 2, 2005). "Interleukin 17-producing CD4+ Effector T Cells Develop Via a Lineage Distinct From the T Helper Type 1 and 2 Lineages," *Nat. Immunol.* 6(11):1123-1132.
Horvath, P. et al. (2008, e-pub. Dec. 7, 2007). "Diversity, Activity, and Evolution of CRISPR Loci in *Streptococcus thermophilus*," *Journal of Bacteriology* 190(4):1401-1412.
Kaulich, M. et al. (2015, e-pub. Jan. 13, 2015). "Efficient CRISPR-rAAV Engineering of Endogenous Genes to Study Protein Function by Allele-Specific RNAi," *Nucleic Acids Research* 43(7):e45, 8 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Kochenderfer, J.N. et al. (Sep. 2009). "Construction and Pre-clinical Evaluation of an Anti-CD19 Chimeric Antigen Receptor," *J. Immunother.* 32(7):689-702, 26 pages.
- Ludwig, E.S. et al. (1985). "The Phylogenetic Position of *Streptococcus* and Enterococcus," *Journal of General Microbiology* 131:543-551.
- López, P. et al. (Apr. 5, 2016). "Th17 Responses and Natural IgM Antibodies are Related to Gut Microbiota Composition in Systemic Lupus Erythematosus Patients," *Sci. Rep.* 6:24072, 12 pages.
- Magee, M.S. et al. (Nov. 2014). "Challenges to Chimeric Antigen Receptor (CAR)-T Cell Therapy for Cancer," *Discov. Med.* 18(100):265-271, 6 pages.
- Mercenier, A. (1990). "Molecular Genetics of *Streptococcus thermophilus*," *FEMS Microbiology Letters* 87(1-2):61-77.
- Park, H. et al. (2005). "A Distinct Lineage of CD4 T Cells Regulates Tissue Inflammation by Producing Interleukin 17," *Nat. Immunol.* 6(11):1133-1141, 24 pages.
- Ran, F.A. et al. (Apr. 9, 2015). "In vivo Genome Editing Using *Staphylococcus aureus* Cas9," *Nature* 520(7546):186-191, 28 pages.
- Roberts, A.P. et al. (Jun. 2009, e-pub. May 20, 2009). "A Modular Master on the Move: The Tn916 Family of Mobile Genetic Elements," *Trends Microbiol.* 17(6):251-258. Abstract Only.
- Rong, Z. et al. (Mar. 14, 2014). "Homologous Recombination in Human Embryonic Stem Cells Using CRISPR/Cas9 Nickase and a Long DNA Donor Template," *Protein & Cell* 5(4):258-260.
- Samaržija, D. et al. (2001). "Taxonomy, Physiology and Growth of *Lactococcus Lactis*: A Review," *Mljekarstvo* 51(1):35-48.
- Sapranaukas, R. et al. (Nov. 1, 2011, e-pub. Aug. 3, 2011). "The *Streptococcus thermophilus* CRISPR/Cas System Provides Immunity in *Escherichia coli*," *Nucleic Acids Research* 39(21):9275-9282.
- Somkuti, G. A. et al. (Apr. 1988). "Genetic Transformation of *Streptococcus thermophilus* by Electroporation," *Biochimie* 70(4):579-585. Abstract Only.
- Sorg, R.A. et al. (Mar. 20, 2014, e-pub. Jun. 4, 2014). "Gene Expression Platform for Synthetic Biology in the Human Pathogen *Streptococcus pneumoniae*," *ACS Synthetic Biology* 4(3):228-239, 38 pages.
- Suvorov, A. (1988). "Transformation of Group a *Streptococci* by Electroporation," *FEMS Microbiology Letters* 56(1):95-100.
- Turnbaugh, P.J. et al. (Dec. 2006). "An Obesity-Associated Gut Microbiome With Increased Capacity for Energy Harvest," *Nature* 444:1027-1131.
- USPTO Interference 106,123—Declaration to Declare Interference Jun. 11, 2020, 11 pages.
- USPTO Interference 106,123—Junior Party Annotated Claims Jul. 9, 2020, 31 pages.
- USPTO Interference 106,123—Junior Party List of Motions Jul. 16, 2020, 6 pages.
- USPTO Interference 106,123—Redeclaration Jul. 21, 2020, 6 pages.
- USPTO Interference 106,123—Rockefeller Clean Claims Jun. 25, 2020, 7 pages.
- USPTO Interference 106,123—Rockefeller Notice of Lead and Backup Counsel Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Notice of Real Party in Interest Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Notice of Related Proceedings Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Power of Attorney Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Request for File Copies Jun. 25, 2020, 10 pages.
- USPTO Interference 106,123—Rockefeller Revised List of Proposed Motions Aug. 13, 2020, 4 pages.
- USPTO Interference 106,123—Senior Party List of Proposed Motions Jul. 16, 2020, 5 pages.
- USPTO Interference 106,123—SNIPR Clean Claims Jun. 25, 2020, 27 pages.
- USPTO Interference 106,123—SNIPR Notice of Lead and Backup Counsel Jun. 25, 2020, 4 pages.
- USPTO Interference 106,123—SNIPR Notice of Related Proceedings Jun. 25, 2020, 4 pages.
- USPTO Interference 106,123—SNIPR Real Party in Interest Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—SNIPR Request for File Copies Jun. 25, 2020, 10 pages.
- USPTO Interference 106,123—Standing Order Jun. 11, 2020, 81 pages.
- USPTO Interference 106,123—Junior Party Revised List of Motions Aug. 13, 2020, 6 pages.
- USPTO Interference 106,123—Order Authorizing Motions and Setting Times 37 C.F.R. 11.104(c) and 121 Aug. 24, 2020, 10 pages.
- USPTO Interference 106,123—Order—Bd.R. 109(b)—Authorizing Copies of Office Records Jul. 21, 2020, 3 pages.
- Wegmann, U. et al. (2007). "Complete Genome Sequence of the Prototype Lactic Acid Bacterium *Lactococcus lactis* Subsp. *Cremoris* MG1363," *Journal of Bacteriology*, 189(8):3256-3270.
- Xie, Z. et al. (2013, e-pub. Aug. 9, 2013). "Development of a Tunable Wide-Range Gene Induction System Useful for the Study of *Streptococcal* Toxin-Antitoxin Systems," *Applied and Environmental Microbiology* 79(20):6375-6384.
- Yang, Y. et al. (Jun. 5, 2014, e-pub. Apr. 13, 2014). "Focused Specificity of Intestinal Th17 Cells Towards Commensal Bacterial Antigens," *Nature* 510(7503):152-156, 29 pages.
- Yu, Z. et al. (Mar. 21, 2014). "Various Applications of TALEN- and CRISPR/Cas9-Mediated Homologous Recombination to Modify the *Drosophila* Genome," *Biology Open* 3(4):271-280.
- Zhang, R. et al. (2009, e-pub. Oct. 30, 2008). "DEG 5.0, A Database of Essential Genes in Both Prokaryotes and Eukaryotes," *Nucleic Acids Research* 37:D455-D458.
- Zhang, X.Z. (2011). "Simple, Fast and High-Efficiency Transformation System for Directed Evolution of Cellulase in *Bacillus Subtilis*," *Microbial Biotechnology* 4(1):98-105.
- USPTO Interference 106,123—Order Additional Applications 37 C.F.R. § 41.104(a), Sep. 3, 2020, 6 pages.
- Abernethy, J. K. et al. (Mar. 2015, e-pub. Jan. 14, 2015). "Thirty Day All-Cause Mortality in Patients With *Escherichia coli* Bacteraemia in England," *Clin. Microbial. Infect.* 21:251.e1-251.e8.
- Aklujkar et al. (2010) "Interference With Histidyl-tRNA Synthetase by a CRISPR Spacer Sequence as a Factor in the Evolution of *Pelobacter Carbinolicus*," *BMC Evolutionary Biology* 10:203, 15 pages.
- American Lung Association (2019). "Preventing COPD," retrieved from <https://www.lung.org/lung-health-and-diseases/lung-disease-lookup/copd/symptoms-causes-risk-factors/preventing-copd.html>, last visited Aug. 5, 2019, 1 page.
- Anatoliotaki, M. et al. (2004). "Bloodstream Infections in Patients with Solid Tumors: Associated Factors, Microbial Spectrum and Outcome," *Infection* 2004, 32(2):65-71.
- Ang, Y.L.E. et al. (2015). "Best Practice in the Treatment of Advanced Squamous Cell Lung Cancer," *Ther. Adv. Respir. Dis.* 9(5):224-235.
- Anonymous (Apr. 2016). "Checkpoint Inhibition: A Promising Immunotherapeutic Approach for Colorectal Cancer," *Oncology*, 5(3):1-5, retrieved from <http://www.personalizedmedonc.com/publications/pmo/april-2016-vol-5-no-3/checkpoint-inhibition-a-promising-immunotherapeutic-approach-for-colorectal-cancer-2/>, last visited Aug. 27, 2019, 5 pages.
- Arnold, I.C. et al. (Apr. 8, 2015, e-pub. Mar. 4, 2015). "Helicobacter Hepaticus Infection in BALB/c Mice Abolishes Subunit-Vaccine-Induced Protection Against *M. Tuberculosis*," *Vaccine* 33(15):1808-1814.
- Arslan, Z. et al. (May 7, 2013). "RcsB-BglJ-Mediated Activation of Cascade Operon Does Not Induce the Maturation of CRISPR RNAs in *E. coli* K12," *RNA Biology* 10(5):708-715.
- Arumugam et al. (May 12, 2011). "Enterotypes of the human gut microbiome," *Nature* 473(7346):174-180, 16 pages.
- Beisel, C.L. et al. (2014). "A CRISPR Design for Next-Generation Antimicrobials," *Genome Biology* 15:516, 4 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Belizario, J.E. et al. (Oct. 6, 2015). "Human Microbiomes and Their Roles in Dysbiosis, Common Diseases, and Novel Therapeutic Approaches," *Frontiers in Microbiology* 6(1050):1-16.
- Bikard, D. et al. (2013, e-pub. Jun. 12, 2013). "Programmable Repression and Activation of Bacterial Gene Expression Using an Engineered CRISPR-Cas System," *Nucleic Acids Research* 41(15):7429-7437.
- Bikard, D. et al. (2017, e-pub. Sep. 6, 2017). "Using CRISPR-Cas Systems as Antimicrobials," *Current Opinion in Microbiology* 37:155-160.
- Bikard, D. et al. (Aug. 16, 2012). "CRISPR Interference Can Prevent Natural Transformation and Virulence Acquisition during In Vivo Bacterial Infection," *Cell Host & Microbe* 12(2):177-186.
- Bikard, D. et al. (Nov. 2014). "Development of Sequence-Specific Antimicrobials Based on Programmable CRISPR-Cas Nucleases," *Nature Biotechnology* 32(11):1146-1151, 16 pages.
- Broaders, E. et al. (Jul./Aug. 2013). "Mobile Genetic Elements of the Human Gastrointestinal Tract," *Gut Microbes* 4(4):271-280.
- Brouns, S.J.J. et al. (Aug. 15, 2008). Supplemental Material for "Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes," *Science* 321:960-964.
- Brouns, S.J.J. et al. (Aug. 15, 2008). "Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes," *Science* 321:960-964.
- Bugrysheva, J.V. et al. (Jul. 2011, E-Pub. Apr. 29, 2011). "The Histone-Like Protein HIOP is Essential for Growth of *Streptococcus pyogenes*: Comparison of Genetic Approaches to Study Essential Genes," *Appl. Environ. Microbiol.* 77(13):4422-4428.
- Bullman, S. et al. (Nov. 23, 2017). "Analysis of Fusobacterium Persistence and Antibiotic Response in Colorectal Cancer," *Science* pp. 1443-1448, 10 pages.
- Chan, B.K. et al. (2013). "Phage Cocktails and the Future of Phage Therapy," *Future Microbiol.* 8(6):769-783.
- Citorik, R.J. et al. (Nov. 2014, e-pub Sep. 21, 2014). "Sequence-Specific Antimicrobials Using Efficiently Delivered RNA-Guided Nucleases," *Nat. Biotechnol.* 32(11):1141-1145, 18 pages.
- Cochrane, K. et al. (2016, e-pub. Nov. 3, 2015). "Complete Genome Sequences and Analysis of the *Fusobacterium nucleatum* Subspecies *Animalis* 7-1 Bacteriophage Φ funu1 and Φ funu2," *Anaerobe* 38:125-129. Abstract Only.
- Consumer Updates (2019). "Combating Antibiotic Resistance," retrieved from <https://www.fda.gov/ForConsumers/ConsumerUpdates/ucm092810.htm>, last visited Jan. 28, 2019.
- Coyne, M.J. et al. (2014). "Evidence of Extensive DNA Transfer between Bacteroidales Species Within the Human Gut," *mBio* 5(3):e01305-14, 12 pages.
- Dailere, R. et al. (Oct. 18, 2016). "Enterococcus hirae and Bacteriophage Intestinihominis Facilitate Cyclophosphamide-Induced Therapeutic Immunomodulatory Effects," *Immunity* 95:931-943.
- De Filippo, C. et al. (Aug. 33 2010). "Impact of Diet in Shaping Gut Microbiota Revealed by a Comparative Study in Children From Europe and Rural Africa," *Proc. Natl. Acad. Sci. USA* 107(33):14691-14696, 6 pages.
- De Paepe, M. et al. (Mar. 28, 2014). "Bacteriophages: An Underestimated Role in Human and Animal Health?" *Frontiers in Cellular and Infection Microbiology* 4(39):1-11.
- Deeks, E.D. (2014, e-pub. Jul. 15, 2014). "Nivolumab: A Review of its Use in Patients With Malignant Melanoma," *Drugs* 74:1233-1239.
- Deghorain, M. et al. (Nov. 23, 2012). "The *Staphylococci* Phages Family: An Overview," *Viruses* 4:3316-3335.
- Del Castillo, M. et al. (Dec. 1, 2016). The Spectrum of Serious Infections Among Patients Receiving Immune Checkpoint Blockade for the Treatment of Melanoma *Clin. Infect. Dis.* 63:1490-1493.
- Denham, J.D. et al. (2018). "Case Report: Treatment of Enteropathogenic *Escherichia coli* Diarrhea in Cancer Patients: A Series of Three Cases," *Case Reports in Infectious Diseases* Article ID 8438701:1-3.
- Derosa, L. et al. (2018, e-pub. Mar. 30, 2018). "Negative Association of Antibiotics on Clinical Activity of Immune Checkpoint Inhibitors in Patients With Advanced Renal Cell and Non-Small-Cell Lung Cancer," *Annals of Oncology*. 2 pages.
- Dhar, A.D. (Jul. 20, 2018). "Overview of Bacterial Skin Infections," Merck Manual retrieved from <https://www.merckmanuals.com/home/skin-disorders/bacterial-skin-infections/overview-of-bacterial-skin-infections>, last visited Jul. 20, 2018, 3 pages.
- Dickson, R.P. et al. (Jan./Feb. 2017). "Bacterial Topography of the Healthy Human Lower Respiratory Tract," *American Society for Microbiology* 8(1):e02287-6, 12 pages.
- Diez-Villasenor, C. et al. (May 2013). "CRISPR-Spacer Integration Reporter Plasmids Reveal Distinct Genuine Acquisition Specificities Among CROSPR-Cas 1-E Variants of *Escherichia coli*," *RNA Biology* 10(5):792-802.
- Dutilh, B.E. et al. (Jul. 24, 2014). "A Highly Abundant Bacteriophage Discovered in the Unknown Sequences of Human Faecal Metagenomes," *Nature Communications* 5(4498):1-10.
- Edgar et al. (Dec. 2010). "The *Escherichia coli* CRISPR System Protects From λ Lysogenization, Lysogens, and Prophage Induction," *Journal of Bacteriology* 192(23):6291-6294.
- Ex Parte Re-Exam, mailed Dec. 10, 2018, for U.S. Appl. No. 90/014,184, filed Aug. 10, 2018, for U.S. Pat. No. 9,701,964 102 pages.
- Fact Sheet (Oct. 2010). "Antimicrobial Resistance," National Institutes of Health, 1-2.
- Foca, A. et al. (2015, e-pub. Apr. 7, 2015). Gut Inflammation and Immunity: What is the Role of the Human Gut Virome? *Mediators of Inflammation* 2015(326032):1-7.
- Fujita, K. et al. (2017). "Emerging Concern of Infectious Diseases in Lung Cancer Patients Receiving Immune Checkpoint Inhibitor Therapy," *Eur. Resp. J.* 50, OA1478. (Abstract Only).
- Galperin, M.Y. (Dec. 2013). "Genome Diversity of Spore-Forming Firmicutes," *Microbiology Spectrum* 1(2): TBS-0015-2012, 27 pages.
- Garon, E.B. et al. (Oct. 2015). "Current Perspectives in Immunotherapy for Non-Small Cell Lung Cancer," *Seminars in Oncology* 42(5 Supp. 2):S11-S18.
- Geller, L.T. et al. (Sep. 15, 2017). "Potential Role of Intratumor Bacteria in Mediating Tumor Resistance to the Chemotherapeutic Drug Gemcitabine," *Cancer*, 1156-1160, 6 pages.
- Goldwater, P.N. et al. (2012). "Treatment of Enterohemorrhagic *Escherichia coli* (EHEC) Infection and Hemolytic Uremic Syndrome (HUS)," *BMC Medicine* 10:12, 8 pages.
- Golubovskaya, V. et al. (Mar. 15, 2016). "Different Subsets of T Cells, Memory, Effector Functions, and CAR-T Immunotherapy," *Cancers* 8(36), 12 pages.
- Gomaa et al. (Jan. 28, 2014). "Programmable Removal of Bacterial Strains by Use of Genome-Targeting CRISPR-Cas Systems," *mBio*, 5(1):e000928-13.
- Gomaa, A.A. et al. (Jan./Feb. 2014). Supplemental Material to "Programmable Removal of Bacterial Strains by Use of Genome-Targeting CRISPR-Cas Systems," *American Society for Microbiology* 5(1):1-9.
- Gopalakrishnan, V. et al. (Jan. 5, 2018). "Gut Microbiome Modulates Response to Anti-PD-1 Immunotherapy in Melanoma Patients," *Science* 359:97-103, 20 pages.
- Green, J. (Jul. 20, 2018). Colgate <https://www.colgate.com/en-us/oral-health/conditions/mouth-sores-and-infections/eight-common-oral-infections-0615>, last visited Jul. 20, 2018, 4 pages.
- Gudbergsdottir, S. et al. (2011, e-pub. Nov. 18, 2010). "Dynamic Properties of the *Sulfolobus* CRISPR/Cas and CRISPR/Cmr Systems When Challenged With Vector-Borne Viral and Plasmid Genes and Protospacers," *Molecular Microbiology* 79(1):35-49.
- Gudiol, C. et al. (2016). "Bloodstream Infections in Patients With Solid Tumors," *Virulence* 7(3):298-308.
- Guedan, S. et al. (Aug. 14, 2014). "ICOS-Based Chimeric Antigen Receptors Program Bipolar TH17/TH1 Cells," *Blood* 124(7):1070-1080.
- Ha, Y.E. et al. (2013). "Epidemiology and Clinical Outcomes of Bloodstream Infections Caused by Extended-Spectrum β -Lactamase-Producing *Escherichia coli* in Patients With Cancer," *Int. J. Antimicrob. Agen.* 42(5):403-409.
- Hargreaves, K.R. et al. (Aug. 26, 2014). "Abundant and Diverse Clustered Regularly Interspaced Short Palindromic Repeat Spacers

(56)

References Cited

OTHER PUBLICATIONS

in Clostridium difficile Strains and Prophages Target Multiple Phage Types within This Pathogen,” mBio 5(5):e01045-13.

Hartland, E.L. et al. (Apr. 30, 2013). “Enteropathogenic and Enterohemorrhagic *E. coli*: Ecology, Pathogenesis, and Evolution,” Frontiers in Cellular and Infection Microbiology 3(15):1-3.

Healthline (2019). “Cystic Fibrosis,” retrieved from <https://www.healthline.com/health/cystic-fibrosis#prevention>, last visited Aug. 5, 2019, 14 pages.

Hooper, L.V. et al. (Jun. 8, 2012). “Interactions Between the Microbiota and the Immune System,” Science 336(6086):1268-1273, 16 pages.

Huddleston, J.R. (Jun. 20, 2014). “Horizontal Gene Transfer in the Human Gastrointestinal Tract: Potential Spread of Antibiotic Resistance Genes,” Infection and Drug Resistance 7:167-176.

Hurwitz, A.A. et al. (Aug. 1998). “CTLA-4 Blockade Synergizes With Tumor-Derived Granulocyte—Macrophage Colony-Stimulating Factor for Treatment of an Experimental Mammary Carcinoma,” Proc. Natl. Acad. Sci. USA 95:10067-10071.

International Search Report and the Written Opinion of the International Searching Authority for PCT/EP2018/066954, dated Oct. 23, 2018, filed Jun. 25, 2018, 14 pages.

International Search Report and the Written Opinion of the International Searching Authority for PCT/EP2019/057453, dated Aug. 16, 2019, filed Mar. 25, 2019, 21 pages.

International Search Report for PCT/EP2016/059803, dated Jun. 30, 2016, filed May 3, 2016, 6 pages.

International Search Report for PCT/EP2018/082053, dated Mar. 14, 2019, filed Nov. 21, 2018, 9 pages.

Ivanov, I.I. et al. (May 2010). “Segmented Filamentous Bacteria Take the Stage,” Mucosal Immunol. 3(3):209-212, 7 pages.

Jiang, W. et al. (Nov. 2013). “Demonstration of CRISPR/Cas9/sgrRNA-Mediated Targeted Gene Modification in *Arabidopsis*, Tobacco, Sorghum and Rice,” Nucleic Acids Research 41(20):e188, 12 pages.

Jin, Y. et al. (2019, e-pub. Apr. 23, 2019). “The Diversity of Gut Microbiome is Associated With Favorable Responses to Anti-Programmed Death 1 Immunotherapy in Chinese Patients With NSCLC,” Journal of Thoracic Oncology 14(8):1378-1389.

Jinek et al. (Aug. 17, 2012). “A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity,” Science 337(6096):816-821.

Jones, R.B. et al. (2008). “Tim-3 Expression Defines a Novel Population of Dysfunctional T Cells With Highly Elevated Frequencies in Progressive HIV-1 Infection,” J. Exp. Med. 205(12):2763-2779.

Karch, H. et al. (Jul. 1999). “Epidemiology and diagnosis of Shiga toxin-producing *Escherichia coli* infections,” Diagnostic Microbiology and Infectious Disease 34(3):229-243.

Khoja, L. et al. (2015). “Pembrolizumab,” Journal for ImmunoTherapy of Cancer 3(36):1-13.

Kosiewicz, M.M. et al. (2014, e-pub. Mar. 26, 2014). “Relationship Between Gut Microbiota and Development of T Cell Associated Disease,” FEBS Lett. 588:4195-4206.

Krom, R.J. et al. (Jul. 5, 2015). “Engineered Phagemids for Nonlytic, Targeted Antibacterial Therapies,” Nano Letters 15(7):4808-4813.

Lopez-Sanchez, M.-J. et al. (2012, e-pub. Jul. 27, 2012). “The Highly Dynamic CRISPR1 System of *Streptococcus agalactiae* Controls the Diversity of its Mobilome,” Molecular Microbiology 85(6):1057-1071.

Luo, M.L. et al. (2015, e-pub. Oct. 17, 2014). “Repurposing Endogenous Type I CRISPR-Cas Systems for Programmable Gene Repression,” Nucleic Acids Research 43(1):674-681.

Macon, B.L. et al. (Jan. 2, 2018). “Acute Nephritis,” retrieved from healthline, <https://www.healthline.com/health/acute-nephritis-syndrome#types>, last visited Jul. 20, 2018, 13 pages.

Mahoney, K.M. et al. (2015). “The Next Immune-Checkpoint Inhibitors: PD-1/PD-L1 Blockade in Melanoma,” Clinical Therapeutics 37(4):764-782.

Mancha-Agresti, P. et al. (Mar. 2017). “A New Broad Range Plasmid for DNA Delivery in Eukaryotic Cells Using Lactic Acid

Bacteria: In Vitro and In Vivo Assays,” Molecular Therapy: Methods & Clinical Development 4:83-91.

Manica, A. et al. (2011, e-pub. Mar. 8, 2011). “In vivo Activity of CRISPR-Mediated Virus Defence in a Hyperthermophilic Archaeon,” Molecular Microbiology 80(2):481-491.

Marin, M. et al. (2014). “Bloodstream Infections in Patients With Solid Tumors Epidemiology, Antibiotic Therapy, and Outcomes in 528 Episodes in a Single Cancer Center,” Medicine 93:143-149.

Marraffini, L.A. et al. (Dec. 19, 2008). “CRISPR Interference Limits Horizontal Gene Transfer in *Staphylococci* by Targeting DNA,” Science 322(5909):1843-1845, 12 pages.

Matson, V. et al. (2018). “The Commensal Microbiome is Associated With Anti-PD-1 Efficacy in Metastatic Melanoma Patients,” Science 359(6371):104-108.

Matsushiro, A. et al. (Apr. 1999). “Induction of Prophages of Enterohemorrhagic *Escherichia coli* O157:117 With Norfloxacin,” J. Bacteriology 181(7):2257-2260.

Mayo Clinic (2019). “Pulmonary Embolism,” retrieved from <https://www.mayoclinic.org/diseases-conditions/pulmonary-embolism/synonyms-causes/syc-20354647>, last visited Aug. 5, 2019, 8 pages.

Mayo Clinic (2020). “Infectious Diseases,” retrieved from <https://www.mayoclinic.org/diseases-conditions/infectious-diseases/diagnosis-treatment/drc-20351179>, last visited Jan. 17, 2020, 5 pages.

Mayo Clinic (2020). “Malaria,” retrieved from <https://www.mayoclinic.org/diseases-conditions/malaria/diagnosis-treatment/drc-20351190>, last visited Jan. 17, 2020, 3 pages.

Mayo Clinic (2020). “Sexually Transmitted Diseases (STDs),” retrieved from <https://www.mayoclinic.org/diseases-conditions/sexually-transmitted-diseases/stds/diagnosis-treatment/drc-20351246>, last visited Jan. 17, 2020, 5 pages.

Mayo Clinic (Jul. 20, 2018). “Bacterial Vaginosis,” retrieved from <https://www.mayoclinic.org/diseases-conditions/bacterial-vaginosis/symptoms-causes/syc-20352279>, last visited Jul. 20, 2018, 3 pages.

Mayo Clinic (Jul. 20, 2018). “Cystitis,” retrieved from <https://www.mayoclinic.org/diseases-conditions/cystitis/symptoms-causes/syc-20371306>, last visited Jul. 20, 2018, 10 pages.

Mayo Clinic (Jul. 20, 2018). “Meningitis,” retrieved from <https://www.mayoclinic.org/diseases-conditions/meningitis/symptoms-causes/syc-20350508>, last visited Jul. 20, 2018, 6 pages.

Mayo Clinic (Jul. 20, 2018). “Pneumonia,” retrieved from <https://www.mayoclinic.org/diseases-conditions/pneumonia/symptoms-causes/syc-20354204>, last visited Jul. 20, 2018, 5 pages.

Mayo Clinic (Mar. 29, 2020). “Liver Disease,” retrieved from <https://www.mayoclinic.org/diseases-conditions/liver-problems/diagnosis-treatment/drc-20374507>, last visited Mar. 29, 2020, 8 pages.

Medina-Aparicio, L. et al. (May 2011, e-pub. Mar. 11, 2011). “The CRISPR/Cas Immune System is an Operon Regulated by LeuO, H-NS, and Leucine-Responsive Regulatory Protein in *Salmonella enterica* Serovar Typhi,” Journal of Bacteriology 193(10):2396-2407.

Mick, E. et al. (May 2013). “Holding a Grudge: Persisting Anti-Phage CRISPR Immunity in Multiple Human Gut Microbiomes,” RNA Biology 10(5):900-906.

Mills, S. et al. (Jan./Feb. 2013). “Movers and Shakers: Influence of Bacteriophages in Shaping the Mammalian Gut Microbiota,” Gut Microbes 4(1):4-16.

Nakamura, S. et al. (Nov. 2008). “Metagenomic Diagnosis of Bacterial Infections,” Emerging Infectious Diseases 14(11):1784-1786.

Nale, J.Y. et al. (2012). “Diverse Temperate Bacteriophage Carriage in Clostridium Difficile 027 Strains,” PLoS One 7(5):e37263, 9 pages.

Navarre, L. et al. (2007). “Silencing of Xenogenic DNA by H-NS—Facilitation of Lateral Gene Transfer in Bacteria by a Defense System That Recognizes Foreign DNA,” Genes & Development 21:1456-1471.

Nelson, M.H. et al. (2015). “Harnessing the Microbiome to Enhance Cancer Immunotherapy,” Journal of Immunology Research 2015: Article 368736, 12 pages.

News (May 22, 2018). “UK Government and Bill & Melinda Gates Foundation Join Carb-X Partnership in Fight Against Superbugs: Millions Earmarked to Boost Research into New Life-Saving Pro-

(56)

References Cited

OTHER PUBLICATIONS

- ucts to Address the Global Rise of Drug-Resistant Bacteria," Combating Antibiotic Resistant Bacteria, 7 pages.
- Norris, J.S. et al. (2000). "Prokaryotic Gene Therapy to Combat Multidrug Resistant Bacterial Infection," Gene Therapy 7:723-725. Notice of Intent to Issue Ex Parte Reexamination Certificate, dated Aug. 12, 2019, for U.S. Appl. No. 90/014,184, filed Aug. 10, 2018, 26 pages.
- Nowak, P. et al. (Nov. 28, 2015). "Gut Microbiota Diversity Predicts Immune Status in HIV-1 Infection," AIDS 29(18):2409-2418.
- Okazaki, T. et al. (2007). "PD-1 and PD-1 Ligands: From Discovery to Clinical Application," Intern. Immun. 19(7):813-824.
- Pardoll, D.M. (2012). "The Blockade of Immune Checkpoints in Cancer Immunotherapy," Nat. Rev. Cancer 12(4):252-264.
- Park, A. (Oct. 18, 2011). "A Surprising Link Between Bacteria and Colon Cancer," Cancer retrieved from <http://healthland.time.com/2011/10/18/a-surprising-link-between-bacteria-and-colon-cancer/>, last visited Aug. 27, 2019, 3 pages.
- Patterson, A.G. et al. (2017, e-pub. Mar. 27, 2017). "Regulation of CRISPR-Cas Adaptive Immune Systems," Current Opinion in Microbiology 37:1-7.
- Patterson, A.G. et al. (Dec. 15, 2016). "Quorum Sensing Controls Adaptive Immunity Through the Regulation of Multiple CRISPR-Cas Systems," Mol. Cell 64(6):1102-1108.
- Pawluk, A. et al. (Apr. 15, 2014). "A New Group of Phage Anti-CRISPR Genes Inhibits the Type I-E CRISPR-Cas System of *Pseudomonas aeruginosa*," mBio. 5(2):e00896.
- Pires, D.P. et al. (Sep. 2016, e-pub. Jun. 1, 2016). "Genetically Engineered Phages: A Review of Advances Over the Last Decade," Microbiology and Molecular Biology Reviews 80(3):523-543.
- Purdy, D. et al. (2002). "Conjugative Transfer of Clostridial Shuttle Vectors From *Escherichia coli* to *Clostridium difficile* Through Circumvention of the Restriction Barrier," Molec. Microbiology 46(2):439-452.
- Ramalingam, S.S. et al. (2014). "LB2-Metastatic Non-Small Cell Lung Cancer: Phase II Study of Nivolumab (Anti-PD-1, BMS-936558, ONO-4538) in Patients With Advanced, Refractory Squamous Non-Small Cell Lung Cancer," International Journal of Radiation Oncology Biology Physics Late Breaking Abstract (LB2).
- Rashid, T. et al. (2013). "The Role of *Klebsiella* in Crohn's Disease With a Potential for the Use of Antimicrobial Measures," International Journal of Rheumatology 2013(Article ID 610393):1-9.
- Request for Ex Parte Reexamination dated Aug. 10, 2018, for U.S. Appl. No. 15/160,405, now U.S. Pat. 9,701,964, 42 pages.
- Request for Ex Parte Reexamination dated Nov. 1, 2018, for U.S. Appl. No. 15/160,405, now U.S. Pat. No. 9,701,964, 35 pages.
- Richter, C. et al. (2012, e-pub. Oct. 19, 2012). "Function and Regulation of Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) / CRISPR Associated (Cas) Systems," Viruses 4(12):2291-2311.
- Ridaura, V.K. et al. (Sep. 6, 2013). "Cultured Gut Microbiota From Twins Discordant for Obesity Modulate Adiposity and Metabolic Phenotypes in Mice," Science 341(6150):1241214, 22 pages.
- Routy, B. et al. (Jan. 5, 2018, e-pub. Nov. 2, 2017). "Gut Microbiome Influences Efficacy of PD-1-Based Immunotherapy Against Epithelial Tumors," Science 359(6371):91-97.
- Roy, S. et al. (May 2017, e-pub. Mar. 17, 2017). "Microbiota: A Key Orchestrator of Cancer Therapy," Nat. Rev. Cancer 17(5):271-285.
- Safdar, N. et al. (Jun. 4, 2002). "The Commonality of Risk Factors for Nosocomial Colonization and Infection With Antimicrobial-Resistant *Staphylococcus aureus*, Enterococcus, Gram-Negative Bacilli, *Clostridium difficile*, and *Candida*," Ann. Intern. Med. 136(11):834-844.
- Samonis, G. et al. (Sep. 2013, e-pub. Apr. 27, 2013). "A Prospective Study of Characteristics and Outcomes of Bacteremia in Patients With Solid Organ or Hematologic Malignancies," Support Care Cancer 21(9):2521-2526.
- Seed et al. (Feb. 27, 2013). "A Bacteriophage Encodes its Own CRISPR/Cas Adaptive Response to Evade Host Innate Immunity," Nature 494(7438):489-491.
- Selle et al. (Apr. 1, 2015). "Harnessing CRISPR-Cas Systems for Bacterial Genome Editing," Trends in Microbiology 23(4):225-232.
- Sepsis Alliance, (Dec. 14, 2017). "What are Vaccines," Retrieved from <https://www.sepsis.org/sepsisand/prevention-vaccinations/>; last visited Jul. 8, 2019, 3 pages.
- Sepsis Alliance; (Jul. 8, 2019). "Prevention," Retrieved from <https://www.sepsis.org/sepsisand/prevention/>; accessed last visited Jul. 8, 2019, 5 pages.
- Sharan, S.K. et al. (2009). "Recombineering: A Homologous Recombination-Based Method of Genetic Engineering," Nat. Protoc. 4(2):206-223, 37 pages.
- Shoemaker, N.B. et al. (Feb. 2001). "Evidence for Extensive Resistance Gene Transfer Among *Bacteroides* spp. and Among *Bacteroides* and Other Genera in the Human Colon," Appl. Environ. Microbiol. 67(2):561-68.
- Sivan, A. et al. (Nov. 27, 2015, e-pub Nov. 5, 2015). "Commensal Bifidobacterium Promotes Antitumor Immunity and Facilitates Anti-PD-L1 Efficacy," Science 350(6264):1084-1089, 13 pages.
- Skenneron, C.T. et al. (May 2011). "Phage Encoded H-NS: A Potential Achilles Heel in the Bacterial Defence System," PLoS ONE 6(5):e20095.
- Slutsker, L. et al. (Apr. 1998). "A Nationwide Case-Control Study of *Escherichia coli* O157:H7 Infection in the United States," J. Infect. Dis. 177(4):962-966.
- Soutourina, O.A. et al. (May 9, 2013). "Genome-Wide Identification of Regulatory RNAs in the Human Pathogen *Clostridium difficile*," PLoS Genet. 9(5):e1003493, 20 pages.
- Stern, A. et al. (2012). "CRISPR Targeting Reveals a Reservoir of Common Phages Associated With the Human Gut Microbiome," Genome Research 22(10):1985-1994.
- Stern, A. et al. (Aug. 2010). Self-Targeting by CRISPR: Gene Regulation or Autoimmunity? Trends Genet. 26(8):335-340, 10 pages.
- Stiefel, U. et al. (Aug. 2014, e-pub. May 27, 2014). "Gastrointestinal Colonization With a Cephalosporinase-Producing *Bacteroides* Species Preserves Colonization Resistance Against Vancomycin-Resistant Enterococcus and *Clostridium difficile* in Cephalosporin-Treated Mice," Antimicrob. Agents Chemother. 58(8):4535-4542.
- Stoebel, D.M. et al. (2008). "Anti-Silencing: Overcoming H-NS-Mediated Repression of Transcription in Gram-negative Enteric Bacteria," Microbiology 154:2533-2545.
- Takaishi, H. et al. (2008). "Imbalance in Intestinal Microflora Constitution Could be Involved in the Pathogenesis of Inflammatory Bowel Disease," Int. J. Med. Microbiol. 298:463-472.
- Takeda, T. et al. (2011). "Distribution of Genes Encoding Nucleoid-Associated Protein Homologs in Plasmids," International Journal of Evolutionary Biology 2011:685015, 31 pages.
- Tan, J. (Dec. 17, 2015). "Immunotherapy Meets Microbiota," Cell 163:1561.
- Tarr, P.I. et al. (Mar. 19-25, 2005). "Shiga-Toxin-Producing *Escherichia coli* and Haemolytic Uraemic Syndrome," Lancet 365(9464):1073-1086.
- Topalian, S.L. et al. (Jun. 28, 2012). "Safety, Activity, and Immune Correlates of Anti-PD-1 Antibody in Cancer," N. Engl. J. Med. 366(26):2443-2454, 19 pages.
- U.S. Appl. No. 16/201,736, filed Nov. 27, 2018, for Martinez et al.
- U.S. Appl. 16/682,889, filed Nov. 13, 2019, for Clube et al.
- U.S. Appl. 62/168,355, filed May 29, 2015, Barrangou, R. et al.
- Uchiyama, J. et al. (2013, e-pub. Mar. 8, 2013). "Characterization of *Helicobacter pylori* Bacteriophage KHP30," Applied and Environmental Microbiology 79(10):3176-3184.
- Veeranagouda, Y. et al. (Jun. 4, 2014). "Identification of Genes Required for the Survival of *B. fragilis* Using Massive Parallel Sequencing of a Saturated Transposon Mutant Library," BMC Genomics 15:429, 11 pages.
- Vega, N.M. et al. (Oct. 2014). "Collective Antibiotic Resistance: Mechanisms and Implications," Curr. Opin. Microbiol. 21:28-34, 14 pages.
- Velasco, E. et al. (2006). "Comparative Study of Clinical Characteristics of Neutropenic and Non-Neutropenic Adult Cancer Patients With Bloodstream Infections," Eur. J. Clin. Microbiol. Infect. Dis. 25:1-7.

(56)

References Cited

OTHER PUBLICATIONS

- Vercoe, R.B. et al. (Apr. 18, 2013). "Cytotoxic Chromosomal Targeting by CRISPR/Cas Systems Can Reshape Bacterial Genomes and Expel or Remodel Pathogenicity Islands," *PLOS Genetics* 9(4):e1003454, 13 pages.
- Villarino, N.F. et al. (Feb. 23, 2016, e-pub. Feb. 8, 2016). "Composition of the Gut Microbiota Modulates the Severity of Malaria," *Proc. Natl. Acad. Sci. USA* 113(8):2235-2240.
- Vétizou, M. et al. (Nov. 27, 2015, e-pub. Nov. 5, 2015). "Anticancer Immunotherapy by CTLA-4 Blockade Relies on the Gut Microbiota," *Science* 350(6264):1079-1084, 13 pages.
- Wagner, P.L. (2002). "Bacteriophage Control of Shiga Toxin 1 Production and Release by *Escherichia coli*," *Molecular Microbiology* 44(4):957-970.
- Walters, W.A. et al. (Nov. 17, 2014). "Meta-Analyses of Human Gut Microbes Associated With Obesity and IBD," *FEBS Letters* 588(22):4223-4233, 34 pages.
- Wei, Y. et al. (2015, e-pub. Jan. 14, 2015). "Sequences Spanning the Leader-Repeat Junction Mediate CRISPR Adaptation to Phage in *Streptococcus thermophilus*," *Nucleic Acids Research* 43(3):1749-1758.
- Westra, E.R. et al. (Sep. 1, 2010, e-pub. Aug. 18, 2010). "H-NS-Mediated Repression of CRISPR-Based Immunity in *Escherichia coli* K12 can be Relieved by the Transcription Activator LeuO," *Molecular Microbiology* 77(6):1380-1393.
- Westwater, C. et al. (Apr. 2003). "Use of Genetically Engineered Phage to Deliver Antimicrobial Agents to Bacteria: An Alternative Therapy for Treatment of Bacterial Infections," *Antimicrobial Agents and Chemotherapy* 47(4):1301-1307.
- Wexler, H.M. (Oct. 2007). "Bacteroides: the Good, the Bad, and the Nitty-Gritty," *Clinical Microbiology Reviews* 20(4):593-621.
- Wong, C.S. et al. (Jun. 29, 2000). "The Risk of the Hemolytic-Uremic Syndrome After Antibiotic Treatment of *Escherichia coli* O157:H7 Infections," *N. Engl. J. Med.* 342(26):1930-1936, 13 pages.
- Written Opinion for PCT Application No. PCT/EP2016/059803, dated Jun. 30, 2016, filed May 3, 2016, 6 pages.
- Written Opinion for PCT/EP2018/082053, dated Mar. 14, 2019, filed Nov. 21, 2018, 6 pages.
- Yao, J. et al. (2016, e-pub. May 9, 2016). "A Pathogen-Selective Antibiotic Minimizes Disturbance to the Microbiome," *Antimicrob. Agents Chemother.*, 24 pages.
- Yosef, I. et al. (2011). "High-Temperature Protein G is Essential for Activity of the *Escherichia coli* Clustered Regularly Interspaced Palindromic Repeats (CRISPR)/Cas System," *Proc. Natl. Acad. Sci. USA* 108(50):20136-20141.
- YourGenome: CRISPR/CAS9, retrieved from <https://www.yourgenome.org/facts/what-is-crispr-cas9>, last visited Jan. 6, 2020, 8 pages.
- Zembower, T.R. (2004). "Epidemiology of Infections in Cancer Patients," in *Infectious Complications in Cancer Patients*, Springer International Publishing Switzerland, 48 pages.
- Zhang, T. et al. (Sep. 24, 2016). "The Efficacy and Safety of Anti-PD-1/PD-L1 Antibodies for Treatment of Advanced or Refractory Cancers: A Meta-Analysis," *Oncotarget* 7(45):73068-73079.
- Zimmerhackl, L.B. (Jun. 29, 2000). "*E. coli*, Antibiotics, and the Hemolytic-Uremic Syndrome," *N. Engl. J. Med.* 342(26):1990-1991.
- Zitvogel, L. et al. (Jan. 2015), "Cancer and the Gut Microbiota: An Unexpected Link," *Sci. Transl. Med.* 7(271):271ps1, 10 pages.
- Zitvogel, L. et al. (Mar. 2018). "The Microbiome in Cancer Immunotherapy: Diagnostic Tools and Therapeutic Strategies," *Science* 359(6382):1366-1370.
- Bae, T. et al. (2006). "Prophages of *Staphylococcus aureus* Newman and Their Contribution to Virulence," *Molecular Microbiology* pp. 1-13.
- Cong, L. et al. (Feb. 15, 2013). "Multiplex Genome Engineering Using CRISPR/Cas Systems," *Science* 339(6121):819-823, 9 pages.
- Garneau, J. E. et al. (Nov. 4, 2010). "The CRISPR/Cas Bacterial Immune System Cleaves Bacteriophage and Plasmid DNA," *Nature* 468(7320):67-71, 28 pages.
- Jiang, W. et al. (Mar. 2013). "CRISPR-Assisted Editing of Bacterial Genomes," *Nat. Biotechnol.* 31(3):233-239.
- Mei, J.-M. et al. (1997). "Identification of *Staphylococcus aureus* Virulence Genes in a Murine Model of Bacteraemia Using Signature-Tagged Mutagenesis," *Molecular Microbiology* 26(2):399-407.
- USPTO Interference 106,123—Joint Stipulated Extension of Time, Sep. 4, 2002, 4 pages.
- Westwater, C. et al. (2002). "Development of a PI Phagemid System for the Delivery of DNA into Gram-Negative Bacteria," *Microbiology* 148:943-950.

* cited by examiner




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3




FIG. 4




FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 7




FIG. 8

Progression-free survival by fecal diversity:




FIG. 9A

Progression-free survival by low/high abundance of certain bacteria:




FIG. 9B

1

TREATING AND PREVENTING MICROBIAL INFECTIONS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 15/967,484, filed Apr. 30, 2018, the contents of which are incorporated herein by reference in their entirety.

SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

The content of the following submission on ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing (file name: 786212000301SEQLIST.txt, date recorded: Nov. 26, 2019, size: 21 KB).

TECHNICAL FIELD

The invention provides methods for treating or preventing microbial (eg, bacterial) infections and means for performing these methods. In particular, treatment of infections requiring rapid or durable therapy is made possible, such as for treating acute conditions such as septicemia, sepsis, SIRS or septic shock. The invention is also particularly useful, for example, for treatment of microbes for environmental, food and beverage use. The invention relates inter alia to methods of controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate or fluid in an industrial or domestic system.

The invention also useful for the treatment of pathogenic bacterial infections in subjects receiving a treatment for a disease or condition, such as a transplant or a treatment for cancer, a viral infection or an autoimmune disease.

BACKGROUND

Septicaemia is an acute and serious bloodstream infection. It is also known as bacteraemia, or blood poisoning. Septicaemia occurs when a bacterial infection elsewhere in the body, such as in the lungs or skin, enters the bloodstream. This is dangerous because the bacteria and their toxins can be carried through the bloodstream to a subject's entire body. Septicaemia can quickly become life-threatening. It must be rapidly treated, such as in a hospital. If it is left untreated, septicaemia can progress to sepsis.

Septicaemia and sepsis aren't the same. Sepsis is a serious complication of septicaemia. Sepsis is when inflammation throughout the body occurs. This inflammation can cause blood clots and block oxygen from reaching vital organs, resulting in organ failure. The US National Institutes of Health (NIH) estimates that over 1 million Americans get severe sepsis each year. Between 28 and 50 percent of these patients may die from the condition. When the inflammation occurs with extremely low blood pressure, it's called septic shock. Septic shock is fatal in many cases.

The increase in average age of the population, more people with chronic diseases, on immunosuppressive drugs, and increase in the number of invasive procedures being performed has led to an increased rate of sepsis. People over 65 years old, particularly those who have health issues, are even more susceptible to sepsis than any other group. According to a study published in 2006, while people aged

2

65 years and older make up about 12% of the American population, they make up 65% of sepsis cases in the hospitals.

Septicaemia is caused by an infection in a part of the body. This infection is typically acute. Many types of bacteria can lead to septicaemia. The exact source of the infection often can't be determined. The most common infections that lead to septicaemia are:

- urinary tract infections
- lung infections, such as pneumonia
- kidney infections
- infections in the abdominal area

Bacteria from these infections enter the bloodstream and multiply rapidly, causing acute infection and immediate symptoms.

People who are already in the hospital for something else, such as a surgery, are at a higher risk of developing septicaemia. Secondary infections can occur while in the hospital. These infections are often more dangerous because the bacteria may already be resistant to antibiotics.

There is a higher risk of developing septicaemia if the subject:

- has severe wounds or burns
- is very young or very old
- has a compromised immune system, which can occur from diseases such as HIV or leukaemia
- has a urinary or intravenous catheter
- is on mechanical ventilation
- is receiving medical treatments that weakens the immune system, such as chemotherapy or steroid injections

The symptoms of septicaemia usually start very quickly. Even in the first stages of the illness, a person can look very sick. They may follow an injury, surgery, or another localized (eg, confined to one location) infection, like pneumonia. The most common initial symptoms are:

- chills
- elevated body temperature (fever)
- very fast respiration
- rapid heart rate

More severe symptoms will begin to emerge as the septicaemia progresses without proper treatment. These include the following:

- confusion or inability to think clearly
- nausea and vomiting
- red dots that appear on the skin
- reduced urine volume
- inadequate blood flow (shock)

Septicaemia that has started to affect the organs or tissue function is an acute medical emergency. It must be rapidly treated at a hospital. Many people with septicaemia are admitted to a hospital's ICU for treatment and recovery. It is recommended to never take a "wait and see" approach or try to treat the problem at home. It is crucial to get to the hospital right away if the subject is showing signs of septicaemia.

Septicaemia has a number of serious complications. These complications may be fatal if left untreated or if treatment is delayed for too long.

Septic Shock

One complication of septicaemia is a serious drop in blood pressure. This is called septic shock. Toxins released by the bacteria in the bloodstream can cause extremely low blood flow, which may result in organ or tissue damage. Septic shock is an acute medical emergency. People with septic shock are usually cared for in a hospital's intensive care unit (ICU). The patient may need to be put on a ventilator, or breathing machine, if in septic shock.

Acute Respiratory Distress Syndrome (ARDS)

Another complication of septicemia is acute respiratory distress syndrome (ARDS). This is a life-threatening condition that prevents enough oxygen from reaching your lungs and blood. According to the National Heart, Lung, and Blood Institute (NHLBI), ARDS is fatal in about one-third of cases. It often results in some level of permanent lung damage. It can also damage the brain, which can lead to memory problems.

Sepsis

Sepsis occurs when the body has a strong immune response to the infection. This leads to widespread inflammation throughout the body. It is called severe sepsis if it leads to organ failure. People with chronic diseases, such as HIV or cancer, are at a higher risk of sepsis. This is because they have a weakened immune system and cannot fight off the infection on their own. Sepsis causes millions of deaths globally each year and is the most common cause of death in people who have been hospitalized. The worldwide incidence of sepsis is estimated to be 18 million cases per year. In the United States sepsis affects approximately 3 in 1,000 people, and severe sepsis contributes to more than 200,000 deaths per year. Sepsis occurs in 1-2% of all hospitalizations and accounts for as much as 25% of ICU bed utilization.

Early diagnosis is necessary to properly manage sepsis, as initiation of rapid therapy is key to reducing deaths from severe sepsis. Within the first three hours of suspected sepsis, diagnostic studies should include white blood cell counts, measuring serum lactate, and obtaining appropriate cultures before starting antibiotics, so long as this does not delay their use by more than 45 minutes.

The most common primary sources of infection resulting in sepsis are the lungs, the abdomen, and the urinary tract. Typically, 50% of all sepsis cases start as an infection in the lungs.

Speed of treatment is essential. Two sets of blood cultures (aerobic and anaerobic) should be taken without delaying the initiation of antibiotics. Cultures from other sites such as respiratory secretions, urine, wounds, cerebrospinal fluid, and catheter insertion sites (in-situ more than 48 hours) can be taken if infections from these sites are suspected. In severe sepsis and septic shock, broad-spectrum antibiotics (usually two, a (3-lactam antibiotic with broad coverage, or broad-spectrum carbapenem combined with fluoroquinolones, macrolides, or aminoglycosides) are conventional. However, combination of antibiotics is not recommended for the treatment of sepsis without shock and in immunocompromised persons unless the combination is used to broaden the anti-bacterial activity. The administration of antibiotics is important in determining the survival of the person. Some recommend they be given within one hour of making the diagnosis, stating that for every hour of delay in the administration of antibiotics, there is an associated 6% rise in mortality.

Early goal directed therapy (EGDT) is an approach to the management of severe sepsis during the initial 6 hours after diagnosis. It is a step-wise approach, with the physiologic goal of optimizing cardiac preload, afterload, and contractility. It includes giving early antibiotics.

Neonatal sepsis can be difficult to diagnose as newborns may be asymptomatic. If a newborn shows signs and symptoms suggestive of sepsis, antibiotics are immediately started and are either changed to target a specific organism identified by diagnostic testing or discontinued after an infectious cause for the symptoms has been ruled out.

Approximately 20-35% of people with severe sepsis and 30-70% of people with septic shock die. The Surviving Sepsis Campaign (SSC) is a global initiative to bring together professional organizations in reducing mortality from sepsis. Antibiotics are administered within two hours of admission/diagnosis. For every hour a patient is denied antibiotic therapy after the onset of septic shock, the patient's chance of survival is reduced by 7.9% (Survivesepsis.org 2005)

There is, therefore, a need for a rapid treatment of acute microbial infections, such as bacterial infections associated with septicemia, sepsis or septic shock. It would also be advantageous if the treatment is durable for many hours. Rapid and durable treatment of microbes is also desirable for is for controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate in industrial and domestic systems.

Acute bacterial infections can, in certain circumstances, be health-threatening or even life-threatening. This may be the case, for example, in cancer patients, transplant patients or other subjects. The need for the treatment of the bacterial infection can become urgent and indeed an immediate focus of attention in the medical care. It would be useful to provide methods of treating such pathogenic bacterial infections in a way that does not adversely undermine the efficacy of the cancer or other separate therapy to which the patient also needs to respond.

SUMMARY OF THE INVENTION

The invention provides a solution by using the action of programmable nuclease cutting of microbe genomes; this is different from the metabolic inhibitor and other mechanisms of action used by beta-lactams and other conventional antibiotics for treating infections. The targeted cutting provides selective microbe killing or reduction of growth or proliferation to treat or prevent infection. Moreover, the inventors have surprisingly found a substantial killing (by several logs) can be achieved very rapidly (eg, within 15 minutes) and sustainable effects can be achieved (eg, for more than 1 hour, and even around 3 hours after treatment commenced) in some embodiments. Thus, the invention provides the following configurations.

In a First Configuration

A programmable nuclease for use in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

In a Second Configuration

A plurality of viruses (eg, phage or phagemids for producing phage) for use with a programmable nuclease in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and viruses wherein the nuclease is

5

programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated;

wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject;

wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

In a Third Configuration

A composition comprising a plurality of nucleic acids for programming a programmable nuclease in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and the nucleic acids wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated;

wherein each nucleic acid encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

In a Fourth Configuration

A CRISPR/Cas system comprising a nuclease according to the invention for use in the method of treatment, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of the microbes.

In a Fifth Configuration

A method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

In a Sixth Configuration

A method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a plurality of viruses wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated; wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

6

In a Seventh Configuration

A method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a plurality of nucleic acids wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated; wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein each nucleic acid encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

In a Eighth Configuration

Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of the invention, in the manufacture of a composition for carrying out a method of treatment as defined herein, wherein the subject is an organism other than a human or animal.

In a Ninth Configuration

Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of the invention, in the manufacture of a composition for carrying out an ex vivo or in vitro a method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.

In a Tenth Configuration

Use of a programmable nuclease in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.

In any Configuration:

For example, the infection is an acute infection. For example, the infection is an acute infection that is rapidly treated. For example, the infection is treated rapidly—for example, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. For example, the treatment is durable—for example, the reduction in infection persists for at least 30 minutes immediately after the first 30 minutes of the treatment. Also, optionally a reduction of the infection by at least 100-fold or 1000-fold is maintained for at least 60

minutes (eg, at least 120 minutes) after commencement of the treatment. Exemplification is provided below which surprisingly demonstrates these, such as a rapid killing that was durable around 3 hours after treatment commenced. For example, the method improves survival of the subject, or improves survival rates in humans or human patients suffering from infection by the microbes of the first species or strain.

The invention also provides a solution to the need for effective treatment of pathogenic bacterial infections in subjects undergoing a cancer or other, separate therapy which must also be efficacious. Thus, the invention further provides: —

In an Eleventh Configuration

A method for treating a pathogenic bacterial infection in a human or animal subject caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a programmable nuclease that is programmed to cut the target site, wherein the subject is suffering from a further disease or condition other than the pathogenic bacterial infection and the method comprises administering a therapy to the subject for treating or preventing the further disease or condition, wherein the nuclease treats the infection and the therapy is efficacious in the presence of the programmed nuclease to treat or prevent the disease or condition.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) that are *E. coli*, *Pseudomonas aeruginosa* or *Klebsiella* bacteria, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-PD-1/PD-L1 axis antibody optionally selected from pembrolizumab, nivolumab, atezolizumab, avelumab and durvalumab; and
- (b) The cancer is selected from melanoma; renal cell carcinoma; bladder cancer; a solid tumour; non-small cell lung cancer (NSCLC); forehead and neck squamous cell carcinoma (HNSCC); Hodgkin's lymphoma; a cancer that overexpresses PD-L1 and no mutations in EGFR or in ALK; colorectal cancer and hepatocellular carcinoma.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a programmable nuclease that is programmed to cut the target site, wherein the subject is suffering from a cancer and the method comprises administering a cancer therapy to the subject for treating the cancer, wherein the nuclease treats the infection and the cancer therapy is efficacious in the presence of the programmed nuclease to treat the cancer.

In a Twelfth Configuration

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first

species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer.

In a Thirteenth Configuration

A programmable nuclease for use in the method of the invention.

In a Fourteenth Configuration

A CRISPR/Cas system comprising a nuclease according to the 13th Configuration for use in the method of the 11th or 12th Configuration, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs (gRNAs) or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of first bacteria.

In a fifteenth Configuration

A guide RNA or a DNA encoding a guide RNA for use in the system or method of treating a pathogenic bacterial infection.

In a Sixteenth Configuration

A nucleic acid vector comprising the guide RNA or DNA.

In a Seventeenth Configuration

A pharmaceutical composition comprising a first nucleic acid vector (or a plurality thereof) encoding the nuclease and a second nucleic acid vector (or a plurality thereof) encoding the guide RNA.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B show time-kill curves for *Escherichia coli* (EHEC) ATCC43888 strain harboring the CGV system. FIG. 1A shows that CRISPR induction killed 99.98% of the population in 30 minutes (black line). Growth in absence of induction is shown in dashed lines. CRISPR was induced at time-point 0 and monitored until 60 minutes. FIG. 2B shows dilution series (10^1 - 10^6) of drop spots (5 μ l) on LB agar plates of *E. coli* ATCC43888 harboring the CGV system after 30 minutes of induction.

FIG. 2 shows CRISPR killing of target strain *Escherichia coli* (EHEC) ATCC43888 in *Galleria mellonella* larvae. *G. mellonella* larvae were delivered injections of bacteria behind the final left proleg. Approximately 1 h after the injection, CRISPR inducers were administered behind the final right proleg. Larvae were incubated at 37° C. for 2 h and sacrificed. Control bacteria carrying an off-target single guide RNA plasmid were also injected in the control group.

FIG. 2 shows CRISPR-kill curves of *Escherichia coli* (EHEC) ATCC43888 in *Galleria mellonella*. *G. mellonella* larvae were delivered injections of bacteria behind the final left proleg. Approximately 1 h after the injection, CRISPR inducers were administered behind the final right proleg. Larvae were incubated at 37° C. and sacrificed at 0, 1, and 2 h after induction.

FIG. 3 shows *E. coli* ATCC43888 count over time.

FIG. 4 shows Kaplan-Meier survival curves of *Galleria mellonella* larvae infected with *Escherichia coli* (EHEC) ATCC43888. CRISPR induction significantly improves survival of the larvae (black line) compared to the off-target control carrying an off-target single guide RNA plasmid (dashed line).

FIGS. 5A-5B show time-kill curves for *Escherichia coli* Nissle 1917 harboring the CGV system targeting pks. FIG. 5A shows that CRISPR induction killed 99.98% of the population in 15 minutes (black line). Growth in absence of induction is shown in dashed lines. CRISPR was induced at time-point 0 and monitored over 3 h. FIG. 5B shows dilution series (10^1 - 10^6) of drop spots (5 μ l) on LB agar plates of *E. coli* Nissle 1917 harboring the CGV system after 15 minutes of induction.

FIGS. 6A-6B show time-kill curves for *Escherichia coli* Nissle 1917 harboring the CGV system targeting yapH. FIG. 6A shows that CRISPR induction killed 99.98% of the population in 15 minutes (black line). Growth in absence of induction is shown in dashed lines. CRISPR was induced at time-point 0 and monitored over 3 h. FIG. 6B shows dilution series (10^1 - 10^6) of drop spots (5 μ l) on LB agar plates of *E. coli* Nissle 1917 harboring the CGV system after 15 minutes of induction.

FIG. 7 shows complete killing of transconjugant *C. difficile*. The complete precision killing of *Clostridium difficile* using a gRNA-encoding CRISPR array that was delivered from a probiotic carrier bacterial species by conjugative plasmids as vectors is shown. A carrier bacterium (*E. coli* donor strain containing the vectors) was mated with *Clostridium difficile* which was killed upon delivery of the designed array. This harnessed the endogenous Cas3 machinery of *Clostridium difficile*. A 100% killing of *Clostridium difficile* cells was achieved and is shown in this figure.

FIG. 8 shows the antibiotic treatment during ICI therapy has fatal outcomes. Kaplan Meier curve for overall survival of a validation cohort from the Memorial Sloan Ketterin Cancer Center including n=239 advanced NSCLC patients treated with anti-PD-L1/anti-PD-1 mAb who received (ATB, n=68) or not (no ATB, n=171) antibiotics (ATB) two months before the injection of immune checkpoint blockade. There was a median overall survival of 21.9 months in the absence of antibiotic treatment, compared to an overall survival of 9.8 months with antibiotic treatment. So, the median overall survival in patients treated with classical antibiotics is <50% (or >12 months shorter) that of patients not receiving antibiotic treatment.

FIGS. 9A and 9B show that the gut microbiome modulates the efficacy of anti-PD-1 inhibition in melanoma patients (from Gopalakrishnan et al, *Science* 2018, 359, 97-103).

DETAILED DESCRIPTION

The approach of the present invention is different from conventional antibiotic approaches.

The present invention utilizes targeted cutting of microbial genomes using programmed nucleases, whereas conventional antibiotics rely upon metabolic processes and cell replication cycles—and the inhibition of these—for their activity. By focusing instead on nuclease cutting, the invention surprisingly achieves very quick and efficient microbial killing that also is remarkably durable. This is demonstrated in experiments below with different microbes, different nucleases and different delivery approaches. Typically, 99-100% killing was surprisingly observed many times and killing of 3-4 logs was very quickly achieved and with lasting duration.

The invention provides methods for treating or preventing microbial (eg, bacterial) infections and means for performing these methods. In particular, treatment of infections requiring rapid therapy is made possible, such as for treating

acute conditions such as septicemia, sepsis, SIRS or septic shock. As explained herein, a rapid response is vital to address microbial infection in many settings. Speed is of the essence for many infection scenarios, such as acute infections requiring hospital admission. Benefits of the invention can be one or more of: the reduction in the spread, severity or progression of the infection in the subject; reduction in the development, severity or progression of symptoms of the infection (eg, sepsis or septic shock); and an increase in the likelihood of survival in human or animal patients.

The invention uses programmable nuclease cutting of microbe genomes. The targeted cutting provides selective microbe killing or reduction of growth or proliferation to treat or prevent infection, as opposed to more broad-spectrum microbial killing of several different species as seen with conventional antibiotics. Selective killing is advantageous to leave beneficial microbes untargeted by the treatment, which may be beneficial to the patient. Moreover, the inventors have surprisingly found a substantial (by several logs) killing can be achieved very rapidly (eg, within 15 minutes) and sustainable effects can be achieved (eg, for more than 1 hour) in some embodiments. As exemplified below, the inventors surprisingly could remarkably achieve a fast and durable killing for around 2-3 hours.

Thus, the invention provides the following aspects: —

A programmable nuclease for use in a method of treating a microbial infection (eg, an acute bacterial infection) of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

Another aspect provides: A programmable nuclease for use in a method of rapidly treating an acute microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is rapidly treated.

Another aspect provides: A programmable nuclease for use in a method of treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a nucleic acid that programs the nuclease to recognise and cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

Another aspect provides: A programmable nuclease for use in a method of rapidly treating an acute microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain

and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a nucleic acid that programs the nuclease to recognise and cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is rapidly treated.

Another aspect provides: A programmable nuclease for use in a method of durably treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are durably killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

Another aspect provides: A programmable nuclease for use in a method of durably treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are durably killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a nucleic acid that programs the nuclease to recognise and cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated.

Another aspect provides: A programmable nuclease for use in a method of durably treating an acute microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are durably killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a nucleic acid that programs the nuclease to recognise and cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated.

Surprisingly, as exemplified below, a durable effect of several logs (eg, 3 or 4 logs) using a nuclease (as opposed to conventional means for conventional antibiotic killing) was observed around 3 hours after the first exposure of bacteria with a programmed nuclease. This aspect of the invention, therefore, makes possible dosing regimens for less frequent exposure to a programmed nuclease (ie, less frequent administration of a programmed nuclease, programmable nuclease and/or nucleic acid for programming the nuclease). For example, a Cas and gRNA (or DNA encoding a gRNA) for programming the nuclease are administered with a programmable nuclease (eg, a Cas 9 or Cas3) to the subject at a first time (T1) and at a second time (T2); or gRNA (or DNA encoding a gRNA) is administered on T1 and T2 for programming an endogenous Cas nuclease (eg, a Cas9 or Cas3) of bacteria of said first species or strain, wherein the programmed endogenous Cas cuts the genomes of the bacteria to kill the bacteria or to reduce growth or

proliferation, thus treating the infection. Such less frequent dosing is convenient for the healthcare practitioner and patient, as well as provides for economical therapy. Thus, optionally, the nuclease and/or nucleic acid is administered to the subject on T1 and T2, wherein T2 is at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 24 hours after T1. For example, T2 is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 24 hours after T1. For example, T2 is 2-7 hours after T1. For example, T2 is 1 hour after T1. For example, T2 is 2 hours after T1. For example, T2 is 3 hours after T1. For example, T2 is 4 hours after T1. For example, T2 is 5 hours after T1.

Optionally, the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject on T1 and T2, wherein T2 is at least 1 hour (eg, 1, 1.5, 2, 2.5 or 3 hours) after T1.

Another aspect provides: A Cas nuclease for use in a method of treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable with a guide RNA (gRNA) to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising administering to the subject said a nucleic acid, wherein the nucleic acid is the gRNA or a DNA encoding the gRNA, thereby programming the nuclease to recognise and cut the target site of the microbes comprised by the subject, whereby genomes of the microbes are cut and microbial infection of the subject is treated, wherein the method comprises administering the nucleic acid to the subject on at a first time (T1) and at a second time (T2), whereby the subject is exposed to programmed nuclease on T1 and T2, and wherein T2 is no less than 1 hour after T1.

Optionally, T2 is no less than 2 hours after T1; optionally, T2 is no less than 3 hours after T1; optionally, T2 is no less than 4 hours after T1; optionally, T2 is no less than 5 hours after T1; optionally, T2 is no less than 6 hours after T1; optionally, T2 is no less than 7 hours after T1; optionally, T2 is no less than 8 hours after T1; optionally, T2 is no less than 9 hours after T1; optionally, T2 is no less than 10 hours after T1; optionally, T2 is no less than 11 hours after T1; optionally, T2 is no less than 12 hours after T1; optionally, T2 is no less than 13 hours after T1; optionally, T2 is no less than 14 hours after T1; or optionally, T2 is no less than 24 hours after T1. Additionally or alternatively: Optionally, T2 is no more than 7 hours after T1; optionally, T2 is no more than 12 hours after T1; optionally, T2 is no more than 24 hours after T1; optionally, T2 is 2-7 hours after T1; optionally, T2 is 24 hours after T1; optionally, T2 is 7 hours after T1; optionally, T2 is 6 hours after T1; optionally, T2 is 5 hours after T1; optionally, T2 is 4 hours after T1; optionally, T2 is 3 hours after T1; optionally, T2 is 2 hours after T1; optionally, T2 is 1 hour after T1. For example, T2 is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 24 hours after T1. For example, T2 is 1-7 hours after T1; or T2 is 2-7 hours after T1; or T2 is 3-7 hours after T1; or T2 is 4-7 hours after T1; or T2 is 5-7 hours after T1; or T2 is 6-7 hours after T1.

Optionally, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the method comprises reducing the infection at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the method comprises reducing the infection at least 10000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally, the method comprises reducing the infection such that the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection such that a reduction in infection by at least 100-fold persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection such that a reduction in infection by at least 1000-fold persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection such that a reduction in infection by at least 10000-fold persists for 30 minutes immediately after the first 30 minutes of the treatment.

Optionally, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein a reduction in infection by at least 100-fold persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein a reduction in infection by at least 1000-fold persists for 30 minutes immediately after the first 30 minutes of the treatment. Optionally, the method comprises reducing the infection at least 10000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein a reduction in infection by at least 10000-fold persists for 30 minutes immediately after the first 30 minutes of the treatment.

Optionally, the method comprises maintaining reduction of the infection by at least 100-fold for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, a reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, the method comprises maintaining reduction of the infection by at least 1000-fold for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, a reduction of the infection by at least 1000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, the method comprises maintaining reduction of the infection by at least 10000-fold for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, a reduction of the infection by at least 10000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.

Optionally, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, the method comprises reducing the infection at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 1000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease. Optionally, the method comprises reducing the infection at least 10000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 10000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.

Optionally, the method comprises reducing the infection at least 10000-fold by the first 15 minutes of the treatment; and wherein reduction of the infection by at least 10000-fold is maintained for at least 45 minutes after exposing the subject to the programmed nuclease. This is exemplified below.

In an example, the infection is durably treated, wherein a reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after commencement of the treatment. In an example, the infection is durably treated, wherein a reduction of the infection by at least 1000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after commencement of the treatment. In an example, the infection is durably treated, wherein a reduction of the infection by at least 10000-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after commencement of the treatment.

Optionally, the infection is reduced at least 100,000-fold by the first 30 or 45 minutes of the treatment. Optionally, the infection is reduced at least 100,000-fold by the first 30 or 45 minutes of the treatment and the reduction is maintained until the 60th minute of the treatment.

Optionally, the infection is reduced at least 1000,000-fold by the first 30 or 45 minutes of the treatment. Optionally, the infection is reduced at least 1000,000-fold by the first 30 or 45 minutes of the treatment and the reduction is maintained until the 60th minute of the treatment.

Optionally, the infection is reduced at least 100-fold by the first 15 minutes of the treatment.

Optionally, the infection is reduced at least 1000-fold by the first 15 minutes of the treatment.

Optionally, the infection is reduced at least 100-fold by the first 15 minutes of the treatment and at least 1000-fold by the first 30 minutes of the treatment.

For example, the reduction is maintained for at least 15 further minutes, eg, the infection is reduced at least 100-fold or at least 1000-fold by the first 15 minutes of the treatment and the reduction is maintained from the 15-30th minute or 15-45th minute of the treatment or 15-60th minute of the treatment.

For example, the infection is reduced at least 100-fold or at least 1000-fold or at least 10000-fold by the first 15 minutes of the treatment in the first 15 minutes and the reduction is maintained for from the 15-30th minute or 15-45th minute of the treatment.

Optionally, the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally the method comprises reducing the infection at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally the method comprises reducing the infection at least 10000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally, the method comprises reducing the infection such that the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment, eg, the reduction may persist for at least 60 minutes after the first 30 minutes of the treatment. If the treatment is administered at time zero (T₀), then the reduction in infection may be present at 60 minutes counted after T₀, and indeed may persist after that 60 minutes. In FIGS. 1A, 5A and 6A, for example, reduction is seen at 60-180 minutes after T₀. Optionally, the reduction in infection persists for at least 30 minutes after the first 30 minutes of the treatment.

In an example, the infection is reduced by at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99%, eg, in the first 15 minutes of treatment. In an example, the infection is reduced by at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99%, eg, in the first 30 minutes of treatment.

For determining killing or reduction in growth or proliferation of the target microbes, one can, for example, determine the difference in the number of microbes of the first species or strain in (i) a sample taken from the subject (eg, a blood, gut or leaf sample) immediately before commencement of the treatment and (ii) a sample (of the same type as the sample of (i), eg, a blood, gut or leaf sample respectively) taken from the subject at 30 minutes of the treatment. For example, if the microbes are bacteria, the samples may be assessed for the difference in colony forming units (CFU)/ml sample, eg, when the samples have been plated on agar in respective petri dishes and incubated under identical conditions. Another example may use microscopic counting of microbes in samples, or other routine methods known to the skilled addressee.

In an example, at least 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% killing of the microbes is achieved by the first 30, 60, 90 or 120 minutes (eg, by the first 30 minutes; or by the first 120 minutes) of the treatment. For example, wherein the subject is a human or animal, the killing is determined comparing the prevalence (eg, by standard colony counting on an agar plate) of the microbes (eg, bacteria) in a blood sample taken immediately before commencement of the treatment versus a sample taken after the first 15 or 30 minutes of the treatment. In an example, at least 80, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% killing of the microbes is achieved by the first 0.5, 1 or 2 hours of the treatment. In an example, at least 99% killing of the microbes is achieved by the first 30 minutes of the treatment. In an example, at least 99% killing of the microbes is achieved by the first 2 hours of the treatment. In an example, 100% killing is achieved. These are exemplified below. In an embodiment, less than 100% of the microbes are killed.

Worked examples of killing in bacteria are shown below. Surprisingly, using a programmed nuclease to target the bacteria of choice, specific cutting resulted in rapid killing of the target bacteria—at least 3 or 4 logs of killing (ie, 1000- or 10,000 fold killing) could be observed in very short spaces of time and surprisingly these were sustained for at least to an hour. Optionally, the infection is reduced at least 1000-fold by the first 15, 30 or 45 minutes of the treatment. Optionally, the infection is reduced at least 1000-fold by the first 15, 30 or 45 minutes of the treatment and the reduction is maintained until the 60th, 120th or 180th minute of the treatment. Optionally, the infection is reduced at least 10,000-fold by the first 15, 30 or 45 minutes of the treatment. Optionally, the infection is reduced at least 10,000-fold by the first 15, 30 or 45 minutes of the treatment and the reduction is maintained until the 60th, 120th or 180th minute of the treatment. See, for example, exemplification in FIG. 5A.

In an example, 100% killing is achieved by 24 hours after commencement of the treatment.

In an example, the infection is reduced at least 1000-fold for 2 hours or more (eg, for 2-3 hours). Optionally also the infection is reduced by at least 1000-fold by the first 15 or 13 minutes of the treatment.

In an example, the infection is reduced at least 10,000-fold for 2 hours or more (eg, for 2-3 hours). Optionally also the infection is reduced by at least 10,000-fold by the first 15 or 13 minutes of the treatment.

In an example, the infection is reduced by at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% for 1 hour; or for 1 hour or more; or for 2 hours or more (eg, for 2-3 hours). Optionally, the infection is reduced by at least 90% for 1 hour; or for 1 hour or more; or for 2 hours or more (eg, for 2-3 hours), and optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the infection is reduced by at least 90% for 1 hour or more, and by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the infection is reduced by least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% by the first 15 or 13 minutes of the treatment. Optionally, the infection is reduced by least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% by the first 15 or 13 minutes of the treatment; and wherein the reduction is maintained for 1 hour or more (eg, for 2 hours or more; or for 3 hours or more; or for about 2 hours; or for 2 hours; or for about 3 hours; or for 3 hours). Exemplification below is provided, wherein the bacteria are *E coli*.

Optionally, the subject is a human or animal and the microbes are bacteria (eg, *E coli* or *C difficile*), wherein blood infection of the subject by the bacteria is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the subject is a human or animal and the microbes are bacteria (eg, *E coli* or *C difficile*), wherein blood infection of the subject by the bacteria is reduced at least 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the subject is a human or animal and the microbes are bacteria (eg, *E coli* or *C difficile*), wherein blood infection of the subject by the bacteria is reduced at least 10,00-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the *E coli* are EHEC *E coli*.

Optionally, the programmed nuclease (eg, a Cas9 or Cas3) is capable of cutting a target site comprised by *E. coli* (EHEC) ATCC43888. Optionally, the programmed nuclease (eg, a Cas9 or Cas3) is capable of cutting a target site comprised by *E. coli* Nissle.

Optionally, the blood of the subject is infected with from 10⁷ to 10¹² CFU/ml (eg, from 10⁷ to 10¹¹, from 10⁷ to 10¹⁰, from 10⁷ to 10⁹ or from 10⁷ to 10⁸ CFU/ml) of the bacteria immediately before the treatment.

The worked example below shows improved survival using the method of the invention in an in vivo model. In an example, therefore, the method of the invention is for improving survival of the subject by treating acute microbial infection of a subject. In an example, the programmed nuclease herein is capable of carrying out the killing of bacteria of the first species or strain in *Galleria mellonella* larvae in vivo infection model.

The nuclease may be, for example, a DNase (eg, a Cpf1, Cas9 or Cas3) or a RNase (eg, Cas13b).

In an example, the nuclease is an isolated or recombinant nuclease. For example, the nuclease is a synthetic or non-naturally occurring nuclease.

In an example, the nuclease is ex vivo, eg, in vitro. In an example, the nucleic acid is ex vivo.

In an example, the guide RNA or DNA encoding guide RNA(s) herein is ex vivo, eg, in vitro.

Optionally, the nuclease is a Cas nuclease (eg, a Cpf1, CasX, CasY, Cas13b, Cas 3 or 9), a meganuclease, or a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease. In an example, the Cas is a *Streptococcus* (eg, *pyogenes* or *aureus*) Cas9, *Clostridium* (eg, *difficile*), *Salmonella* (eg, *typhimurium*) or *E coli* Cas3. For example, the Cas is a spCas. In an example, the Cas9 is in combination with a tracrRNA or a DNA encoding a tracrRNA which is

operable with the Cas. For example, the tracrRNA is of the same species as the Cas, eg, a *Spyogenes* tracrRNA or DNA encoding this.

In an example, the nuclease is a Cas 3 encoded by a nucleic acid comprising SEQ ID NO: 9 or a sequence that is at least 80, 85, 90, 95, 96, 97, 98 or 99% identical thereto. Optionally, also the bacteria are *Clostridium* (eg, *C difficile*) bacteria; or any *Clostridium* shown in Table 1. This is exemplified below.

In an example, the nuclease is a Cas 9 encoded by a nucleic acid comprising SEQ ID NO: 10 or a sequence that is at least 80, 85, 90, 95, 96, 97, 98 or 99% identical thereto. In an example, the nuclease is a Cas 9 comprising SEQ ID NO: 11 or a sequence that is at least 80, 85, 90, 95, 96, 97, 98 or 99% identical thereto. Optionally, also the bacteria are *Clostridium* (eg, *C difficile*) bacteria; or any *Clostridium* shown in Table 1. Optionally, also the bacteria are *E coli* (eg, EHEC). This is exemplified below.

Optionally, the method comprises administering to the subject a RNA or a nucleic acid (eg, DNA) that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

Optionally, the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject.

Optionally, subject comprises the nuclease prior to administration of the RNA or nucleic acid to the subject. For example, the nuclease is a Cas nuclease that is an endogenous Cas nuclease of bacterial cells of the first species or strain that are comprised by the subject. Thus, in this example, the RNA or nucleic acid may be administered to the subject and introduced into the bacteria for programming endogenous Cas comprised by the bacteria, thereby forming programmed Cas nuclease that cuts the target site in the genomes of the bacteria, whereby bacteria are killed or growth or proliferation of bacteria is reduced, thus treating or preventing the infection.

Optionally, a plurality of viruses (eg, phage or phagemids) are administered to the subject, wherein each virus comprises a copy (eg, one or more, eg, a plurality of copies) of the nucleic acid, wherein the viruses infect the microbes comprised by the subject to deliver thereto the nucleic acid. For example, viruses herein are phage or phagemids that infect (or are capable of infecting) the bacteria of the first species or strain.

Optionally, the ratio of administered viruses: microbes comprised by the subject is from 10 to 150. For example, the microbes are bacteria and the ratio is from 10 to 100, ie, a multiplicity of infection (MOI) of from 1 to 100 (eg, wherein the viruses are capable of replication, eg, are phage and not phagemid), eg, from 10 to 100. The ratio can be determined, for example, using a sample (eg, a blood or gut sample) from a human or animal subject immediately before the treatment and determining the number of microbes (eg, bacteria per ml of blood or gut sample). The amount of viruses to be administered can then be worked out according to the determination using the sample.

Optionally, the microbes are bacteria. Alternatively, the microbes are archaea. Alternatively, the microbes are viruses. Alternatively, the microbes are fungi. Alternatively, the microbes are algae. Alternatively, the microbes are protozoa.

In an example, the subject is a human and the infection is a nosocomial infection. In an example, the subject is a plant, yeast, protist or amoeba.

Optionally, the subject is a human (eg, an adult, child, neonate, toddler, teenager, male or female) or animal (eg, a

dog, cat, horse, cow, sheep, goat, salmon, chicken, turkey, pig, companion animal or livestock animal).

In an example, the subject is a human or animal and: Optionally, the infection is an infection of the lungs, abdomen or urinary tract. In an example, the subject is suffering from a urinary tract infection, lung infections, such as pneumonia, a kidney infection or an abdominal infection. In an example, the subject is a surgery patient. In an example, the subject is a burns patient. In an example, the subject has an infected wound (eg, a bacterially infected wound). In an example, the patient is suffering from AIDS or is infected by HIV. In an example, the subject is suffering from a cancer, such as a blood cancer, eg, leukaemia, eg, AML or CML or CLL or a lymphoma. In an example, the subject is a tissue or organ transplant patient, eg, a haematopoietic stem cell transplant or bone marrow transplant patient. In an example, the subject has a urinary or intravenous catheter. In an example, the subject is on mechanical ventilation. In an example, the subject has been receiving an immunosuppressant. In an example, the subject is suffering from pneumonia. In an example, the subject is an intensive care unit (ICU) patient. In an example, the subject is an Acute respiratory distress syndrome (ARDS) patient. In an example, the subject is suffering from meningitis, an infection in pregnancy, a ruptured gallbladder (a gallbladder rupture is a medical condition where the gallbladder leaks or bursts. Ruptures are commonly caused by inflammation of the gallbladder), abortion with septic shock (abortion with septic shock can be an acute life-threatening illness), endometritis (endometritis is an inflammatory condition of the lining of the uterus, usually due to an infection), Acute Respiratory Distress Syndrome (Acute respiratory distress syndrome is a lung condition; it occurs when fluid fills up the air sacs in the lungs) or cellulitis.

The increase in average age of the population, more people with chronic diseases, on immunosuppressive drugs, and increase in the number of invasive procedures being performed has led to an increased rate of sepsis. Optionally, the subject has undergone surgery, is on an immunosuppressant medication and/or is suffering from a chronic disease.

Optionally, the subject is a human over 60, 65, 70, 75 or 80 years of age or is a pediatric patient. In an alternative, the subject is a pediatric patient (eg, a human baby or child) or adolescent.

In an example, the method treats or prevents neonatal sepsis in the subject. In an example the subject is an immune-compromised human or animal, eg, suffering from an acute viral infection, such as HIV infection; or the subject is suffering from a cancer, eg, a blood cancer, such as a leukaemia; or the patient is a transplant patient, eg, that has received an organ, tissue or bone marrow transplant. In an example, the subject is a human or animal that is positive for gram negative bacterial lipopolysaccharide or lipid A. In an example, the subject is a human or animal that is positive for gram positive bacterial cell wall lipoteichoic acid.

Optionally, the method treats or prevents septicemia and/or sepsis (eg, septic shock) in the subject.

SIRS (Systemic Inflammatory Response Syndrome) criteria has been used to define sepsis.

SIRS is the presence of two or more of the following: abnormal body temperature, heart rate, respiratory rate, or blood gas, and white blood cell count. Sepsis is, for example, SIRS in response to an infectious process. Severe sepsis is, for example, sepsis with sepsis-induced organ dysfunction or tissue hypoperfusion (manifesting as hypotension, elevated lactate, or decreased urine output). Septic shock is,

for example, severe sepsis plus persistently low blood pressure, despite the administration of intravenous fluids.

In an embodiment, the method prevents or delays progression of end-organ dysfunction in the subject (when the subject is a human or animal).

Examples of end-organ dysfunction include the following:

Lungs: acute respiratory distress syndrome (ARDS) ($\text{PaO}_2/\text{FiO}_2 < 300$)

Brain: encephalopathy symptoms including agitation, confusion, coma; causes may include ischemia, bleeding, formation of blood clots in small blood vessels, microabscesses, multifocal necrotizing leukoencephalopathy

Liver: disruption of protein synthetic function manifests acutely as progressive disruption of blood clotting due to an inability to synthesize clotting factors and disruption of metabolic functions leads to impaired bilirubin metabolism, resulting in elevated unconjugated serum bilirubin levels

Kidney: low urine output or no urine output, electrolyte abnormalities, or volume overload

Heart: systolic and diastolic heart failure, likely due to chemical signals that depress myocyte function, cellular damage, manifest as a troponin leak (although not necessarily ischemic in nature)

More specific definitions of end-organ dysfunction exist for SIRS in pediatrics.

Cardiovascular dysfunction (after fluid resuscitation with at least 40 ml/kg of crystalloid)

hypotension with blood pressure <5th percentile for age or systolic blood pressure <2 standard deviations below normal for age, or

vasopressor requirement, or

two of the following criteria:

unexplained metabolic acidosis with base deficit >5 mEq/l

lactic acidosis: serum lactate 2 times the upper limit of normal

oliguria (urine output <0.5 ml/kg/h)

prolonged capillary refill ≥ 5 seconds

core to peripheral temperature difference $> 3^\circ \text{C}$.

Respiratory dysfunction (in the absence of cyanotic heart disease or known chronic lung disease)

the ratio of the arterial partial-pressure of oxygen to the fraction of oxygen in the gases inspired ($\text{PaO}_2/\text{FiO}_2$) <300 (the definition of acute lung injury), or

arterial partial-pressure of carbon dioxide (PaCO_2) >65 torr (20 mmHg) over baseline PaCO_2 (evidence of hypercapnic respiratory failure), or

supplemental oxygen requirement of greater than FiO_2 0.5 to maintain oxygen saturation $\geq 92\%$

Neurologic dysfunction

Glasgow Coma Score (GCS) ≤ 11 , or

altered mental status with drop in GCS of 3 or more points in a person with developmental delay/intellectual disability

Hematologic dysfunction

platelet count <80,000/ mm^3 or 50% drop from maximum in chronically thrombocytopenic, or

international normalized ratio (INR) >2

Disseminated intravascular coagulation

Kidney dysfunction

serum creatinine ≥ 2 times the upper limit of normal for age or 2-fold increase in baseline creatinine in people with chronic kidney disease

Liver dysfunction (only applicable to infants >1 month) total serum bilirubin ≥ 4 mg/dl, or alanine aminotransferase (ALT) ≥ 2 times the upper limit of normal

5 Table 2 sets out the criteria for a positive diagnosis of sepsis.

Optionally, the method reduces one or more symptoms in the patient selected from fever, low body temperature, rapid breathing, elevated heart rate, confusion, confusion, metabolic acidosis, respiratory alkalosis, low blood pressure, dysfunction of blood coagulation (such as blood clotting in one or more organs, or bruising) and oedema. Optionally, the method reduces septic shock. Optionally, the sepsis is severe sepsis.

15 Optionally, at the start of the treatment, the subject (eg, a human) has a temperature of $< 36^\circ \text{C}$. or $> 38^\circ \text{C}$.; a heart rate of $> 90/\text{min}$, a respiratory rate of > 20 breaths/min or $\text{PaCO}_2 < 4.3$ kPa; and white blood cell count of $< 4000/\text{mm}^3$ or $> 12,000/\text{mm}^3$.

20 Optionally, at the start of the treatment, the subject (eg, a human) has presence of two or more of the following: abnormal body temperature, abnormal heart rate, abnormal respiratory rate, abnormal blood gas and abnormal white blood cell count.

25 Optionally, the subject is a plant. In an example, the subject is a protist, eg, amoeba. Optionally in this example, the microbes are viruses (eg, large or giant viruses, eg, Mimiviruses). The nuclease, for example, is a Cas and is programmable using a guide RNA delivered by a viroplasm that infects the virus microbes

30 In an example the microbes are yeast, eg, *Candida*.

Preferably, the microbes are bacteria. Optionally, the bacteria are gram positive bacteria. Optionally, the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonnorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E. coli* (eg, EHEC *E. coli*), *C. difficile*, *V. cholera*, *Staphylococcus* (eg, *S. aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).

40 Optionally, the first species is selected from the species in Table 1.

Optionally, the first species is enterohemorrhagic *E. coli* (EHEC), *E. coli* Serotype O157:H7 or Shiga-toxin producing *E. coli* (STEC). In an example, the bacteria are selected from

Shiga toxin-producing *E. coli* (STEC) (STEC may also be referred to as Verocytotoxin-producing *E. coli* (VTEC);

Enterohemorrhagic *E. coli* (EHEC) (this pathotype is the one most commonly heard about in the news in association with foodborne outbreaks);

Enterotoxigenic *E. coli* (ETEC);

Enteropathogenic *E. coli* (EPEC);

Enteroaggregative *E. coli* (EAEC);

55 Enteroinvasive *E. coli* (EIEC); and

Diffusely adherent *E. coli* (DAEC).

Enterohemorrhagic *Escherichia coli* (EHEC) serotype O157:H7 is a human pathogen responsible for outbreaks of bloody diarrhoea and haemolytic uremic syndrome (HUS) worldwide. Conventional antimicrobials trigger an SOS response in EHEC that promotes the release of the potent Shiga toxin that is responsible for much of the morbidity and mortality associated with EHEC infection. Cattle are a natural reservoir of EHEC, and approximately 75% of EHEC outbreaks are linked to the consumption of contaminated bovine-derived products. EHEC causes disease in humans but is asymptomatic in adult ruminants. Character-

istics of *E. coli* serotype O157:H7 (EHEC) infection includes abdominal cramps and bloody diarrhoea, as well as the life-threatening complication haemolytic uremic syndrome (HUS). Currently there is a need for a treatment for EHEC infections (Goldwater and Bettelheim, 2012). The use of conventional antibiotics exacerbates Shiga toxin-mediated cytotoxicity. In an epidemiology study conducted by the Centers for Disease Control and Prevention, patients treated with antibiotics for EHEC enteritis had a higher risk of developing HUS (Slutsker et al., 1998). Additional studies support the contraindication of antibiotics in EHEC infection; children on antibiotic therapy for hemorrhagic colitis associated with EHEC had an increased chance of developing HUS (Wong et al., 2000; Zimmerhackl, 2000; Safdar et al., 2002; Tarr et al., 2005). Conventional antibiotics promote Shiga toxin production by enhancing the replication and expression of stx genes that are encoded within a chromosomally integrated lambdoid prophage genome. The approach of the present invention relies on nuclease cutting. Stx induction also promotes phage-mediated lysis of the EHEC cell envelope, allowing for the release and dissemination of Shiga toxin into the environment (Karch et al., 1999; Matsushiro et al., 1999; Wagner et al., 2002). Thus, advantageously, the invention provides alternative means for treating EHEC in human and animal subjects. This is exemplified below with surprising results on the speed and duration of anti-EHEC action produced by nuclease action (as opposed to conventional antibiotic action).

In an example, the subject (eg, a human) is suffering from or at risk of haemolytic uremic syndrome (HUS), eg, the subject is suffering from an *E. coli* infection, such as an EHEC *E. coli* infection.

An aspect of the invention provides: A plurality of viruses (eg, phage or phagemids for producing phage) for use with the nuclease of the invention in the method of treatment, wherein each virus comprises a copy of a nucleic acid described herein, wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

A aspect of the invention provides: A plurality of viruses (eg, phage or phagemids for producing phage) for use with a programmable nuclease in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated;

wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject;

wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

Optionally, the method is for durable treatment, eg, as described herein; and/or optionally, the infection is acute infection.

Optionally, the method is for rapid treatment, eg, as described herein; and/or optionally, the infection is acute infection.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, the nucleic acid is according to any nucleic acid of the invention herein.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, the nucleic acid is according to any nucleic acid of the invention herein.

In an alternative, when the microbes are viruses, the plurality of viruses are phage that are capable of infecting host cells harbouring the microbes, wherein the nucleic acids are introduced into the host cells for expression therein of the RNA. The RNA complexes with the nuclease in the host cells to guide the nuclease to cut a target site of the microbes (ie, to cut viral RNA or DNA), thereby inactivating the viral microbes. For example, the microbes are viruses (eg, in amoeba; or in human or animal or plant cells) and the viruses of said plurality of viruses are capable of targeting the microbes, whereby the nuclease is programmed to cut the microbes (eg, in the amoeba or in said cells).

An aspect of the invention provides: A composition comprising a plurality of nucleic acids for programming the nuclease of the invention in the method of treatment, wherein each nucleic acid is a nucleic acid as defined herein.

An aspect of the invention provides: A composition comprising a plurality of nucleic acids for programming a programmable nuclease in a method of treating a microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and the nucleic acids wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is treated; wherein each nucleic acid encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

Optionally, the method is for durable treatment, eg, as described herein; and/or optionally, the infection is acute infection.

Optionally, the method is for rapid treatment, eg, as described herein; and/or optionally, the infection is acute infection.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, each nucleic acid is according to any nucleic acid of the invention herein.

Optionally, the composition is a pharmaceutical composition comprising the nucleic acids and a pharmaceutically acceptable diluent, carrier or excipient. Optionally, the composition is for oral, intravenous, pulmonary, rectal, topical, buccal, ocular, intranasal, or subcutaneous administration to a human or animal subject. Optionally, the composition is a herbicide or pesticide or insecticide or nematocide or aracnicide. Optionally, the composition is toxic to yeast. Optionally, the composition is toxic to giant viruses.

An aspect of the invention provides: A CRISPR/Cas system comprising a nuclease according to the invention for use in the method of treatment, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9 or any other Cas mentioned herein) and the system comprises one or more guide RNAs or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of the microbes.

In an example, each guide RNA mentioned herein is a single guide RNA (ie, a chimaeric guide RNA). In another example, each guide RNA comprises a crRNA that is hybridised to a tracrRNA.

In an example, a target site mentioned herein is comprised by an essential gene, virulence gene or antibiotic resistance gene of the bacteria. In an example, a target site mentioned herein is comprised by a multi-copy sequence (ie, a sequence that is present in more than one (eg, 2, 3, 4, 5, 6, 7, 8 or 9, or more) copies in each bacterial genome). For example, the target site is comprised by a ribosomal RNA gene. In an example, a target site mentioned herein is comprised by a ribosomal RNA gene (eg, a 23S ribosomal RNA gene), a yapH gene; or a pks gene; or homologue or orthologue thereof.

Optionally, each guide RNA herein is capable of hybridizing to a protospacer sequence comprising the target site, wherein the protospacer sequence is 15-45 nucleotides in length, eg, 15-25; 18-21; 20; or about 20 nucleotides in length. Optionally, each guide RNA herein comprises a spacer sequence that is 15-45 nucleotides in length, eg, 15-25; 18-21; 20; or about 20 nucleotides in length.

Optionally, each guide RNA herein is cognate to a 5'-NGG protospacer adjacent motif (PAM), eg, wherein the bacteria are *E. coli*. Optionally, each guide RNA herein is cognate to a 5'-CCA or 5'-CCT protospacer adjacent motif (PAM), eg, wherein the bacteria are *C. difficile*.

An aspect of the invention provides: A guide RNA or a DNA encoding a guide RNA for use in the system of the invention for use in the method of treating an acute microbial infection in the subject, eg, septicaemia or sepsis.

An aspect of the invention provides: A nucleic acid vector comprising the guide RNA or DNA.

Optionally, the vector is a phage, phagemid, viriophage, virus, plasmid (eg, conjugative plasmid) or transposon. The example below shows almost complete killing can be achieved using a conjugative plasmid as the vector. Thus, in an embodiment, each vector is a conjugative plasmid that is delivered from carrier bacteria eg, probiotic carrier bacteria for administration to the human or animal subject. In an example, the carrier bacteria are *Lactobacillus* (eg, *L. reuteri*) or *E. coli*. This is exemplified below and achieved complete (100%) killing.

An aspect of the invention provides: An anti-sepsis or anti-septicaemia composition for administration to a human or animal for treating sepsis or septicaemia, the composition comprising a plurality of vectors, wherein each vector a vector of the invention.

An aspect of the invention provides: A method of treating (eg, rapidly and/or durably treating) an acute microbial infection of a subject, wherein the method is as defined herein.

An aspect of the invention provides: A method of treating (eg, rapidly and/or durably treating) an acute microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated (eg, rapidly and/or durably treated).

An aspect of the invention provides: A method of treating (eg, rapidly and/or durably treating) an acute microbial

infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a plurality of viruses wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated (eg, rapidly and/or durably treated); wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, the nucleic acid is according to any nucleic acid of the invention herein.

An aspect of the invention provides: A method of treating (eg, rapidly and/or durably treating) an acute microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease and a plurality of nucleic acids wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated (eg, rapidly and/or durably treated); wherein each virus comprises a copy of a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject; wherein each nucleic acid encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.

Optionally, the nuclease is according to any nuclease of the invention herein. Optionally, each nucleic acid is according to any nucleic acid of the invention herein.

In an example, the invention is for medical or dental or ophthalmic use (eg, for treating or preventing an infection in an organism or limiting spread of the infection in an organism).

In an example, the invention is for cosmetic use (eg, use in a cosmetic product, eg, make-up), or for hygiene use (eg, use in a hygiene product, eg, soap).

In an example, the vectors and/or nuclease prior to administration to the subject are comprised by a composition which is as any of the following (host here refers to the microbes of the first species or strain): In an example, the composition is a medical, ophthalmic, dental or pharmaceutical composition (eg, comprised by an anti-host vaccine). In an example, the composition is an antimicrobial composition, eg, an antibiotic or antiviral, eg, a medicine, disinfectant or mouthwash. In an example, the composition is a cosmetic composition (eg, face or body make-up composition). In an example, the composition is a herbicide. In an example, the composition is a pesticide (eg, when the host is a *Bacillus* (eg, *thuringiensis*) host). In an example, the composition is a beverage (eg, beer, wine or alcoholic

beverage) additive. In an example, the composition is a food additive (eg, where the host is an *E coli*, *Salmonella*, *Listeria* or *Clostridium* (eg, botulinum) host). In an example, the composition is a water additive. In an example, the composition is a additive for aquatic animal environments (eg, in a fish tank). In an example, the composition is an oil or petrochemical industry composition or comprised in such a composition (eg, when the host is a sulphate-reducing bacterium, eg, a *Desulfovibrio* host). In an example, the composition is a oil or petrochemical additive. In an example, the composition is a chemical additive. In an example, the composition is a disinfectant (eg, for sterilizing equipment for human or animal use, eg, for surgical or medical use, or for baby feeding). In an example, the composition is a personal hygiene composition for human or animal use. In an example, the composition is a composition for environmental use, eg, for soil treatment or environmental decontamination (eg, from sewage, or from oil, a petrochemical or a chemical, eg, when the host is a sulphate-reducing bacterium, eg, a *Desulfovibrio* host). In an example, the composition is a plant growth stimulator. In an example, the composition is a composition for use in oil, petrochemical, metal or mineral extraction. In an example, the composition is a fabric treatment or additive. In an example, the composition is an animal hide, leather or suede treatment or additive. In an example, the composition is a dye additive. In an example, the composition is a beverage (eg, beer or wine) brewing or fermentation additive (eg, when the host is a *Lactobacillus* host). In an example, the composition is a paper additive. In an example, the composition is an ink additive. In an example, the composition is a glue additive. In an example, the composition is an anti-human or animal or plant parasitic composition. In an example, the composition is an air additive (eg, for air in or produced by air conditioning equipment, eg, where the host is a *Legionella* host). In an example, the composition is an anti-freeze additive (eg, where the host is a *Legionella* host). In an example, the composition is an eyewash or ophthalmic composition (eg, a contact lens fluid). In an example, the composition is comprised by a dairy food (eg, the composition is in or is a milk or milk product; eg, wherein the host is a *Lactobacillus*, *Streptococcus*, *Lactococcus* or *Listeria* host). In an example, the composition is or is comprised by a domestic or industrial cleaning product (eg, where the host is an *E coli*, *Salmonella*, *Listeria* or *Clostridium* (eg, botulinum) host). In an example, the composition is comprised by a fuel. In an example, the composition is comprised by a solvent (eg, other than water). In an example, the composition is a baking additive (eg, a food baking additive). In an example, the composition is a laboratory reagent (eg, for use in biotechnology or recombinant DNA or RNA technology). In an example, the composition is comprised by a fibre retting agent. In an example, the composition is for use in a vitamin synthesis process. In an example, the composition is an anti-crop or plant spoiling composition (eg, when the host is a saprotrophic bacterium). In an example, the composition is an anti-corrosion compound, eg, for preventing or reducing metal corrosion (eg, when the host is a sulphate-reducing bacterium, eg, a *Desulfovibrio* host, eg for use in reducing or preventing corrosion of oil extraction, treatment or containment equipment; metal extraction, treatment or containment equipment; or mineral extraction, treatment or containment equipment). In an example, the composition is an agricultural or farming composition or comprised in such a composition. In an example, the composition is a silage additive. The invention provides a CRISPR array, gRNA-encoding nucleotide sequence, vector or plurality of vectors

described herein for use in any of the compositions described in this paragraph or for use in any application described in this paragraph, eg, wherein the host cell is a bacterial or archaeal cell. The invention provides a method for any application described in this paragraph, wherein the method comprises combining a CRISPR array, gRNA-encoding nucleotide sequence, vector or plurality of the invention with a host cell (eg, bacterial or archaeal cell). In an embodiment, the host cell is not present in or on a human (or human embryo) or animal.

Any aspect of the present invention is, for example, for an industrial or domestic use, or is used in a method for such use. For example, it is for or used in agriculture, oil or petroleum industry, food or drink industry, clothing industry, packaging industry, electronics industry, computer industry, environmental industry, chemical industry, aerospace industry, automotive industry, biotechnology industry, medical industry, healthcare industry, dentistry industry, energy industry, consumer products industry, pharmaceutical industry, mining industry, cleaning industry, forestry industry, fishing industry, leisure industry, recycling industry, cosmetics industry, plastics industry, pulp or paper industry, textile industry, clothing industry, leather or suede or animal hide industry, tobacco industry or steel industry.

Host cells herein refers to the microbes of the first species or strain. Optionally, any host cell(s) herein is/are bacterial or archaeal cells. In an example, the cell(s) is/are in stationary phase. In an example, the cell(s) is/are in exponential phase. In an example, the cell(s) is/are in lag phase. In an example, the cell(s) is/are wild-type cells or naturally-occurring cells, eg, comprised by a naturally-occurring microbiome, eg, of a human, animal, plant, soil, water, sea, waterway or environment. In an example, the cell(s) is/are artificially genetically modified.

In an example, a plurality of vectors of the invention are introduced into a plurality of said host cells, wherein the host cells are comprised by a bacterial population, eg, ex vivo, in vivo or in vitro. In an example, the host cells are comprised by a microbiota population comprised by an organism or environment (eg, a waterway microbiota, water microbiota, human or animal gut microbiota, human or animal oral cavity microbiota, human or animal vaginal microbiota, human or animal skin or hair microbiota or human or animal armpit microbiota), the population comprising first bacteria that are symbiotic or commensal with the organism or environment and second bacteria comprising said host cells, wherein the host cells are detrimental (eg, pathogenic) to the organism or environment. In an embodiment, the population is ex vivo. In an example, the ratio of the first bacteria sub-population to the second bacteria sub-population is increased. In an example, the first bacteria are *Bacteroides* (eg, *B fragalis* and/or *B thetaiotamicron*) bacteria. Optionally, the *Bacteroides* comprises one, two, three or more *Bacteroides* species selected from *caecae*, *capillosus*, *cellulosilyticus*, *coprocola*, *coprophilus*, *coprosuis*, *distansis*, *dorei*, *eggerthii*, *faecis*, *fingoldii*, *fluxus*, *fragalis*, *intestinalis*, *melaninogenicus*, *nordii*, *oleiciplenus*, *oralis*, *ovatus*, *pectinophilus*, *plebeius*, *stercoris*, *thetaitaomicron*, *uniformis*, *vulgatus* and *xylanisolvans*. For example, the *Bacteroides* is or comprises *B thetaiotamicron*. For example, the *Bacteroides* is or comprises *B fragalis*.

In an example, the host, first or second cells are any bacterial species disclosed in US20160333348, GB1609811.3, PCT/EP2017/063593 and all US equivalent applications. The disclosures of these species (including specifically, Table 1 of PCT/EP2017/063593), are incorpo-

rated herein in their entirety and for potential inclusion of one or more disclosures therein in one or more claims herein.

In an example, the host cell(s) or bacterial population is harboured by a beverage or water (eg, a waterway or drinking water) for human consumption. In an example, the host cell(s) or said population is comprised by a composition (eg, a medicament (eg, bacterial gut transplant), beverage, mouthwash or foodstuff) for administration to a human or non-human animal for populating and rebalancing the gut or oral microbiota thereof (eg, wherein said use of the medicament is to treat or prevent a disease or condition in the human or animal). In an example, the host cell(s) or said population are on a solid surface or comprised by a biofilm (eg, a gut biofilm or a biofilm on an industrial apparatus). In an example of the invention for in vitro treating an industrial or medical fluid, solid surface, apparatus or container (eg, for food, consumer goods, cosmetics, personal healthcare product, petroleum or oil production); or for treating a waterway, water, a beverage, a foodstuff or a cosmetic, wherein the host cell(s) are comprised by or on the fluid, surface, apparatus, container, waterway, water, beverage, foodstuff or cosmetic.

In an example, the invention provides a container for medical or nutritional use, wherein the container comprises the vectors for use in the method. For example, the container is a sterilised container, eg, an inhaler or connected to a syringe or IV needle.

In an example, the vectors or composition is for administration (or is administered) to the human or non-human animal subject by mucosal, gut, oral, intranasal, intrarectal, intravaginal, ocular or buccal administration.

Optionally, each host cell is of a strain or species found in human microbiota, optionally wherein the host cells are mixed with cells of a different strain or species, wherein the different cells are Enterobacteriaceae or bacteria that are probiotic, commensal or symbiotic with humans (eg, in the human gut. In an example, the host cell is an *E coli* or *Salmonella* cell.

The invention is optionally for altering the relative ratio of sub-populations of first and second bacteria in a mixed population of bacteria, eg, for altering human or animal microbiomes, such as for the alteration of the proportion of Bacteroidetes (eg, *Bacteroides*, eg, *fragalis* and/or *thetaita-micron*), Firmicutes and/or gram positive or negative bacteria in microbiota of a human.

In an example, the vectors or composition of the invention comprises a nucleotide sequence for expressing in the host cell an endolysin for host cell lysis, optionally wherein the endolysin is a phage phil 1, phage Twort, phage P68, phage phiWMY or phage K endolysin (eg, MV-L endolysin or P-27/HP endolysin).

In an example, the target site is comprised by a chromosome of each microbe host cell, eg, wherein the sequence is comprised by an antibiotic resistance gene, virulence gene or essential gene of the host cell. An example, provides the vectors of the invention in combination with an antibiotic agent (eg, a beta-lactam antibiotic), eg, wherein the vectors target a protospacer sequence comprised by an antibiotic resistance gene comprised by host cell genome or episome (eg, a plasmid comprised by the host cells). In an example, the episome is a plasmid, transposon, mobile genetic element or viral sequence (eg, phage or prophage sequence).

In an example, the target is a chromosomal sequence, an endogenous host cell sequence, a wild-type host cell sequence, a non-viral chromosomal host cell sequence, not an exogenous sequence and/or a non-phage sequence (ie,

one more or all of these), eg, the sequence is a wild-type host chromosomal cell sequence such as antibiotic resistance gene or essential gene sequence comprised by a host cell chromosome. In an example, the sequence is a host cell plasmid sequence, eg, an antibiotic resistance gene sequence.

Optionally, the nuclease is a Cas and the target site is comprised by a protospacer sequence that is adjacent a NGG, NAG, NGA, NGC, NGGNG, NNGRRT or NNA-GAAW protospacer adjacent motif (PAM), eg, a AAAGAAA or TAAGAAA PAM (these sequences are written 5' to 3'). In an embodiment, the PAM is immediately adjacent the 3' end of the protospacer sequence. In an example, the Cas is a *S aureus*, *S thermophilus* or *Spyogenes* Cas. In an example, the Cas is Cpf1 and/or the PAM is TTN or CTA. Optionally, the Cas is a Type I (eg, Type I-A, I-B, I-C, I-D, I-E, or I-F) CRISPR system Cas. Optionally, the Cas is a Type II CRISPR system Cas. Optionally, the Cas is a Type III CRISPR system Cas. Optionally, the Cas is a Type IV CRISPR system Cas. Optionally, the Cas is a Type V CRISPR system Cas. Optionally, the Cas is a Type VI CRISPR system Cas.

Optionally, the nuclease is a Cas and each vector comprises a cognate CRISPR array that comprises multiple copies of the same spacer for targeting the target site. Optionally, there is provide a vector or plurality of vectors of the invention, wherein the vector(s) comprises a plurality of CRISPR arrays of said gRNA-encoding sequences for host cell protospacer sequence targeting, wherein the protospacers comprise the target site. Optionally, the or each vector comprises two, three or more of copies of nucleic acid sequences encoding crRNAs (eg, gRNAs), wherein the copies comprise the same spacer sequence for targeting a host cell target site (eg, a site comprised by a virulence, resistance or essential gene sequence).

In an example, at least two target sequences are modified by Cas, for example an antibiotic resistance gene and an essential gene. Multiple targeting in this way may be useful to reduce evolution of escape mutant host cells.

In an example, the Cas is a wild-type endogenous host cell Cas nuclease. In an example, target site cutting is carried out by a dsDNA Cas nuclease (eg, a Cas9, eg, a spCas9 or saCas9), whereby repair of the cut is by non-homologous end joining (NHEJ); alternatively the Cas is an exonuclease or Cas3

In an example, the array, gRNA-encoding sequence or vector is not in combination with a Cas endonuclease-encoding sequence that is naturally found in a cell together with repeat sequences of the array or gRNA-encoding sequence.

A tracrRNA sequence may be omitted from an array or vector of the invention, for example for Cas systems of a Type that does not use tracrRNA, or an endogenous tracrRNA may be used with the crRNA encoded by the vector.

In an example, the host target site is comprised by at least 5, 6, 7, 8, 9, 10, 20, 30 or 40 contiguous nucleotides.

In an example, the or each vector comprises an exogenous promoter functional for transcription of the crRNA or gRNA in the microbes.

Optionally, each vector is a plasmid, cosmid, virus, a virion, phage, phagemid or prophage.

For example, the invention provides a plurality of bacteriophage comprising a plurality of vectors of the invention, eg, wherein the vectors are identical. In an example, the vector is a viral vector. Viral vectors have a particularly limited capacity for exogenous DNA insertion, thus virus

packaging capacity needs to be considered. Room needs to be left for sequences encoding vital viral functions, such as for expressing coat proteins and polymerase. In an example, the vector is a phage vector or an AAV or lentiviral vector. Phage vectors are useful where the host is a bacterial cell. In an example, the vector is a virus capable of infecting an archaea host cell.

Optionally, vector components are comprised by a transposon that is capable of transfer into and/or between host cells. The transposon can be a transposon as described in US20160333348, GB1609811.3 and all US equivalent applications; the disclosures of these, including these specific transposon disclosures, are incorporated herein in its entirety and for potential inclusion of one or more disclosures therein in one or more claims herein.

In an example, the or each vector is provided by a nanoparticle or in liposomes.

In an example, the or each host cell (or first and/or second bacteria) is a gram positive cell. In an example, the or each host cell is an Enterobacteriaceae, eg, *Salmonella*, *Yersinia pestis*, *Klebsiella*, *Shigella*, *Proteus*, *Enterobacter*, *Serratia*, or *Citrobacter* cells. Optionally, the or each cell is an *E coli* (eg, *E coli* K12) or *Salmonella* (eg, *S enteric* serovar *typhimurium*) cell. Optionally, the or each host cell (or first and/or second bacteria) is a gram negative cell.

Optionally, the host (or first and/or second bacteria) is a *mycoplasma*, chlamydiae, spirochete or *mycobacterium*. Optionally, the host (or first and/or second bacteria) is a *Streptococcus* (eg, *pyogenes* or *thermophilus*) host. Optionally, the host (or first and/or second bacteria) is a *Staphylococcus* (eg, *aureus*, eg, MRSA) host. Optionally, the host (or first and/or second bacteria) is an *E. coli* (eg, O157: H7) host. Optionally, the host (or first and/or second bacteria) is a *Pseudomonas* (eg, *aeruginosa*) host. Optionally, the host (or first and/or second bacteria) is a *Vibro* (eg, *cholerae* (eg, O139) or *vulnificus*) host. Optionally, the host (or first and/or second bacteria) is a *Neisseria* (eg, gonorrhoeae or meningitidis) host. Optionally, the host (or first and/or second bacteria) is a *Bordetella* (eg, pertussis) host. Optionally, the host (or first and/or second bacteria) is a *Haemophilus* (eg, *influenzae*) host. Optionally, the host (or first and/or second bacteria) is a *Shigella* (eg, *dysenteriae*) host. Optionally, the host (or first and/or second bacteria) is a *Brucella* (eg, *abortus*) host. Optionally, the host (or first and/or second bacteria) is a *Francisella* host. Optionally, the host (or first and/or second bacteria) is a *Xanthomonas* host. Optionally, the host (or first and/or second bacteria) is a *Agrobacterium* host. Optionally, the host (or first and/or second bacteria) is a *Erwinia* host. Optionally, the host (or first and/or second bacteria) is a *Legionella* (eg, *pneumophila*) host. Optionally, the host (or first and/or second bacteria) is a *Listeria* (eg, *monocytogenes*) host. Optionally, the host (or first and/or second bacteria) is a *Campylobacter* (eg, *jejuni*) host. Optionally, the host (or first and/or second bacteria) is a *Yersinia* (eg, *pestis*) host. Optionally, the host (or first and/or second bacteria) is a *Borelia* (eg, *burgdorferi*) host. Optionally, the host (or first and/or second bacteria) is a *Helicobacter* (eg, *pylori*) host. Optionally, the host (or first and/or second bacteria) is a *Clostridium* (eg, *dificile* or *botulinum*) host. Optionally, the host (or first and/or second bacteria) is a *Erlchia* (eg, *chaffeensis*) host. Optionally, the host (or first and/or second bacteria) is a *Salmonella* (eg, *typhi* or *enterica*, eg, serotype *typhimurium*, eg, DT 104) host. Optionally, the host (or first and/or second bacteria) is a *Chlamydia* (eg, *pneumoniae*) host. Optionally, the host (or first and/or second bacteria) is a *Parachlamydia* host. Optionally, the host (or first and/or second bacteria) is a

Corynebacterium (eg, *amycolatum*) host. Optionally, the host (or first and/or second bacteria) is a *Klebsiella* (eg, *pneumoniae*) host. Optionally, the host (or first and/or second bacteria) is a *Enterococcus* (eg, *faecalis* or *faecium*, eg, linezolid-resistant) host. Optionally, the host (or first and/or second bacteria) is a *Acinetobacter* (eg, *baumannii*, eg, multiple drug resistant) host.

Optionally, the invention is for reducing the growth or proliferation of host cell(s) in an environment (eg, soil, a composition comprising said host cells and yeast cells), human, animal or plant microbiome. This is useful, for example, when the microbiome is naturally-occurring.

Optionally, the nuclease is a Cas and the target is comprised by a protospacer sequence comprising at least 5, 6, 7, 8, 9 or 10 contiguous nucleotides immediately 3' of a cognate PAM in the genome of the host cell, wherein the PAM is selected from AWG, AAG, AGG, GAG and ATG. Non-Medical, Ex vivo & In Vitro Uses Etc

In certain configurations, the inventive observation of rapid and durable microbial killing and growth or proliferation inhibition using nuclease cutting finds application in subjects other than humans and animals (eg, to treat plants or yeast cultures), or for ex vivo or in vitro treatment of substrates, such as industrial surfaces, fluids and apparatus. Thus, the invention further provides the following Concepts. Any other feature herein of the invention, its configurations, aspects, embodiments, options and examples above and elsewhere herein are combinable mutatis mutandis with these Concepts (including for providing combinations of features in the claims herein).

A Concept provides: Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of the invention, in the manufacture of a composition for carrying out a method of treatment as defined herein, wherein the subject is an organism other than a human or animal.

A Concept provides: Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of the invention, in the manufacture of a composition for carrying out an ex vivo or in vitro a method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.

Herein, treatment of an infection of a substrate may mean the treatment of a bacterial population (eg, one or more colonies) on a surface of the substrate and/or incorporated in the material of the substrate. For example, the treatment may be the treatment to kill bacteria on the surface of an industrial apparatus or equipment (eg, medical equipment, such as a scalpel or medical device or tubing). In another example, the substrate is a fluid (eg, a liquid or a gas), such as a medical fluid or petroleum product in fluid form (eg, an oil or hydrocarbon fluid or liquid).

A Concept provides: Use of a programmable nuclease in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first

species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.

Optionally the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject or substrate at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1. T1 and T2 may be as defined herein.

Optionally, the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment. Optionally, the infection is maintained by at least 100-fold for at least 60 minutes (eg, at least 120 minutes) after exposing the subject to the programmed nuclease.

Optionally, the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.

Optionally, the method comprises administering to the subject or substrate a RNA or a nucleic acid that encodes an RNA for expression of the RNA in or on the subject or substrate, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject or substrate.

Optionally, the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject or substrate.

Optionally, the subject or substrate comprises the nuclease prior to administration of the RNA or nucleic acid.

Optionally, a plurality of viruses (eg, phage) are administered to the subject or substrate, wherein each virus comprises a copy of the nucleic acid, wherein the viruses infect the microbes comprised by the subject or substrate to deliver thereto the nucleic acid.

Optionally, the ratio of administered viruses: microbes is from 10 to 150.

Optionally, the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.

Optionally, the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject or substrate to the programmed nuclease.

Optionally, the subject is a plant; or wherein the substrate is a metallic, plastic, concrete, stone, wood, glass or ceramic substrate. Optionally, the subject is a fluid (eg, a liquid or a gas).

Optionally, the microbes are bacteria or archaea. Optionally, the bacteria are gram positive bacteria. Optionally, the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).

Optionally, the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.

Reference is made to WO2016177682, which discusses aspects of microbiologically influenced corrosion (MIC) or biofouling of substrates and discloses methods for controlling MIC or biofouling of a substrate. The methods, nucleases, arrays, RNAs, vectors and viruses disclosed in that

document can be employed in the present invention, for example for carrying out the method or use of the present invention and the disclosures of these parts and the substrates and bacteria disclosed in WO2016177682 are incorporated herein by reference for potentially providing disclosure of features possible to be used in one or more claims herein.

Optionally, the use of the present invention is for controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate in an industrial or domestic system (eg, a system disclosed in WO2016177682, which disclosure is incorporated herein by reference). In an example, the system comprises equipment (eg, for use in an industrial process) and the surface is a surface of said equipment. In an example, the biofouling comprises microbial biofilm and/or sludge formation, proliferation or maintenance. In an example, the microbes are sessile. In an example “controlling” comprises preventing, reducing or eliminating said MIC or biofouling, or reducing spread of said MIC or biofouling in the system. Cell growth or proliferation or maintenance is, for example, a characteristic of cell viability. Thus, in an example, the method reduces microbe proliferation and/or maintenance.

Optionally, the microbes are comprised by a microbial biofilm that is in contact with said substrate. Optionally, said surface and host cells are in contact with a fluid, such as an aqueous liquid (eg, sea water, fresh water, stored water or potable water).

Fresh water is naturally occurring water on the Earth’s surface in ice sheets, ice caps, glaciers, icebergs, bogs, ponds, lakes, rivers and streams, and underground as groundwater in aquifers and underground streams. Fresh water is generally characterized by having low concentrations of dissolved salts and other total dissolved solids. The term specifically excludes sea water and brackish water, although it does include mineral-rich waters such as chalybeate springs. In an example said fresh water is any of these fresh water types. Potable water is water for human or animal (eg, livestock) consumption. In an example, the fluid is selected from industrial cooling water wherein the system is a cooling system; sewage water wherein the system is a sewage treatment or storage system; drinking water wherein the system is a drinking water processing, storage, transportation or delivery system; paper making water wherein the system is a paper manufacture or processing system; swimming pool water wherein the system is a swimming pool or swimming pool water treatment or storage system; fire extinguisher water wherein the system is a fire extinguishing system; or industrial process water in any pipe, tank, pit, pond or channel.

Optionally, the use is for controlling bacterial souring of a liquid in a reservoir or container), wherein the fluid comprises a population of first host cells of a first microbial species that mediates said biofouling, the method comprising

(i) contacting the population with a plurality of vectors that are capable of transforming or transducing the cells, each vector comprising a CRISPR array whereby CRISPR arrays are introduced into the host cells, wherein

(a) each CRISPR array comprises one or more sequences for expression of a crRNA and a promoter for transcription of the sequence(s) in a host cell; and

(b) each crRNA is capable of hybridising to a target sequence of a host cell to guide Cas (eg, a Cas nuclease) in the host cell to modify the target sequence (eg, to cut the target sequence); the target sequence being a gene sequence for mediating host cell viability; and

wherein the method comprises allowing expression of said cRNAs in the presence of Cas in host cells, thereby modifying target sequences in host cells, resulting in reduction of host cell viability and control of said biofouling.

In an example, the fluid is a liquid. In an example, the fluid is a gaseous fluid. Systems.

An example system is selected from the group consisting of a: —

Petrochemical recovery, processing, storage or transportation system; hydrocarbon recovery, processing, storage or transportation system; crude oil recovery, processing, storage or transportation system; natural gas recovery, processing, storage or transportation system, (eg, an oil well, oil rig, oil drilling equipment, oil pumping system, oil pipeline, gas rig, gas extraction equipment, gas pumping equipment, gas pipeline, oil tanker, gas tanker, oil storage equipment or gas storage equipment); Water processing or storage equipment; water reservoir (eg, potable water reservoir); Air or water conditioning (eg, cooling or heating) equipment, eg, a coolant tube, condenser or heat exchanger; Medical or surgical equipment; Environmental (eg, soil, waterway or air) treatment equipment; Paper manufacturing or recycling equipment; Power plant, eg, a thermal or nuclear power plant; Fuel (eg, hydrocarbon fuel, eg, petroleum, diesel or LPG) storage equipment; Mining or metallurgical, mineral or fuel recovery system, eg, a mine or mining equipment; Engineering system; Shipping equipment; Cargo or goods storage equipment (eg, a freight container); Food or beverage manufacturing, processing or packaging equipment; Cleaning equipment (eg, laundry equipment, eg, a washing machine or dishwasher); Catering (eg, domestic or commercial catering) equipment; Farming equipment; Construction (eg, building, utilities infrastructure or road construction) equipment; Aviation equipment; Aerospace equipment; Transportation equipment (eg, a motor vehicle (eg, a car, lorry or van); a railcar; an aircraft (eg, an aeroplane) or a marine or waterway vehicle (eg, a boat or ship, submarine or hovercraft)); Packaging equipment, eg, consumer goods packaging equipment; or food or beverage packaging equipment; Electronics (eg, a computer or mobile phone or an electronics component thereof); or electronics manufacture or packaging equipment; Dentistry equipment; Industrial or domestic piping (eg, a sub-sea pipe) or storage vessel (eg, a water tank or a fuel tank (eg, gasoline tank, eg, a gasoline tank of a vehicle)); Underground equipment; Building (eg, a dwelling or office or commercial premises or factory or power station); Roadway; Bridge; Agricultural equipment; Factory system; Crude oil or natural gas exploration equipment; Office system; and a Household system.

In an example, the system is used in an industry or business selected from the group consisting of agriculture, oil or petroleum industry, food or drink industry, clothing industry, packaging industry, electronics industry, computer industry, environmental industry, chemical industry, aerospace industry, automotive industry, biotechnology industry, medical industry, healthcare industry, dentistry industry, energy industry, consumer products industry, pharmaceutical industry, mining industry, cleaning industry, forestry industry, fishing industry, leisure industry, recycling industry, cosmetics industry, plastics industry, pulp or paper industry, textile industry, clothing industry, leather or suede or animal hide industry, tobacco industry and steel industry. In an example, the surface or fluid to be treated is a surface or fluid of equipment used in said selected industry. In an example, the system is used in the crude oil industry. In an example, the system is used in the natural gas industry. In an

example, the system is used in the petroleum industry. In an example, the system is a sea container, platform or rig (eg, oil or gas platform or rig for use at sea or at sea), ship or boat. In an embodiment, such a system is anchored at sea; eg, non-temporarily anchored at sea, eg, has been anchored at sea for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or more months (eg, contiguous months). In an embodiment, such a system is in the waters of a country or state; eg, non-temporarily at sea in such waters, eg, has been in waters of said country for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or more months (eg, contiguous months).

In an example, the substrate surface to be treated comprises stainless steel, carbon steel, copper, nickel, brass, aluminium, concrete, a plastic or wood. In an example, the substrate is a metal weld or join. In an example, the surface is a metallic (eg, steel or iron) or non-metallic (eg, plastic, concrete, asphalt, wood, rubber or stone) surface. In an example, the metal is an alloy (eg, stainless steel, brass or a nickel-, zinc-, copper-, nickel- or aluminium-alloy). In an example, the surface is a man-made polymer surface. In an example, the surface is a substrate coating. In an example, the substrate is in contact with soil, fresh water or sea water.

In an example, the fluid is potable water; a waterway; brackish water; or a liquid fuel, eg, gasoline or diesel (eg, for a car or motorised vehicle), LPG, kerosine, an alcohol (eg, ethanol, methanol or butanol), liquid hydrogen or liquid ammonia), in an example, the fuel is stored liquid fuel. In an example the fluid is an oil or non-aqueous liquid. In an example, the fluid is a liquid comprised by a waterway or body of water, eg, sea water, fresh water, potable water, a river, a stream, a pond, a lake, a reservoir, stored water (eg, in a water storage tank or cooling equipment), groundwater, well water, water in a rock formation, soil water or rainwater. In an example, the liquid is sea water. In an example, the substrate is in contact with a liquid mentioned in this paragraph. In an example, the fluid or liquid is selected from the group consisting of an oil, an aqueous solution, a hydraulic fracturing fluid, a fuel, carbon dioxide, a natural gas, an oil/water mixture, a fuel/water mixture, water containing salts, ocean or sea water, brackish water, sources of fresh water, lakes, rivers, stream, bogs, ponds, marshes, runoff from the thawing of snow or ice, springs, groundwater, aquifers, precipitation, any substance that is a liquid at ambient temperature (eg, at rtp) and is hydrophobic but soluble in organic solvents, hexanes, benzene, toluene, chloroform, diethyl ether, vegetable oils, petrochemical oils, crude oil, refined petrochemical products, volatile essential oils, fossil fuels, gasoline, mixtures of hydrocarbons, jet fuel, rocket fuel, biofuels. In an example the fluid is an oil/water mixture.

The terms “microbiologically influenced corrosion” or “MIC” as used herein, unless otherwise specified, refer to processes in which any element (substrate) of a system is structurally compromised due to the action of at least one member of a microbial population, eg, bacterial or archaeal population. The term “biofouling” as used herein, unless otherwise specified, refers to processes in which microorganisms (such as bacteria and/or archaea) accumulate on a substrate surface in contact with a fluid (eg, water or an aqueous liquid, or a hydrocarbon, or a petrochemical).

Also included is the undesirable accumulation and proliferation of microorganisms (such as bacteria and/or archaea) in a fluid (eg, water or an aqueous liquid, or a hydrocarbon, or a petrochemical), ie, “souring” of the fluid. In an example, the bacteria are comprised by ship or boat ballast water and the bacteria are environmentally undesir-

able. The term “substrate” as used herein refers to any type of surface on which cells can attach and a biofilm can form and grow or on which biofouling (eg slime or sludge formation) can occur. The substrate may be an “industrial” substrate such as the surface of equipment in an petrochemical, fuel, crude oil or gas piping system, or a “non-industrial” (eg, domestic, eg, household or office) substrate such as a kitchen counter or a shower substrate or a garden substrate.

In an alternative, instead of a population of host bacterial cells, the population is a population of archaeal cells of a first species.

Optionally, said fluid is an aqueous liquid (eg, sea water, fresh water, stored water or potable water).

In an alternative, instead the microbes are algal cells.

Optionally, the microbes are sulphate reducing bacteria (SRB) cells (eg, *Desulfovibrio* or *Desulfotomaculum* cells). In an example, the cells are selected from the group consisting of *Desulfotomaculum nigrificans*, *Desulfacinum infernum*, *Thermodesulfobacterium mobile*, *Thermodesulforhabdus norvegicus*, *Archaeoglobus fulgidus*, *Desulfomicrobium apsheronum*, *Desulfovibrio gabonensis*, *Desulfovibrio longus*, *Desulfovibrio vietnamensis*, *Desulfobacterium cetonicum*, *Desulphomaculum halophilum*, *Desulfobacter vibriiformis* and *Desulfotomaculum thermocisternum* cells. In an example, the population comprises a mixture of two or more of these cell species.

Optionally, the surface or fluid is comprised by a crude oil, gas or petrochemicals recovery, processing, storage or transportation equipment. Crude oil is one of the most important energetic resources in the world. It is used as raw material in numerous industries, including the refinery-petrochemical industry, where crude oil is refined through various technological processes into consumer products such as gasoline, oils, paraffin oils, lubricants, asphalt, domestic fuel oil, vaseline, and polymers. Oil-derived products are also commonly used in many other chemical processes. In an alternative, the fluid is a said consumer product or the surface is in contact with such a consumer product.

Optionally, the surface is in contact with sea water, a fracking liquid or liquid in a well; or wherein the fluid is sea water, a fracking liquid or liquid in a well.

Optionally, step (i) of the method comprises providing a population of microbial cells of a second species (second host cells), the second cells comprising said vectors, wherein the vectors are capable of transfer from the second host cells to the first host cells; and combining the second host cells with the first host cells, whereby vectors are introduced into the first host cells. In an example, the second cell(s) are environmentally-, industrially-, or domestically-acceptable in an environment (eg, in a water or soil environment) and the first host cell(s) are not acceptable in the environment.

Optionally, the first host cells are comprised by a mixture of microbial cells (eg, comprised by a microbial biofilm) before contact with said vectors, wherein the mixture comprises cells of said second species.

Optionally, said second species is a species of *Bacillus* or nitrate-reducing bacteria or nitrate reducing sulfide oxidizing bacteria (NRB)

Optionally, the NRB is selected from the group consisting of *Campylobacter* sp., *Nitrobacter* sp., *Nitrosomonas* sp., *Thiomicrospira* sp., *Sulfurospirillum* sp., *Thauera* sp., *Paracoccus* sp., *Pseudomonas* sp., *Rhodobacter* sp. and *Desulfovibrio* sp; or comprises at least 2 of said species.

Optionally, the NRB is selected from the group consisting of *Nitrobacter vulgaris*, *Nitrosomonas europea*, *Pseudomo-*

nas stutzeri, *Pseudomonas aeruginosa*, *Paracoccus denitrificans*, *Sulfurospirillum deleyianum*, and *Rhodobacter sphaeroides*.

Optionally, the method comprises contacting the host cells of said first species with a biocide simultaneously or sequentially with said vectors. In an example, the vectors and biocide are provided pre-mixed in a composition that is contacted with the host cells.

Optionally, the biocide is selected from the group consisting of tetrakis hydroxymethyl phosphonium sulfate (THPS), glutaraldehyde, chlorine monoxide, chlorine dioxide, calcium hypochlorite, potassium hypochlorite, sodium hypochlorite, dibromonitripropionamide (DBNPA), methylene bis(thiocyanate) (MBT), 2-(thiocyanomethylthio) benzothiazole (TCMTB), bronopol, 2-bromo-2-nitro-1,3-propanediol (BNPD), tributyl tetradecyl phosphonium chloride (TTPC), taurinamide and derivatives thereof, phenols, quaternary ammonium salts, chlorine-containing agents, quinaldinium salts, lactones, organic dyes, thiosemicarbazones, quinones, carbamates, urea, salicylamide, carbamylide, guanide, amidines, imidazolines, acetic acid, benzoic acid, sorbic acid, propionic acid, boric acid, dehydroacetic acid, sulfurous acid, vanillic acid, p-hydroxybenzoate esters, isopropanol, propylene glycol, benzyl alcohol, chlorobutanol, phenylethyl alcohol, formaldehyde, iodine and solutions thereof, povidone-iodine, hexamethylenetetramine, noxythiolin, 1-(3-chloroallyl)-3,5,7-triazo-1-azoniaadamantane chloride, taurolidine, taurultam, N-(5-nitro-2-furfurylidene)-1-amino-hydantoin, 5-nitro-2-furaldehyde semicarbazone, 3,4,4'-trichlorocarbonyl, 3,4', 5-tribromosalicylanilide, 3-trifluoromethyl-4,4'-dichlorocarbonyl, 8-hydroxyquinoline, 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid, 1,4-dihydro-1-ethyl-6-fluoro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid, hydrogen peroxide, peracetic acid, sodium oxychlorosene, parachlorometaxyleneol, 2,4,4'-trichloro-2'-hydroxydiphenol, thymol, chlorhexidine, benzalkonium chloride, cetylpyridinium chloride, silver sulfadiazine, silver nitrate, bromine, ozone, 40 quinoxalones, polyoxyethylene (dimethylimino) ethylene (dimethylimino) ethylene dichloride, 2-(tert-butylamino)-4-chloro-6-ethylamino-5'-triazine (terbutylazine), and combinations thereof. In an example the biocide is tetrakis hydroxymethyl phosphonium sulfate (THPS). In an example, the biocide is a quaternary ammonium compound.

Optionally, the system is used in an industry operation selected from the group consisting of mining; shipping; crude oil, gas or petrochemicals recovery or processing; hydraulic fracturing; air or water heating or cooling; potable water production, storage or delivery; transportation of hydrocarbons; and wastewater treatment.

Optionally, the surface is a surface of equipment used in said selected industry; or wherein the fluid is a fluid comprised by equipment used in said selected industry.

Optionally, the surface is a surface of kitchen, bathing or gardening equipment; or wherein the fluid is comprised by kitchen, bathing or gardening equipment. For example, the equipment is used in a domestic setting.

Optionally, the fluid is a potable liquid contained in a container (eg, water tank or bottle) and the surface is a surface of the container in contact with the liquid.

Optionally, each vector comprises a mobile genetic element (MGE), wherein the MGE comprises an origin of transfer (oriT) and a said CRISPR array; wherein the MGE is capable of transfer between a host cell of said first species and a further microbial host cell in said industrial or domestic system. For example, the further cell(s) are environmen-

tally-, industrially-, or domestically-acceptable in an environment (eg, in a water or soil environment) and the first host cell(s) are not acceptable in the environment. Optionally, the oriT is functional in the first and further host cells.

Optionally, the first and further host cells are comprised by a biofilm of fluid in contact with said surface; or wherein said cells are comprised by said fluid.

Optionally, each MGE is or comprises an integrative and conjugative element (ICE); or wherein each vector is a phage that is capable of infecting host cells of said first species and each MGE is a phage nucleic acid that is capable of said transfer between the cells. Optionally, each ICE is a transposon, eg, a conjugative transposon. Optionally, each vector is a plasmid, optionally comprising an MGE as described herein. Optionally, the sequences are comprised by a conjugative transposon of the first cell and/or further cell.

In an example, the method is a method of controlling microbiologically influenced corrosion (MIC) or biofouling of a substrate comprised by a crude oil, gas or petrochemicals recovery, processing, storage or transportation equipment (eg, a crude oil tanker, oil rig or oil drilling equipment), wherein a surface of the substrate is in contact with a population of first host cells, wherein the first host cells are sulphur- or sulphate-reducing bacteria (SRB), extracellular polymeric substance-producing bacteria (EPSB), acid-producing bacteria (APB), sulphur- or sulphide-oxidizing bacteria (SOB), iron-oxidising bacteria (IOB), manganese-oxidising bacteria (MOB), ammonia producing bacteria (AmPB) or acetate producing bacteria (AcPB) of a first species that mediates MIC or biofouling of the substrate, wherein the surface and cell population are in contact with a liquid selected from sea water, fresh water, a fracking liquid or liquid in a well (eg, oil or natural gas well), the method comprising

(i) contacting the cell population with vectors by mixing the liquid with a plurality of vectors that are capable of transforming or transducing first host cells, each vector comprising a CRISPR array whereby CRISPR arrays are introduced into the host cells, wherein

(a) each CRISPR array comprises one or more sequences for expression of a crRNA and a promoter for transcription of the sequence(s) in a host cell;

(b) each crRNA is capable of hybridising to a target sequence of a host cell to guide Cas (eg, a Cas nuclease, eg, a Cas9 or Cfp1) in the host cell to modify the target sequence (eg, to cut the target sequence); the target sequence being a gene sequence for mediating host cell viability;

(c) wherein each sequence of (a) comprises a sequence R1-S1-R1' for expression and production of the respective crRNA in a first host cell, wherein R1 is a first CRISPR repeat, R1' is a second CRISPR repeat, and R1 or R1' is optional; and S1 is a first CRISPR spacer that comprises or consists of a nucleotide sequence that is 70, 75, 80, 85, 90 or 95% or more identical to a target sequence of a said first host cell and

(ii) allowing expression of said crRNAs in the presence of Cas in host cells, thereby modifying target sequences in host cells, resulting in reduction of host cell viability and control of MIC or biofouling of said substrate. In an embodiment, both R1 and R1' are present.

In an example, the method is a method of controlling bacterial biofouling in ballast water of a ship or boat, wherein the water comprises a population of first host cells of a first microbial species that mediates said biofouling, the method comprising

(i) contacting the population with a plurality of vectors that are capable of transforming or transducing the cells, each vector comprising a CRISPR array whereby CRISPR arrays are introduced into the host cells, wherein

(a) each CRISPR array comprises one or more sequences for expression of a crRNA and a promoter for transcription of the sequence(s) in a host cell; and

(b) each crRNA is capable of hybridising to a target sequence of a host cell to guide Cas (eg, a Cas nuclease) in the host cell to modify the target sequence (eg, to cut the target sequence); the target sequence being a gene sequence for mediating host cell viability; and

(ii) allowing expression of said crRNAs in the presence of Cas in host cells, thereby modifying target sequences in host cells, resulting in reduction of host cell viability and control of said biofouling.

Optionally, the first host cells are *Vibrio cholerae*, *E coli* or Enterococci sp cells.

Optionally, step (i) comprises mixing the ballast water with the vectors, eg, in the hull of a ship or boat. Optionally, the ship or boat is a marine vehicle and the water is sea water. Optionally, instead of a ship or boat, the ballast water is comprised by a container or a drilling platform at sea, eg, an oil platform or oil rig. In an example, the ship, boat, container, platform or rig is anchored at sea (ie, not temporarily in its location).

In an example, the method is a method of discharging ballast water from a ship or boat, wherein the discharged ballast water comprises water treated by the method. Optionally, the water is discharged into a body of water, eg, a sea, ocean or waterway (eg, a river, canal, lake or reservoir) or into a container.

Paragraphs:

The invention provides the following Paragraphs, which are supported by the Examples below: —

1. A programmable Cas (eg, Cas3 or Cas9) nuclease for use in a method of treating *E coli* or *C difficile* infection of a subject, wherein the Cas nuclease is programmable with a guide RNA to cut a target site comprised by the genomes of *E coli* or *C difficile* bacteria that have infected the subject, whereby *E coli* or *C difficile* cells are killed, or growth or proliferation of the cells is reduced, the treatment method comprising exposing the subject to the Cas nuclease wherein the nuclease is programmed with guide RNA to cut the target site, whereby genomes of the *E coli* or *C difficile* bacteria comprised by the subject are cut and the infection of the subject is reduced by at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
2. A programmable Cas (eg, Cas3 or Cas9) nuclease (optionally according to paragraph 1) for use in a method of treating *E coli* or *C difficile* infection of a subject, wherein the Cas nuclease is programmable with a guide RNA to cut a target site comprised by the genomes of *E coli* or *C difficile* bacteria that have infected the subject, whereby *E coli* or *C difficile* cells are killed, or growth or proliferation of the cells is reduced, the treatment method comprising exposing the subject to the Cas nuclease wherein the nuclease is programmed with guide RNA to cut the target site, whereby genomes of the *E coli* or *C difficile* bacteria comprised by the subject are cut and the infection of the subject is reduced, wherein a reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.

3. The nuclease of any preceding Paragraph, wherein at least 60% of the infection is reduced by 60 minutes after exposing the subject to the programmed nuclease.
4. The nuclease of any preceding Paragraph, wherein the nuclease (eg, programmed nuclease) and/or a nucleic acid encoding the guide RNA is administered to the subject at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1.
5. The nuclease of any preceding Paragraph, wherein the method comprises reducing the infection such that the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.
6. The nuclease of any preceding Paragraph, wherein the method comprises administering to the subject the RNA or a nucleic acid that encodes the RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.
7. The nuclease of any preceding Paragraph, wherein the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid encoding the RNA to the subject.
8. The nuclease of Paragraph 7, wherein the subject comprises the nuclease prior to administration of the RNA or nucleic acid to the subject.
9. The nuclease of any preceding Paragraph, wherein a plurality of viruses (eg, phage) are administered to the subject, wherein each virus comprises a copy of a nucleic acid encoding the RNA, wherein the viruses infect the microbes comprised by the subject to deliver thereto the nucleic acid.
10. The nuclease of Paragraph 9, wherein the ratio of administered viruses: microbes comprised by the subject is from 10 to 150.
11. The nuclease according to any preceding Paragraph, wherein the subject is a human or animal, optionally wherein the subject is a human over 65 years of age or is a pediatric patient.
12. The nuclease according to Paragraph 11, wherein the infection is an infection of the lungs, abdomen or urinary tract; or wherein the subject has undergone surgery, is on an immunosuppressant medication and/or is suffering from a chronic disease.
13. The nuclease according to any preceding Paragraph, wherein the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
14. The nuclease according to any preceding Paragraph, wherein the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.
15. The nuclease according to any one of Paragraphs 11 to 14, wherein the method treats or prevents septicaemia and/or sepsis (eg, septic shock) in the subject.
16. The nuclease of Paragraph 16, wherein at the start of the treatment, the subject (eg, a human) has a temperature of $<36^{\circ}\text{C}$. or $>38^{\circ}\text{C}$.; a heart rate of $>90/\text{min}$, a respiratory rate of >20 breaths/min or $\text{PaCO}_2 < 4.3\text{ kPa}$; and white blood cell count of $<4000/\text{mm}^3$ or $>12,000/\text{mm}^3$.
17. The nuclease of Paragraph 15 or 16, wherein at the start of the treatment, the subject (eg, a human) has presence of two or more of the following: abnormal

- body temperature, abnormal heart rate, abnormal respiratory rate, abnormal blood gas and abnormal white blood cell count.
18. The nuclease of any preceding Paragraph, wherein the subject is a human or animal and the microbes are bacteria (eg, *E coli* or *C difficile*), wherein blood infection of the subject by the bacteria is reduced at least 100- or 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
19. The nuclease of any one of Paragraphs 11 to 18, wherein the blood of the subject is infected with from 10^7 to 10^{12} CFU/ml of the bacteria immediately before the treatment.
20. The nuclease according to any one of Paragraphs 1 to 10, wherein the subject is a plant.
21. The nuclease according to any preceding Paragraph, wherein the bacteria are comprised by a microbiome.
22. The nuclease according to Paragraph 21, wherein the microbiome comprises *Lactobacillus* and/or *Streptococcus* bacteria.
23. The nuclease according to any preceding Paragraph, wherein the *E coli* are EHEC *E coli*.
24. The nuclease according to any preceding Paragraph, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.
25. A plurality of viruses (eg, phage or phagemids for producing phage) for use with the nuclease of any preceding Paragraph in the method of treatment, wherein each virus comprises a copy of a nucleic acid encoding the RNA, wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.
26. A composition comprising a plurality of nucleic acids for programming the nuclease of any one of Paragraphs 1 to 24 in the method of treatment, wherein each nucleic acid is a nucleic acid as defined in any one of Paragraphs 6 to 9.
27. A CRISPR/Cas system comprising a nuclease according to any preceding Paragraph for use in the method of treatment, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of the microbes.
28. A guide RNA or a DNA encoding a guide RNA for use in the system of Paragraph 27 for use in the method of treating an acute microbial infection in the subject, eg, septicaemia or sepsis.
29. A nucleic acid vector comprising the guide RNA or DNA recited in Paragraph 27 or 28.
30. The vector of Paragraph 29 wherein the vector is a phage, phagemid, viriophage, virus, plasmid (eg, conjugative plasmid) or transposon.
31. An anti-sepsis or anti-septicaemia composition for administration to a human or animal for treating sepsis or septicaemia, the composition comprising a plurality of vectors, wherein each vector is according to Paragraph 29 or 30.
32. A method of treating an acute microbial infection of a subject, wherein the method is as defined by any preceding Paragraph.
33. Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of any one of Paragraphs 1 to 25 and 27 to 30, in the manufacture of a composition for carrying out a method of treatment as defined by any

- preceding Paragraph, wherein the subject is an organism other than a human or animal.
34. Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of any one of Paragraphs 1 to 25 and 27 to 30, in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.
35. Use of a programmable nuclease in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.
36. The use of Paragraph 33, 34, 35, wherein the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject or substrate at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1.
37. The use of any one of Paragraphs 33 to 36, wherein the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
38. The use of any one of Paragraphs 33 to 37, wherein the reduction of the infection is maintained by at least 100-fold for at least 60 minutes (eg, at least 120 minutes) after exposing the subject to the programmed nuclease.
39. The use of any one of Paragraphs 33 to 38, wherein the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.
40. The use of any one of Paragraphs 33 to 39, wherein the method comprises administering to the subject or substrate a RNA or a nucleic acid that encodes an RNA for expression of the RNA in or on the subject or substrate, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject or substrate.
41. The use of Paragraph 40, wherein the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject or substrate.
42. The use of Paragraph 40, wherein the subject or substrate comprises the nuclease prior to administration of the RNA or nucleic acid.
43. The use of any one of Paragraphs 40 to 42, wherein a plurality of viruses (eg, phage) are administered to the subject or substrate, wherein each virus comprises a copy of the nucleic acid, wherein the viruses infect the

- microbes comprised by the subject or substrate to deliver thereto the nucleic acid.
44. The use of Paragraph 43, wherein the ratio of administered viruses: microbes is from 10 to 150.
45. The use of any one of Paragraphs 33 to 44, wherein the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
46. The use of any one of Paragraphs 44 to 45, wherein the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject or substrate to the programmed nuclease.
47. The use of any one of Paragraphs 33 to 46, wherein the subject is a plant; or wherein the substrate is a metallic, plastic, concrete, stone, wood, glass or ceramic substrate.
48. The use of any one of Paragraphs 33 to 47, wherein the microbes are bacteria.
49. The use according to Paragraph 48, wherein the bacteria are gram positive bacteria.
50. The use according to Paragraph 48 or 49, wherein the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonnorrhoea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).
51. The use of any one of Paragraphs 33 to 50, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.

Treatment of Pathogenic Bacterial Infections

Infectious complications are a serious cause of morbidity and mortality in cancer patients, especially those with underlying haematological malignancies where autopsy studies demonstrate that approximately 60% of deaths are infection related. Although fewer data exist on infectious mortality in patients with solid organ tumours, approximately 50% of these patients are estimated to have an infection as either the primary or an associated cause of death (“Epidemiology of Infections in Cancer Patients”, in “Infectious Complications in Cancer Patients”, Springer International Publishing Switzerland (2014)). Bacterial infections dominate. These infectious complications remain a significant limitation of cancer treatment modalities.

The detrimental effects of classic antibiotic treatment with broad-spectrum antibiotics have been demonstrated in immune checkpoint inhibitor (ICI)-treated cancer patients. Routy et al investigated how the gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumours (Routy et al Science 2018, 359, 91-97). In this work, the authors also analyzed datasets for infections/antibiotic use in patients with advanced NSCLC (n=140), renal cell carcinoma (n=67), or urothelial carcinoma (n=42) who received antibody ICI against PD-1/PD-L1 interaction after one or several prior therapies. Among these patients, they were prescribed broad-spectrum antibiotics (beta-lactam+/-inhibitors, fluoroquinolones, or macrolides) within 2 months before, or 1 month after, the first administration of PD-1/PD-L1 mAb. Patients generally took antibiotic orally for common indications (dental, urinary, and pulmonary infections). The detrimental effect of treating

infections in cancer patients undergoing ICI therapy with classical, broad-spectrum antibiotics was observed. See FIG. 8, which shows that the antibiotic treatment during ICI therapy has fatal outcomes: there was a medial overall survival of 21.9 months in the absence of antibiotic treatment, compared to an overall survival of 9.8 months with antibiotic treatment. So, the median overall survival in patients treated with classical antibiotics is <50% (or >12 months shorter) that of patients not receiving antibiotic treatment.

The work by Gopalakrishnan et al is another recent example lending support to the importance of a “healthy” microbiome in immuno-oncology therapy outcomes (Gopalakrishnan et al, *Science* 2018, 359, 97-103). See FIGS. 9A and 9B. It was observed that the gut microbiome modulates the efficacy of anti-PD-1 inhibition in melanoma patients.

Several other studies add to the expanding evidence base of the critical link between the microbiome and immuno-oncology outcomes:

“Microbiota: a key orchestrator of cancer therapy”, *Nat. Rev. Cancer* 2017, 17, 271-285

Matson et al, *Science* 2018, 359, 104-108

L. Derosa et al *Annals of Oncology* 2018 (epub 30 Mar. 2018)

M. Vétizou et al *Science*. 2015, 350, 1079-84

Sivan et al *Science* 2015, 350, 1084-1089

Another report—claiming to be the first systematic review of infection among patients receiving immune checkpoint blockade for cancer therapy—investigated serious infections in melanoma patients treated with immune checkpoint inhibitors (against CTLA-4, PD-1, and/or PD-L1) (M. Del Castillo et al *Clin. Infect. Dis.* 2016, 63, 1490-1493). Serious infections were defined as infections requiring hospitalization or parenteral antimicrobials. Of 740 patients (898 courses of immune checkpoint blockade), serious infection developed in 54 patients (7.3%). Nine patients (17%) were deemed to have died of an infection. Total number of infections was 58, as some patients developed >1 infection. The majority of infections were bacterial in origin (~80%; i.e., bacterial infections: 80% of 7.3%: 5.8% of patients). Pneumonia and bloodstream infections were the two dominating bacterial infection types.

Immune checkpoint-blocking drugs are associated with immune-related adverse effects (irAEs) related to the upregulated immune system. The complications are managed with immunosuppressive drugs, such as steroids (immunosuppression is a risk factor for subsequent opportunistic infections). Of the 740 patients, 46% received steroids during the course of treatment. Risk of serious infections was 13.5% in the cohort receiving corticosteroids or infliximab (vs. 7.3% in the overall population).

In yet another report, the emerging concern of infectious diseases in lung cancer patients receiving ICI therapy was investigated. Of 84 NSCLC patients receiving nivolumab (a PD-1 inhibitor), 20 patients (23.8%) developed an infectious disease. Bacterial infections accounted for 75% of infections; i.e., bacterial infections in 18% of patients. Most common type of bacterial infection was pneumonia. See K. Fujita et al *Eur. Resp. J.* 2017, 50, OA1478.

The Gram-negative *bacillus E. coli* is one of the most common causes of bacteraemia in patients with cancer. The all-cause 30-day mortality rate for this pathogen is high (~15%) (Y. E. Ha et al *Int. J. Antimicrob. Agen.* 2013, 42, 403-409). Published estimates of 30-day all-cause mortality among *E. coli* bacteraemia patients (cancer/non-cancer) vary from around 10 to 35% (J. K. Abernethy et al *Clin. Micro-*

biol. Infect. 2015, 21, 251.e1-251.e8), clearly highlighting the high burden associated with just this pathogen. Overall, causative pathogens in bacteraemia are primarily Gram-negative bacteria (65%), with *E. coli* (18.3%), *P. aeruginosa* (18.3%), and *K. pneumoniae* (17.3%) being the most common organisms encountered; the three pathogens together account for 54% of the bacteraemia cases, or 85% of Gram-negative cases, according to a study investigating >100 bacteraemia cases in cancer patients (G. Samonis et al *Support Care Cancer* 2013, 21, 2521-2526). In-hospital mortality was 26.2% in this study. Comparable numbers can be found elsewhere. For example, a study of neutropenic and non-neutropenic adult cancer patients with bloodstream infections investigated 399 cases of bloodstream infections in 344 cancer patients: The largest causative pathogen group was Gram-negative bacilli (45%). Of the clinical isolates, *E. coli* (35%) accounted for the most cases within Gram-negatives, followed by *K. pneumoniae* (20%) and *P. aeruginosa* (19%) (E. Velasco et al *Eur. J. Clin. Microbiol. Infect. Dis.* 2006, 25, 1-7). The three pathogens collectively account for 33% of the bacteraemia cases (or 74% of Gram-negative cases). The overall 30-day mortality rate was 32% in this study. Two other reports looked at causative agents of bloodstream infection in patients with solid tumours and also found Gram-negative bacteria to be the dominating pathogen type (47-55% of the infections, across several hundred patients) (M. Marin et al *Medicine* 2014, 93, 143-149; M. Anatoliotaki et al *Infection* 2004, 32, 65-71; see also C. Gudiol et al *Virulence* 2016, 7, 298-308). In the larger of the two studies (with more robust numbers for individual pathogens), the three main pathogens within the Gram-negative group were again *E. coli* (55%), *P. aeruginosa* (18%), and *Klebsiella* spp. (11%)—corresponding to 92% of the Gram-negative cases, or 51% of the 528 total cases of bloodstream infections studied.

The above data on specific causative infectious pathogens in cancer patients are summarized in Table 5 below.

Thus, available data on bloodstream infections in cancer indicate that Gram-negative pathogens are involved in 45-65% of the infection cases, with three key pathogens—*E. coli*, *K. pneumoniae*, and *P. aeruginosa*—being the culprits in the vast majority of Gram-negative cases (73-92%).

The inventors, thus, formulated an oncologist’s dilemma: Reduction in efficacy of the cancer therapy is likely due to the reduced microbiome diversity resulting from antibiotic therapy

At least 1/3 of patients on checkpoint inhibitors get serious and life-threatening infections

Not treating these infections could result in death from the infection (1-2 weeks)

Treatment with classic antibiotics leads to reduction in progression free survival after 4 years from >40% to around 10%.

The choice is to treat the immediate need of a potentially fatal infection (which must be addressed) at the risk of seriously undermining the cancer therapy.

The inventors realised, therefore, that there is a need for methods that can treat a bacterial pathogenic infection in a different way that minimizes compromise to the cancer therapy. The inventors realised that this need would also be useful in other therapy settings where the microbiome composition can modulate therapy outcomes, eg, in transplant settings.

Whilst not wishing to be bound by any particular theory, the inventors believe that alleviating the detrimental effect of traditional antibiotic therapy on overall survival in ICI patients using the invention may, in some embodiments,

translate to as much as a doubling of overall survival (or >12 months). Capturing a treatment effect of several months in terms of median overall survival is a very substantial achievement in this space. In fact, an effect size of this order of magnitude is comparable to the outcomes reported for ICI trials (i.e., where benefits usually are measured in months, not years). Additionally, PD-1/PD-L1 drugs are projected to dominate the ICI market. In 2023, PD-1/PD-L1 are projected to account for 94% of \$46B USD global sales of ICIs (CTLA-4 blockers only account for 6%), source: “Landscape & Forecast: Immune Checkpoint Inhibitors”, Decision Resources, December 2017. Thus, a need for improving treatment using immune checkpoint inhibitors of PD-1 or PD-L1 is particularly pressing in medicine, and we believe that the present invention finds particular benefit in this respect.

In an example, the method removes the need to administer a classic antibiotic, such as a broad-spectrum antibiotic (or any other one disclosed herein). In another example, the invention reduces the amount or dosing frequency of a classic antibiotic, such as a broad-spectrum antibiotic (or any other one disclosed herein) that is administered to the subject for treating the infection. For example, the subject can be administered a low-dose broad-spectrum antibiotic (eg, 50, 40, 30, 20, 10% or less of a conventional dose) whilst the guided nuclease cutting is used, and thus treatment of the infection in this setting. The invention may be particularly beneficial for patients on immunosuppressants, eg, for cancer patients, transplant patients or patients suffering from a viral infection (eg, HIV (human immunodeficiency virus), CMV (cytomegalovirus) or RSV (respiratory syncytial virus) infection).

The term “broad-spectrum antibiotic” can refer to an antibiotic that acts on the two major bacterial groups, gram-positive and gram-negative, or any antibiotic that acts against a wide range of disease-causing bacteria. These medications are used when a bacterial infection is suspected but the group of bacteria is unknown (also called empiric therapy) or when infection with multiple groups of bacteria is suspected. Although powerful, broad-spectrum antibiotics pose specific risks, particularly the disruption of native, normal bacteria and the development of antimicrobial resistance. Examples of commonly used broad-spectrum antibiotics are: Aminoglycosides (except for streptomycin), Ampicillin, Amoxicillin, Amoxicillin, clavulanic acid (Augmentin), Carbapenems (e.g. imipenem), Piperacillin, tazobactam, Quinolones (e.g. ciprofloxacin), Tetracyclines, Chloramphenicol, Ticarcillin, Trimethoprim and sulfamethoxazole (Bactrim). In veterinary medicine, examples are co-amoxiclav, (eg, in small animals), penicillin, streptomycin, oxytetracycline and potentiated sulfonamides.

Clauses: —

The invention, therefore, in one aspect provides the following Clauses that are directed to the treatment of a pathogenic bacterial infection using a programmed nuclease.

1. A method for treating a pathogenic bacterial infection in a human or animal subject caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a programmable nuclease that is programmed to cut the target site, wherein the subject is suffering from a further disease or condition other than the pathogenic bacterial infection and the method comprises administering a therapy to the subject for treating or preventing the further disease or condition,

wherein the nuclease treats the infection and the therapy is efficacious in the presence of the programmed nuclease to treat or prevent the disease or condition.

In an example, Clause 1 provides: —

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-PD-1 antibody optionally selected from pembrolizumab (or KEYTRUDA™) and nivolumab (or OPDIVO™); and
- (b) The cancer is selected from metastatic melanoma; renal cell carcinoma; bladder cancer; a solid tumour; non-small cell lung cancer (NSCLC); forehead and neck squamous cell carcinoma (HNSCC); Hodgkin’s lymphoma; a cancer that overexpresses PD-L1 and no mutations in EGFR or in ALK; colorectal cancer and hepatocellular carcinoma; and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-PD-L1 antibody optionally selected from atezolimumab (or TECENTRIQ™), avelumab (or BAVENCIO™) and durvalumab (or IMFINZI™); and
- (b) The cancer is selected from metastatic melanoma; renal cell carcinoma; a solid tumour; non-small cell lung cancer (NSCLC); forehead and neck squamous

cell carcinoma (HNSCC); Merkel cell carcinoma; Hodgkin's lymphoma; a cancer that overexpresses PD-L1 and no mutations in EGFR or in ALK; colorectal cancer and hepatocellular carcinoma; and

- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-CD52, antibody optionally alemtuzumab (or CAMPATH™); and
- (b) The cancer is B-cell chronic lymphocytic leukemia (CLL); and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-CD20 antibody, optionally ofatumumab (or ARZERRA™) or rituximab (or RITUXAN™); and
- (b) The cancer is B-cell chronic lymphocytic leukemia (CLL) (eg, refractory CLL) or non-Hodgkin lymphoma; and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the

subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-KIR antibody, optionally lirilumab; and
- (b) The cancer is optionally acute myeloid leukaemia or squamous cell carcinoma of the head and neck (SCCHN); and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer;

Wherein

- (a) The immunotherapy comprises administering to the patient an anti-CD19 CAR-T optionally selected from axicabtagene ciloleucel (or YESCARTA™) and tisagenlecleucel (or KYMRIA™); and
- (b) The cancer is selected from a B-cell lymphoma (eg, non-Hodgkin's lymphoma (NHL); diffuse large B-cell lymphoma (DLBCL); primary mediastinal large B-cell lymphoma; or high grade B-cell lymphoma); B-cell acute lymphoblastic leukaemia (ALL); or central nervous system lymphoma; and
- (c) The first bacteria are selected from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E coli*, *Salmonella* (eg, *S typhimurium*), *Clostridium difficile*, *Staphylococcus* (eg, *S aureus* or *S epidermis*), *Streptococcus* (eg, *S viridans* or *S thermophilus*), *Pneumococcus* and *Enterococcus* bacteria.

Alternatively, the CAR-T is an anti-CD30, CD38 or CD22 CAR-T. In an example the cancer is large B-cell lymphoma after at least two other kinds of treatment failed. In an example the cancer is high grade B-cell lymphoma and DLBCL arising from follicular lymphoma. In an example the cancer is relapsing/remitting B cell acute lymphoblastic leukaemia. In an example the cancer is primary central nervous system lymphoma.

In an example, the nuclease treats the infection without causing reduction in efficacy of the therapy. In an embodiment, "without causing reduction in efficacy of the therapy" means the efficacy of the therapy compared to a reduction caused in patients by the administration of a broad-spectrum antibiotic (or an antibiotic disclosed herein) that kills a plurality of different species, wherein the plurality comprises the first species. In an embodiment, "without causing reduction in efficacy of the therapy" means the efficacy of the therapy is reduced by no more than 70, 80, 90 or 95% compared to administration of the therapy in the absence of treatment of the pathogenic bacterial infection (or compared to therapy as typically achieved in patients suffering from the disease or condition and receiving said therapy therefor).

This may be assessed, for example, by determining the duration of progression-free survival of the subject or treatment of the disease or condition, or overall survival of the subject; and/or by determining a reduction in one or more symptoms of the disease or condition.

In an example, the infection is treated completely or substantially completely. In another example, the infection is reduced (eg, by at least 80, 90 or 95% as determined by a marker of the infection or a symptom thereof). A marker may, for example, be CFUs of bacteria of the first species or strain per ml of a blood sample taken from the patient after the method has been carried out, eg, within 24 hours of that method being carried out, eg, from 1-12 hours or 1-24 hours after carrying out the method or from 1-12 hours or 1-24 hours after administering a RNA or DNA encoding the RNA to programme the nuclease in the subject. For example, the RNA is a guide RNA and the nuclease is Cas (eg, a Cas3 or a Cas9). The reduction may be compared to a sample taken from the subject immediately prior to the commencement of the method. Alternatively, the sample may be a stool, saliva or urine sample.

In an example, the invention increases overall survival rate in a human subject (compared to median overall survival rate in humans suffering from the same cancer and receiving the same cancer therapy treatment (eg, administration of the same immune checkpoint inhibitor, such as nivolumab, pembrolizumab or another antibody disclosed herein)). In an example any composition, or other product of the invention herein is provided for use in such method of treatment.

In an example, the method is practised on a population of human subjects and the median overall survival rate for the population is 120-250% (eg, 150-200%) of the median overall survival rate in humans suffering from the same cancer and receiving the same cancer therapy treatment (eg, administration of the same immune checkpoint inhibitor, such as nivolumab, pembrolizumab or another antibody disclosed herein). In an example any composition, or other product of the invention herein is provided for use in such method of treatment.

A "pathogenic bacterial infection" is a health-threatening infection of the subject, for example, a life-threatening infection. In an embodiment, a pathogenic bacterial infection is an infection requiring hospitalization or parenteral antimicrobials. The infection may be an acute bacterial infection, such as a systemic infection or a localised infection. Bacterial pathogens often cause infection in specific areas of the body. Others are generalists. A pathogenic bacterial infection is contrasted with an infection of commensal bacteria, such as commensal gut bacteria; in this case the bacteria do not cause an immediate health- or life-threatening situation.

The infection (or symptom thereof) can be any of the following: —

Bacterial vaginosis: this is caused by bacteria that change the vaginal microbiota caused by an overgrowth of bacteria that crowd out the Lactobacilli species that maintain healthy vaginal microbial populations.

Bacterial meningitis: this is a bacterial inflammation of the meninges, that is, the protective membranes covering the brain and spinal cord.

Bacterial pneumonia: this is a bacterial infection of the lungs.

Urinary tract infection: this is predominantly caused by bacteria. Symptoms include the strong and frequent sensation or urge to urinate, pain during urination, and urine that is cloudy. The main causal agent is *Escheri-*

chia coli. Bacteria can ascend into the bladder or kidney and causing cystitis and nephritis.

Bacterial gastroenteritis: this is caused by enteric, pathogenic bacteria. These pathogenic species are usually distinct from the usually harmless bacteria of the normal gut flora. But a different strain of the same species may be pathogenic.

Bacterial skin infections: these include:

Impetigo, which is a highly contagious bacterial skin infection commonly seen in children. It is caused by *Staphylococcus aureus*, and *Streptococcus pyogenes*.

Erysipelas, which is an acute *streptococcus* bacterial infection of the deeper skin layers that spreads via with lymphatic system.

Cellulitis, which is a diffuse inflammation of connective tissue with severe inflammation of dermal and subcutaneous layers of the skin. Cellulitis can be caused by normal skin flora or by contagious contact, and usually occurs through open skin, cuts, blisters, cracks in the skin, insect bites, animal bites, burns, surgical wounds, intravenous drug injection, or sites of intravenous catheter insertion. In most cases it is the skin on the face or lower legs that is affected, though cellulitis can occur in other tissues.

In an example, the first bacteria are *Streptococcus* and the patient is suffering from chest infection, cellulitis or tonsillitis. In an example, the first bacteria are *Enterococcus* and the patient is suffering from bladder infection or septicaemia. In an example, the first bacteria are *Pseudomonas aeruginosa* and the patient is suffering from diarrhoea. In an example, the first bacteria are *E coli* and the patient is suffering from diarrhoea.

2. The method of Clause 1, wherein the subject is a cancer patient and the therapy is a cancer therapy.

3. The method of Clause 2, wherein the therapy comprises administration of a haematopoietic stem cell transplant, chemotherapeutic agent, immune checkpoint inhibitor, immune checkpoint agonist or an immune cell (eg, T-cell and/or NK cell) enhancer; adoptive cell therapy (eg, CAR-T therapy); radiation or surgery.

In an example, the therapy is immunotherapy. Examples of suitable immunotherapy are administration of adoptive cell therapy (eg, CAR-T therapy), an immune checkpoint inhibitor, an immune checkpoint agonist or an immune cell (eg, T-cell and/or NK cell) enhancer. For example, administration of an anti-CTLA4, PD-1, PD-L1, PD-L2, LAG3, OX40, CD28, BTLA, CD137, CD27, HVEM, KIR, TIM-3, VISTA, ICOS, GITR, TIGIT or SIRPa antibody, such as administration of an antibody selected from ipilimumab (or YERVOY™), tremelimumab, nivolumab (or OPDIVO™), pembrolizumab (or KEYTRUDA™), pidilizumab, BMS-936559, durvalumab and atezolizumab, or a CAR-T therapy such as axicabtagene ciloleucel (Yescarta™) or tisagenlecleucel (Kymriah™).

In an example, the immune enhancer comprises an interleukin-2 (IL-2) or fragment or deletion mutant thereof.

In an example, the surgery comprises the removal of necrotic or cancerous tissue.

In an example, the chemotherapy comprises administration of a platinum-containing chemotherapy drug. In an example, the chemotherapy comprises administration of gefitinib.

In an example, the therapy comprises administering Cyclophosphamide, methotrexate and 5-fluorouracil (CMF); or doxorubicin and cyclophosphamide (AC); docetaxel, doxorubicin and cyclophosphamide (TAC); or doxorubicin, bleomycin, vinblastine and dacarbazine (ABVD); or mus-

tine, vincristine, procarbazine and prednisolone (MOPP); cyclophosphamide, doxorubicin, vincristine and prednisolone (CHOP); bleomycin, etoposide and cisplatin (BEP); epirubicin, cisplatin and 5-fluorouracil (ECF); or epirubicin, cisplatin and capecitabine (ECX); methotrexate, vincristine, doxorubicin and cisplatin (MVAC); cyclophosphamide, doxorubicin and vincristine (CAV); or 5-fluorouracil, folinic acid and oxaliplatin (FOLFOX).

In an example, the cancer is breast cancer and the therapy comprises administering CMF or AC. In an example, the cancer is Hodgkin's lymphoma and the therapy comprises administering TAC, ABVD or MOPP. In an example, the cancer is Non-Hodgkin's lymphoma and the therapy comprises administering CHOP. In an example, the cancer is germ cell cancer and the therapy comprises administering BEP. In an example, the cancer is stomach cancer and the therapy comprises administering ECF or ECX. In an example, the cancer is bladder cancer and the therapy comprises administering MVAC. In an example, the cancer is lung cancer and the therapy comprises administering CAV. In an example, the cancer is colorectal cancer and the therapy comprises administering FOLFOX.

4. The method of Clause 3, wherein the therapy is an immune checkpoint inhibitor antibody.

Optionally the antibody is an anti-CTLA4, PD-1, PD-L1, PD-L2, LAG3, OX40, CD28, BTLA, CD137, CD27, HVEM, KIR, TIM-3, VISTA, ICOS, GITR, TIGIT or SIRPa antibody. In an example, the antibody is an anti-PD-1 antibody. In an example, the antibody is an anti-PD-L1 antibody. In an example, the antibody is an anti-CTLA4 antibody.

5. The method of Clause 3, wherein the therapy is administration of an antibody selected from ipilimumab (or YERVOY™), tremelimumab, nivolumab (or OPDIVO™), pembrolizumab (or KEYTRUDA™), pidilizumab, BMS-936559, durvalumab and atezolizumab.

Optionally, the antibody (eg, anti-PD-L1 antibody) is administered with an anti-CTLA4 antibody (eg, ipilimumab or tremelimumab).

In an example, the an anti-PD-1 antibody herein is selected from nivolumab, pembrolizumab, pidilizumab, OPDIVO®, KEYTRUDA®, AMP-514, REGN2810, CT-011, BMS 936559, MPDL3280A and AMP-224.

In an example, the an anti-CTLA4 antibody herein is selected from tremelimumab, YERVOY® and ipilimumab.

In an example the therapy is administration of an anti-KIR antibody, eg, lirilumab.

In an example, the checkpoint inhibitor is selected from an inhibitor of CTLA-4, PD-1, PD-L1, PD-L2, LAG-3, BTLA, B7H3, B7H4, TIM3, KIR, or A2aR. In certain aspects, the immune checkpoint inhibitor is a human programmed cell death 1 (PD-1) axis-binding antagonist. In some aspects, the PD-1 axis-binding antagonist is selected from the group consisting of a PD-1 binding antagonist, a PD-L1-binding antagonist and a PD-L2-binding antagonist. In certain aspects, the PD-1 axis-binding antagonist is a PD-1-binding antagonist. In some aspects, the PD-1-binding antagonist inhibits the binding of PD-1 to PD-L1 and/or PD-L2.

In some embodiments, the immune checkpoint inhibitor is a PD-L1 antagonist such as durvalumab, also known as MEDI4736, atezolizumab, also known as MPDL3280A, or avelumab, also known as MSB00010118C. In certain aspects, the immune checkpoint inhibitor is a PD-L2 antagonist such as rHlgM12B7. In some aspects, the immune checkpoint inhibitor is a LAG-3 antagonist such as IMP321

or BMS-986016. The immune checkpoint inhibitor may be an adenosine A2a receptor (A2aR) antagonist such as PBF-509.

In some embodiments, the antibody described herein (such as an anti-PD-1 antibody, an anti-PD-L1 antibody, or an anti-PD-L2 antibody) further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic, CHO, Cos or HEK cells. In a still further specific aspect the minimal effector function results from an "effector-less Fc mutation" or aglycosylation.

For example, the therapy comprises a hemopoietic stem cell transplant, eg, a bone marrow transplant (such as when the patient is a cancer patient, eg, a blood cancer or leukaemia patient).

For example, the therapy comprises a stem cell transplant, a skin graft, or an organ transplant, eg, a heart, liver, kidney or lung transplant.

6. The method of Clause 1 or 2, wherein the therapy is a tissue, organ or cell transplant.

7. The method of any preceding Clause, wherein the treatment of the bacterial infection is carried out simultaneously with the administration of the therapy to the subject.

8. The method of any one of Clauses 1 to 6, wherein the treatment of the bacterial infection is carried out immediately before administering the therapy to the subject.

In an example, the treatment of the bacterial infection is carried out no more than 7, 6, 5, 4, 3, 2, or 1 day, or 24, 12, 6, 5, 4, 3, 2, 1 or 0.5 hours before the therapy of the further disease or condition.

In an example, the treatment of the bacterial infection is carried out no more than 7, 6, 5, 4, 3, 2, or 1 day, or 24, 12, 6, 5, 4, 3, 2, 1 or 0.5 hours after the therapy of the further disease or condition.

The treatment of the infection and the administration of the therapy may be carried out simultaneously or sequentially.

9. The method of any one of Clauses 1 to 6, wherein the treatment of the bacterial infection is carried out immediately after administering the therapy to the subject.

10. The method of any preceding Clause, wherein the method comprises administering to the subject a RNA (eg, a gRNA) or a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in first bacteria comprised by the subject, thereby killing the first bacteria.

The RNA or nucleic acid is, for example, administered to the subject or patient orally, by IV injection, by subcutaneous injection or by inhalation.

11. The method of any preceding Clause, comprising administering a vector (eg, phage or plasmids) to the subject, wherein the vector encodes the programmable nuclease.

The nuclease is, for example, administered to the subject or patient orally, by IV injection, by subcutaneous injection or by inhalation.

12. The method of any one of Clauses 1 to 10, wherein the programmable nuclease is an endogenous nuclease (eg, Cas nuclease) of the first cells.

13. The method of any preceding Clause, wherein the efficacy of the therapy in the presence of the programmed nuclease is greater than the efficacy of the therapy in the presence of a broad-spectrum antibiotic.

In an example, the efficacy being greater is assessed by determining the duration of progression-free survival or treatment of the disease or condition; and/or by determining a reduction in one or more symptoms of the disease or condition. For example, this determination is compared to an analogous determination in a patient suffering from the disease or condition as well as the bacterial infection and being treated with the therapy and the antibiotic (rather than the nuclease killing of first bacteria as per the invention).

14. The method of any preceding Clause, wherein the efficacy of the therapy in the presence of the programmed nuclease is greater than the efficacy of the therapy in the presence of an antibiotic selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbopristin; teicoplanin; cephalosporin; carbapenem; fluoroquinolone; aminoglycoside; colistin; erythromycin; clindamycin; beta-lactam; macrolide; amoxicillin; azithromycin; penicillin; ceftriaxone; azithromycin; ciprofloxacin; isoniazid (INH); rifampicin (RMP); amikacin; kanamycin; capreomycin; trimethoprim; itrofurantoin; cefalexin; amoxicillin; metronidazole (MTZ); cefixime; tetracycline; and meropenem.

15. The method of any preceding Clause, wherein the first bacteria is selected from (i) *Staphylococcus aureus* that is resistant to an antibiotic selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbopristin and teicoplanin; (ii) *Pseudomonas aeruginosa* that is resistant to an antibiotic selected from cephalosporins, carbapenems, fluoroquinolones, aminoglycosides and colistin; (iii) *Klebsiella* species that is resistant to carbapenem; (iv) *Streptococcus* species that is resistant to an antibiotic selected from erythromycin, clindamycin, beta-lactam, macrolide, amoxicillin, azithromycin and penicillin; (v) *Salmonella* species that is resistant to an antibiotic selected from ceftriaxone, azithromycin and ciprofloxacin; (vi) *Shigella* species that is resistant to ciprofloxacin or azithromycin; (vii) *Mycobacterium tuberculosis* that is resistant to an antibiotic selected from Resistance to isoniazid (INH), rifampicin (RMP), fluoroquinolone, amikacin, kanamycin, capreomycin and azithromycin; (viii) *Enterococcus* species that is resistant to vancomycin; (ix) Enterobacteriaceae species that is resistant to an antibiotic selected from cephalosporin and carbapenem; (x) *E coli* that is resistant to an antibiotic selected from trimethoprim, itrofurantoin, cefalexin and amoxicillin; (xi) *Clostridium* species that is resistant to metronidazole (MTZ), fluoroquinolone or carbapenem; (xii) *Neisseria gonorrhoea* that is resistant to an antibiotic selected from cefixime, ceftriaxone, azithromycin and tetracycline; (xiii) *Acinetobacter baumannii* that is resistant to an antibiotic selected from beta-lactam, meropenem and carbapenem; and (xiv) *Campylobacter* species that is resistant to ciprofloxacin or azithromycin.

16. The method of any preceding Clause, wherein the treatment of the infection treats or prevents in the subject a condition selected from vaginosis, meningitis, pneumonia, urinary tract infection, cystitis, nephritis,

gastroenteritis, a skin infection, impetigo, erysipelas, dental infection and cellulitis.

17. The method of any preceding Clause, wherein the treatment of the infection treats or prevents septicaemia or sepsis in the subject.

In an example, the infection is a bloodstream infection.

18. The method of any preceding Clause, wherein the further disease or condition is a cancer; autoimmune disease or condition; viral infection or GI tract disease or condition. In an example, the cancer is metastatic. In an example, the cancer is melanoma. In an example, the cancer is a solid tumour with mismatch repair deficiency or microsatellite instability. In an example, the cancer is NSCLC. In an example, the cancer is HNSCC. In an example, the cancer is Hodgkin's lymphoma. In an example, the cancer is urothelial cancer. In an example, the cancer is lung cancer. In an example, the cancer is head and neck cancer. In an example, the cancer is head cancer. In an example, the cancer is neck cancer. In an example, the viral infection is a HIV, CMV or RSV infection.

19. The method of any preceding Clause, wherein the subject comprises bacteria (second bacteria) of one or more strains or species that are different to the first strain or species, wherein the genomes of the second bacteria do not comprise the target site, wherein the genomes of the second bacteria are not cut by the programmed nuclease in the subject, whereby second bacteria survive in the presence of the programmed nuclease in the patient; and wherein the therapy is efficacious in the presence of the second bacteria.

20. The method of Clause 19, wherein reduction in the second bacteria in patients (eg, in the gut microbiome) is associated with reduced efficacy of the therapy.

Optionally, the therapy is efficacious in the presence of the second bacteria in the gut of the subject.

Optionally, the first and/or second bacteria are present in the gut of the subject immediately prior to carrying out the method.

Optionally, the first and/or second bacteria are present in the blood of the subject immediately prior to carrying out the method.

Optionally, the first bacteria are present in the blood of the subject and the second bacteria are present in the gut of the subject immediately prior to carrying out the method.

Optionally, the first bacteria are present in the gut of the subject and the second bacteria are present in the blood of the subject immediately prior to carrying out the method.

Optionally, first bacteria in the blood of the subject is killed.

Optionally, the bacteria are gram positive bacteria. Optionally, the bacteria are gram negative bacteria.

Optionally, the first and second bacteria are capable of being killed by the same antibiotic.

Optionally, the method does not comprise administering the antibiotic to the subject. In an example, the antibiotic is selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbopristin; teicoplanin; cephalosporin; carbapenem; fluoroquinolone; aminoglycoside; colistin; erythromycin; clindamycin; beta-lactam; macrolide; amoxicillin; azithromycin; penicillin; ceftriaxone; azithromycin; ciprofloxacin; isoniazid (INH); rifampicin (RMP); amikacin; kanamycin; capreomycin; trimethoprim; itrofurantoin; cefalexin; amoxicillin; metronidazole (MTZ); cefixime; tetracycline; and meropenem. In an example, the antibiotic is selected from Aminoglycosides, Ampicillin, Amoxicillin, Amoxicillin or clavulanic acid, Carbapenems (eg. imipe-

nem), Piperacillin or tazobactam, Quinolones (e.g. ciprofloxacin), Tetracyclines, Chloramphenicol, Ticarcillin, Trimethoprim or sulfamethoxazole, penicillin, streptomycin, oxytetracycline and potentiated sulfonamides. In an example, the first bacteria are resistant to an antibiotic selected from Aminoglycosides, Ampicillin, Amoxicillin, Amoxicillin or clavulanic acid, Carbapenems (e.g. imipenem), Piperacillin or tazobactam, Quinolones (e.g. ciprofloxacin), Tetracyclines, Chloramphenicol, Ticarcillin, Trimethoprim or sulfamethoxazole, penicillin, streptomycin, oxytetracycline and potentiated sulfonamides. In an alternative, the antibiotic is selected from a beta-lactam, fluoroquinolone and macrolide.

Optionally, the first and second bacteria are bacteria of the same species, but are different strains of the species.

Optionally, the first and second bacteria are bacteria of the same genus, but are bacteria of different species of the genus.

Optionally, the first and second bacteria are bacteria of the same family, but are bacteria of different genera of the family.

Optionally, the first and second bacteria are gram positive bacteria.

Optionally, the first and second bacteria are gram-negative bacteria.

Optionally, the therapy is efficacious in the presence of the second bacteria.

Optionally, reduction in the second bacteria in patients is associated with reduced efficacy of the therapy. Optionally, reduction in the second bacteria in patients reduces efficacy of the therapy.

Optionally, the presence of the second bacteria in patients is associated with enhanced efficacy of the therapy. Optionally, the presence of the second bacteria in patients enhances efficacy of the therapy. For example, enhanced efficiency is efficiency compared to therapy in the absence or a reduced presence of the second bacteria, such as in the presence of an antibiotic that kills the second bacteria.

In an example, the therapy is efficacious in the presence of the second bacteria, wherein the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the progression of the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein disease-free progression of the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the duration of the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the severity of the disease or condition (or a symptom thereof) is reduced in the subject by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the disease or condition (or a symptom thereof) is reduced for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 21 or 28 days or for at least 1, 2, 3, 4, 5, 6 or 12 months in the patient by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the disease or condition (or a symptom thereof) is treated for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 21 or 28 days or for at least 1, 2, 3, 4

5, 6 or 12 months in the patient by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%. In an example, the therapy is efficacious in the presence of the second bacteria, wherein the disease or condition (or a symptom thereof) is undetectable for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 21 or 28 days or for at least 1, 2, 3, 4, 5, 6 or 12 months in the patient by at least 20, 30, 40, 50, 60, 70, 80, 90 or 95%.

21. The method of Clause 19 or 20, wherein the second bacteria are selected from the group consisting of *Akkermansia*, *Alistipes*, *Bacteroides*, *Barnesiella*, *Bifidobacterium*, *Clostridium*, *Collinsella*, *Enterococcus*, *Fusobacterium*, *Lactobacillus*, *Propionibacterium*, *Ruminococcus*, *Segmented filamentous bacteria* (SFB); *Veillonella*, *Prevotella*, *Escherichia* and *Streptococcus* bacteria.

In an example the second bacteria bacteria that produce short chain fatty acids (eg, butyrate-producing bacteria). In particular aspects, the species of bacteria produce butyrate. For example, the second bacteria are Clostridiales. The Clostridiales bacteria may be substantially or include bacteria in spore form. In particular aspects, the second bacteria are of the family Ruminococcaceae, Christensenellaceae, Clostridiaceae or Coriobacteriaceae. In some embodiments, the Clostridiales (eg, *Clostridium*) bacteria comprise a first family and a second family. In some embodiments, the first family is selected from the group consisting of Ruminococcaceae, Christensenellaceae, Clostridiaceae and Coriobacteriaceae, and the second family is not identical to the first family. In an example, the second bacteria are *Faecalibacterium prausnitzii*, *Ruminococcus albus*, *Ruminococcus bromii*, *Ruminococcus callidus*, *Ruminococcus flavefaciens*, *Ruminococcus champanellensis*, *Ruminococcus faecis*, *Ruminococcus gauvreauii*, *Ruminococcus gnavus*, *Ruminococcus hanseni*, *Ruminococcus hydrogenotrophicus*, *Ruminococcus lactaris*, *Ruminococcus luti*, *Ruminococcus obeum*, *Ruminococcus palustris*, *Ruminococcus pasteurii*, *Ruminococcus productus*, *Ruminococcus schinkii*, *Ruminococcus torques*, *Subdoligranulum variabile*, *Butyrivibrio fibrisolvens*, *Roseburia intestinalis*, *Anaerostipes caccae*, *Blautia obeum*, *Eubacterium nodatum* or *Eubacterium oxidoreducens*. In particular aspects, the second bacteria are *Faecalibacterium prausnitzii*. In an example the second bacteria are Firmicutes.

In certain embodiments, the first bacteria are *Bacteroidia* or *Prevotellaceae*, eg, *Bacteroidetes* or *Bacteroides*.

In an embodiment, the treatment results in or maintains a microbiome (eg, gut and/or blood microbiome) of the subject, which is beneficial for the immune checkpoint inhibition or other therapy. In an example, the microbiome comprises a high relative abundance of one or more bacterial species from the phylum Firmicutes, class Clostridia, order Clostridiales, family Ruminococcaceae, genus *Ruminococcus*, genus *Hydrogenoanaerobacterium*, genus *Faecalibacterium*, phylum Actinobacteria, class Coriobacteriia, order Coriobacteriales, family Coriobacteriaceae, domain Archaea, phylum Cyanobacteria, phylum Euryarchaeota or family Christensenellaceae. Additionally or alternatively, the microbiome comprises a low relative abundance of bacteria from the genus *Dialister*, family Veillonellaceae, phylum Bacteroidetes, class Bacteroida, order Bacteroidales or family Prevotellaceae. Accordingly, a favorable microbial profile would have a higher relative abundance of one or more bacterial species from the phylum Firmicutes, class Clostridia, order Clostridiales, family Ruminococcaceae, genus *Ruminococcus*, genus *Hydrogenoanaerobacterium*, phylum Actinobacteria, class Coriobacteriia, order Coriobacteriales, family Coriobacteriaceae, domain Archaea, phy-

lum Cyanobacteria, phylum Euryarchaeota or family Christensenellaceae, and/or has a decreased abundance of one or more bacterial species from genus *Dialister*, family Veillonellaceae, phylum Bacteroidetes, class Bacteroidia, order Bacteroidales and/or family Prevotellaceae.

For example, the microbiome comprises a higher relative abundance of Firmicutes compared to Bacteroidetes, Bacteroidia, Bacteroidales or Prevotellaceae. For example, the microbiome comprises a higher relative abundance of Firmicutes compared to Bacteroidetes, Bacteroidia, Bacteroidales and Prevotellaceae.

Optionally, the second bacteria are selected from the group consisting of *Akkermansia muciniphila*; *Alistipes shahii*; *Bacteroides fragilis*; *Bacteroides uniformis*; *Barnesiella intestinihominis*; *Bacteroides dorei*; *Bifidobacterium adolescentis*; *Bifidobacterium breve*; *Bifidobacterium longum*; *Clostridium orbiscindens*; *Clostridium novyi*; *Clostridium perfringens*; *Collinsella aerofaciens*; *Enterococcus hirae*; *Fusobacterium nucleatum*; *Lactobacillus casei* Shirota; *L. casei* AO47; *Lactobacillus rhamnosus*; *Propionibacterium granulosum*; *Ruminococcus gnavus*; *Segmented filamentous bacteria* (SFB); *Veillonella*; *Lactobacilli*; *Bacteroides*; *Clostridia*; *Prevotella*; *E. coli* Nissle; *Lactobacillus plantarum*; *Lactobacillus delbrueckii* (eg, subsp. *Bulgarius*); *Lactobacillus paracasei*; *Lactobacillus acidophilus*; *Bifidobacterium infantis*; and *Streptococcus salivarius* (eg, subsp. *Thermophilus*). See “The microbiome in cancer immunotherapy: Diagnostic tools and therapeutic strategies”; Laurence Zitvogel et al; Science 23 Mar. 2018: Vol. 359, Issue 6382, pp. 1366-1370; DOI: 10.1126/science.aar6918.

In an example, the second bacteria are commensal bacteria in humans.

In an example, the first bacteria are comprised by gut microbiota, skin microbiota, oral cavity microbiota, throat microbiota, hair microbiota, armpit microbiota, vaginal microbiota, rectal microbiota, anal microbiota, ocular microbiota, nasal microbiota, tongue microbiota, lung microbiota, liver microbiota, kidney microbiota, genital microbiota, penile microbiota, scrotal microbiota, mammary gland microbiota, ear microbiota, urethra microbiota, labial microbiota, organ microbiota or dental microbiota.

In an example, the second bacteria are comprised by gut microbiota, skin microbiota, oral cavity microbiota, throat microbiota, hair microbiota, armpit microbiota, vaginal microbiota, rectal microbiota, anal microbiota, ocular microbiota, nasal microbiota, tongue microbiota, lung microbiota, liver microbiota, kidney microbiota, genital microbiota, penile microbiota, scrotal microbiota, mammary gland microbiota, ear microbiota, urethra microbiota, labial microbiota, organ microbiota or dental microbiota.

In an example, the first and/or second bacteria are blood-borne bacteria.

22. The method of any preceding Clause, wherein the first bacteria are selected from the group consisting of *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Helicobacter*, *Legionella*, *Heamophilus*, *Ghonorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria.

H pylori has been implicated in gastric cancer and gastric ulcers. Thus, in an example, the first bacteria are *H pylori* and optionally the disease is a cancer, such as gastric cancer. In an embodiment, the therapy is chemotherapy or therapy with an immune checkpoint inhibitor (eg, an antibody). In an example, the first bacteria are *H pylori* and the disease is gastric ulcer(s). In an embodiment, triple therapy for gastric ulcers is administered to the subject.

In an example, the first bacteria are Gram-negative bacteria and optionally the infection is a blood infection. In an example, the first bacteria are selected from *E. coli*, *P. aeruginosa* and *K. pneumoniae*, and optionally the infection is a blood infection.

23. The method of Clause 22, wherein the first bacteria are selected from the group consisting of *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Helicobacter pylori*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* bacteria.

In an example, the subject has been administered an immunosuppressant drug, or is on a course of an immunosuppressant drug, eg, a steroid, such as a corticosteroid.

24. A programmable nuclease for use in the method of any preceding Clause.

25. The method or nuclease of any preceding Clause, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.

26. A CRISPR/Cas system comprising a nuclease according to Clause 24 or 25 for use in the method of any one of Clauses 1 to 23, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs (gRNAs) or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of first bacteria.

27. A guide RNA or a DNA encoding a guide RNA for use in the system of Clause 26.

28. A guide RNA or a DNA encoding a guide RNA for use in the method of treating a pathogenic bacterial infection according to any one of Clauses 1 to 23, wherein the guide RNA is capable of programming the nuclease, wherein the nuclease is a Cas nuclease (eg, a Cas9, Cas3, Cas13, CasX, CasY or Cpf1 nuclease).

29. A nucleic acid vector comprising the guide RNA or DNA recited in any one of Clauses 26 to 28.

30. A nucleic acid vector encoding the nuclease of Clause 24 or 25 and optionally the guide RNA of Clause 29.

31. The vector of Clause 29 or 30 wherein the vector is a phage, phagemid plasmid (eg, conjugative plasmid) or transposon.

The phage are capable of infecting first bacteria and the phagemids are capable of producing such phage in the presence of a helper phage.

32. A pharmaceutical composition comprising a first nucleic acid vector (or a plurality thereof) encoding the nuclease of Clause 24 or 25 and a second nucleic acid vector (or a plurality thereof) encoding the guide RNA of Clause 29, the composition further comprising a pharmaceutically acceptable diluent, excipient or carrier.

33. A pharmaceutical composition comprising the CRISPR/Cas system of claim 26 and a pharmaceutically acceptable diluent, excipient or carrier.

34. A pharmaceutical composition comprising the vector of claim 31 and a pharmaceutically acceptable diluent, excipient or carrier.

Preventing a disease or condition herein may, for example, be reducing the risk of the disease or condition in the subject or patient.

In an alternative, instead of first bacteria, the infection is caused by first archaea and in this embodiment all of the features of the method and other configurations of the

invention relating to killing first bacteria instead relate mutatis mutandis to killing first archaea.

In an embodiment, the method comprises carrying out the method of treating an acute microbial infection as described herein, and thus features of that method as described herein are combinable with the present method of treating a pathogenic bacterial infection (ie, where the pathogenic bacterial infection is the acute microbial infection in the first method). In an embodiment, the method comprises carrying out the method of durably treating a microbial infection as described herein, and thus features of that method as described herein are combinable with the present method of treating a pathogenic bacterial infection (ie, where the pathogenic bacterial infection is the microbial infection in the first method). Any of the optional features of the first method herein may apply mutatis mutandis to the present method of treating a pathogenic bacterial infection.

Aspects: —

Thus, the invention provides the following Aspects, which are optional features of Clauses above: —

1. The method of any one of Clauses 1-23, wherein the infection is reduced at least 100-fold by the first 30 minutes of carrying out step (b). Optionally, the infection is reduced at least 1000-fold by the first 30 minutes of carrying out step (b). Optionally, the reduction in infection persists for 30 minutes immediately after the first 30 minutes of carrying out step (b). For example, the reduction can be assessed by determining the difference in the number of bacteria of the first species or strain in (i) a sample taken from the subject (eg, a blood sample) immediately before commencement of the method and (ii) a sample (of the same type as the sample of (i), eg, a blood sample) taken from the subject at 30 minutes of the treatment. For example, the samples may be assessed for the difference in colony forming units (CFU)/ml sample, eg, when the samples have been plated on agar in respective petri dishes and incubated under identical conditions. Another example may use microscopic counting of bacteria in samples, or other routine methods known to the skilled addressee.
2. The method of any one of Clauses 1-23, wherein blood infection of the subject by the first bacteria is reduced at least 100- or 1000-fold by the first 30 minutes of carrying out step (b).
3. The method of Aspect 2, wherein the blood is infected with from 10^5 to 10^{12} (eg, 10^7 to 10^{12}) CFU/ml of the first bacteria immediately before the treatment.
4. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the method comprises administering to the subject a nucleic acid (eg, a RNA) and nuclease, wherein the nucleic acid complexes with the nuclease to program the nuclease to cut the target site in the first bacteria comprised by the subject.
5. The method of Aspect 4, wherein the nuclease is administered simultaneously or sequentially with the nucleic acid to the subject.
6. The method of Aspect 4, wherein the subject comprises the nuclease prior to administration of the nucleic acid to the subject.
7. The method of any one of Aspects 4 to 6, wherein a plurality phage are administered to the subject, wherein each phage comprises a copy of the nucleic acid, wherein the phage infect first bacteria comprised by the subject to deliver thereto the nucleic acid.
8. The method of Aspect 7, wherein the ratio of administered phage: first bacteria comprised by the subject is

from 10 to 150. For example, the ratio is from 10 to 100, ie, a multiplicity of infection (MOI) of from 10 to 100.

The ratio can be determined, for example, using a sample (eg, a blood or gut sample) from a human or animal subject immediately before the treatment and determining the number of bacteria per ml of blood or gut sample. The amount of phage to be administered can then be worked out according to the determination using the sample.

9. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the infection is an infection of the lungs, brain, skin, abdomen or urinary tract.

10. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the subject has undergone surgery, is on an immunosuppressant medication, suffering from burns, suffering from diabetes, suffering from cancer or is suffering from a chronic disease.

11. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the subject is a human over 65 years of age or is a pediatric patient.

12. The method of any one of Clauses 1-23 or any preceding Aspect, wherein the method treats or prevents sepsis in the subject.

13. The method of Clause 12, wherein at the start of the treatment, the subject (eg, a human) has a temperature of $<36^\circ\text{C}$. or $>38^\circ\text{C}$.; a heart rate of $>90/\text{min}$, a respiratory rate of >20 breaths/min or $\text{PaCO}_2 < 4.3$ kPa; and white blood cell count of $<4000/\text{mm}^3$ or $>12,000/\text{mm}^3$.

14. The method of Clause 12 or 13, wherein at the start of the treatment, the subject (eg, a human) has presence of two or more of the following: abnormal body temperature, abnormal heart rate, abnormal respiratory rate, abnormal blood gas and abnormal white blood cell count.

Immune Checkpoint Modulation

Immune checkpoints of the invention either turn up a signal (e.g., co-stimulatory molecules) or turn down a signal. Inhibitory immune checkpoint molecules that may be targeted by immune checkpoint modulation in the invention include adenosine A2A receptor (A2AR), B7-H3 (also known as CD276), B and T lymphocyte attenuator (BTLA), cytotoxic T-lymphocyte-associated protein 4 (CTLA-4, also known as CD 152), indoleamine 2,3-dioxygenase (IDO), killer-cell immunoglobulin (KIR), lymphocyte activation gene-3 (LAG3), programmed death 1 (PD-1), T-cell immunoglobulin domain and mucin domain 3 (TIM-3) and V-domain Ig suppressor of T cell activation (VISTA). In particular, the immune checkpoint inhibitors target the PD-1 axis and/or CTLA-4.

The immune checkpoint inhibitors may be drugs such as small molecules, recombinant forms of ligand or receptors, or, antibodies, such as human antibodies (e.g., WO2015016718; Pardoll, Nat Rev Cancer, 12(4): 252-64, 2012; both incorporated herein by reference). Known inhibitors of the immune checkpoint proteins or analogs thereof may be used, in particular chimerised, humanised or human forms of antibodies may be used. As the skilled person will know, alternative and/or equivalent names may be in use for certain antibodies mentioned in the present disclosure. Such alternative and/or equivalent names are interchangeable in the context of the present invention. For example it is known that lambrolizumab is also known under the alternative and equivalent names MK-3475 and pembrolizumab.

It is contemplated that any of the immune checkpoint inhibitors that are known in the art to stimulate immune responses may be used. This includes inhibitors that directly

or indirectly stimulate or enhance antigen-specific T-lymphocytes. These immune checkpoint inhibitors include, without limitation, agents targeting immune checkpoint proteins and pathways involving PD-L2, LAG3, BTLA, B7H4 and TIM3. For example, LAG3 inhibitors known in the art include soluble LAG3 (IMP321, or LAG3-Ig disclosed in WO2009044273) as well as mouse or humanized antibodies blocking human LAG3 (e.g., IMP701 disclosed in WO2008132601), or fully human antibodies blocking human LAG3 (such as disclosed in EP 2320940). Another example is provided by the use of blocking agents towards BTLA, including without limitation antibodies blocking human BTLA interaction with its ligand (such as 4C7 disclosed in WO2011014438). Yet another example is provided by the use of agents neutralizing B7H4 including without limitation antibodies to human B7H4 (disclosed in WO 2013025779, and in WO2013067492) or soluble recombinant forms of B7H4 (such as disclosed in US20120177645). Yet another example is provided by agents neutralizing B7-H3, including without limitation antibodies neutralizing human B7-H3 (e.g. MGA271 disclosed as BRCA84D and derivatives in US 20120294796). Yet another example is provided by agents targeting TIM3, including without limitation antibodies targeting human TIM3 (e.g. as disclosed in WO 2013006490 A2 or the anti-human TIM3, blocking antibody F38-2E2 disclosed by Jones et al, J Exp Med. 2008; 205(12):2763-79).

A. PD-1 Axis Antagonists

T cell dysfunction or anergy occurs concurrently with an induced and sustained expression of the inhibitory receptor, programmed death 1 polypeptide (PD-1). Thus, therapeutic targeting of PD-1 and other molecules which signal through interactions with PD-1, such as programmed death ligand 1 (PD-L1) and programmed death ligand 2 (PD-L2) is provided herein. PD-L1 is overexpressed in many cancers and is often associated with poor prognosis (Okazaki T et al, Intern. Immun. 2007 19(7):813). Thus, improved methods of treating cancer by inhibiting the PD-L1/PD-1 interaction in combination with modulating the microbiome is provided herein.

For example, PD-1 axis binding antagonists include a PD-1 binding antagonist, a PD-L1 binding antagonist and a PD-L2 binding antagonist. Alternative names for "PD-1" include CD279 and SLEB2. Alternative names for "PD-L1" include B7-H1, B7-4, CD274, and B7-H. Alternative names for "PD-L2" include B7-DC, Btdc, and CD273. In some embodiments, PD-1, PD-L1, and PD-L2 are human PD-1, PD-L1 and PD-L2.

In some embodiments, the PD-1 binding antagonist is a molecule that inhibits the binding of PD-1 to its ligand binding partners. In a specific aspect, the PD-1 ligand binding partners are PD-L1 and/or PD-L2. In another embodiment, a PD-L1 binding antagonist is a molecule that inhibits the binding of PD-L1 to its binding partners. In a specific aspect, PD-L1 binding partners are PD-1 and/or B7-1. In another embodiment, the PD-L2 binding antagonist is a molecule that inhibits the binding of PD-L2 to its binding partners. In a specific aspect, a PD-L2 binding partner is PD-1. The antagonist may be an antibody, an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide. Exemplary antibodies are described in U.S. Pat. Nos. 8,735,553, 8,354,509, and 8,008,449, all incorporated herein by reference. Other PD-1 axis antagonists for use in the methods provided herein are known in the art such as described in U.S. Patent Application No. US20140294898, US2014022021, and US20110008369, all incorporated herein by reference.

In some embodiments, the PD-1 binding antagonist is an anti-PD-1 antibody {e.g., a human antibody, a humanized antibody, or a chimeric antibody}. In some embodiments, the anti-PD-1 antibody is selected from the group consisting of nivolumab, pembrolizumab, and CT-011. In some embodiments, the PD-1 binding antagonist is an immunoadhesin (e.g., an immunoadhesin comprising an extracellular or PD-1 binding portion of PDL1 or PD-L2 fused to a constant region (e.g., an Fc region of an immunoglobulin sequence). In some embodiments, the PD-1 binding antagonist is AMP-224. Nivolumab, also known as MDX-1106-04, MDX-1106, ONO-4538, BMS-936558, and OPDIVO®, is an anti-PD-1 antibody described in WO2006/121168. Pembrolizumab, also known as MK-3475, Merck 3475, lambrolizumab, KEYTRUDA®, and SCH-900475, is an anti-PD-1 antibody described in WO2009/114335. CT-011, also known as hBAT or hBAT-1, is an anti-PD-1 antibody described in WO2009/101611. AMP-224, also known as B7-DCIg, is a PD-L2-Fc fusion soluble receptor described in WO2010/027827 and WO2011/066342. Additional PD-1 binding antagonists include pidilizumab, also known as CT-011, MEDI0680, also known as AMP-514, and REGN2810.

In some embodiments, the immune checkpoint inhibitor is a PD-L1 antagonist such as durvalumab, also known as MEDI4736, atezolizumab, also known as MPDL3280A, or avelumab, also known as MSB00010118C. In certain aspects, the immune checkpoint inhibitor is a PD-L2 antagonist such as rHIgM12B7. In some aspects, the immune checkpoint inhibitor is a LAG-3 antagonist such as, but not limited to, IMP321, and BMS-986016. The immune checkpoint inhibitor may be an adenosine A2a receptor (A2aR) antagonist such as PBF-509.

In some embodiments, any antibody described herein (such as an anti-PD-1 antibody, an anti-PD-L1 antibody, or an anti-PD-L2 antibody) further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic cells. In a still further specific aspect the minimal effector function results from an "effector-less Fc mutation" or aglycosylation. Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylglucosamine, galactose, or xylose to a hydroxy amino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxy lysine may also be used. Removal of glycosylation sites from an antibody is conveniently accomplished by altering the amino acid sequence such that one of the above-described tripeptide sequences (for N-linked glycosylation sites) is removed. The alteration may be made by substitution of an asparagine, serine or threonine residue within the glycosylation site another amino acid residue (e.g., glycine, alanine or a conservative substitution).

The antibody or antigen binding fragment thereof, may be made using methods known in the art, for example, by a process comprising culturing a host cell containing nucleic acid encoding any of the previously described anti-PD-L1, anti-PD-1, or anti-PD-L2 antibodies or antigen-binding fragment in a form suitable for expression, under conditions suitable to produce such antibody or fragment, and recovering the antibody or fragment.

B. CTLA-4

Another immune checkpoint that can be targeted in the methods provided herein is the cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), also known as CD152. The complete cDNA sequence of human CTLA-4 has the Genbank accession number L15006. CTLA-4 is found on the surface of T cells and acts as an "off switch when bound to CD80 or CD86 on the surface of antigen-presenting cells. CTLA4 is a member of the immunoglobulin superfamily that is expressed on the surface of Helper T cells and transmits an inhibitory signal to T cells. CTLA4 is similar to the T-cell co-stimulatory protein, CD28, and both molecules bind to CD80 and CD86, also called B7-1 and B7-2 respectively, on antigen-presenting cells. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T cells and may be important to their function. T cell activation through the T cell receptor and CD28 leads to increased expression of CTLA-4, an inhibitory receptor for B7 molecules.

In some embodiments, the immune checkpoint inhibitor is an anti-CTLA-4 antibody (e.g., a human antibody, a humanized antibody, or a chimeric antibody), an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide.

Anti-human-CTLA-4 antibodies (or VH and/or VL domains derived therefrom) suitable for use in the present methods can be generated using methods well known in the art. Alternatively, art recognized anti-CTLA-4 antibodies can be used. For example, the anti-CTLA-4 antibodies disclosed in: U.S. Pat. No. 8,119,129, WO 01/14424, WO 98/42752; WO 00/37504 (CP675,206, also known as tremelimumab; formerly ticilimumab), U.S. Pat. No. 6,207,156; Hurwitz et al, 1998; can be used in the methods disclosed herein. The teachings of each of the aforementioned publications are hereby incorporated by reference. Antibodies that compete with any of these art-recognized antibodies for binding to CTLA-4 also can be used. For example, a humanized CTLA-4 antibody is described in International Patent Application No. WO2001014424, WO2000037504, and U.S. Pat. No. 8,017,114; all incorporated herein by reference.

An exemplary anti-CTLA-4 antibody is ipilimumab (also known as 10D1, MDX-010, MDX-101, and Yervoy®) or antigen binding fragments and variants thereof (see, e.g., WO01/14424). In other embodiments, the antibody comprises the heavy and light chain CDRs or VRs of ipilimumab.

Accordingly, in one embodiment, the antibody comprises the CDR1, CDR2, and CDR3 domains of the VH region of ipilimumab, and the CDR1, CDR2 and CDR3 domains of the VL region of ipilimumab. In another embodiment, the antibody competes for binding with and/or binds to the same epitope on CTLA-4 as the above-mentioned antibodies. In another embodiment, the antibody has at least about 90% variable region amino acid sequence identity with the above-mentioned antibodies (e.g., at least about 90%, 95%, or 99% variable region identity with ipilimumab).

Other molecules for modulating CTLA-4 include soluble CTLA-4 ligands and receptors such as described in U.S. Pat. Nos. 5,844,905, 5,885,796 and International Patent Application Nos. WO1995001994 and WO1998042752; all incorporated herein by reference, and immunoadhesins such as described in U.S. Pat. No. 8,329,867, incorporated herein by reference.

C. Killer Immunoglobulin-Like Receptor (KIR)

Another immune checkpoint inhibitor for use in the present invention is an anti-KIR antibody. Anti-human-KIR antibodies (or VH/VL domains derived therefrom) suitable for use in the present methods can be generated using methods well known in the art.

Alternatively, art recognized anti-KIR antibodies can be used. The anti-KIR antibody can be cross-reactive with multiple inhibitory KIR receptors and potentiates the cytotoxicity of NK cells bearing one or more of these receptors. For example, the anti-KIR antibody may bind to each of KIR2D2DL1, KIR2DL2, and KIR2DL3, and potentiate NK cell activity by reducing, neutralizing and/or reversing inhibition of NK cell cytotoxicity mediated by any or all of these KIRs. In some aspects, the anti-KIR antibody does not bind KIR2DS4 and/or KIR2DS3. For example, monoclonal antibodies 1-7F9 (also known as IPH2101), 14F1, 1-6F1 and 1-6F5, described in WO 2006/003179, the teachings of which are hereby incorporated by reference, can be used. Antibodies that compete with any of these art-recognized antibodies for binding to KIR also can be used. Additional art-recognized anti-KIR antibodies which can be used include, for example, those disclosed in WO 2005/003168, WO 2005/009465, WO 2006/072625, WO 2006/072626, WO 2007/042573, WO 2008/084106, WO 2010/065939, WO 2012/071411 and WO 2012/160448.

An exemplary anti-KIR antibody is lirilumab (also referred to as BMS-986015 or IPH2102). In other embodiments, the anti-KIR antibody comprises the heavy and light chain complementarity determining regions (CDRs) or variable regions (VRs) of lirilumab. Accordingly, in one embodiment, the antibody comprises the CDR1, CDR2, and CDR3 domains of the heavy chain variable (VH) region of lirilumab, and the CDR1, CDR2 and CDR3 domains of the light chain variable (VL) region of lirilumab. In another embodiment, the antibody has at least about 90% variable region amino acid sequence identity with lirilumab.

Examples of cancers contemplated for treatment include lung cancer, head and neck cancer, breast cancer, pancreatic cancer, prostate cancer, renal cancer, bone cancer, testicular cancer, cervical cancer, gastrointestinal cancer, lymphomas, pre-neoplastic lesions in the lung, colon cancer, melanoma, metastatic melanoma, basal-cell skin cancer, squamous-cell skin cancer, dermatofibrosarcoma protuberans, Merkel cell carcinoma, Kaposi's sarcoma, keratoacanthoma, spindle cell tumours, sebaceous carcinomas, microcystic adnexal carcinoma, Paget's disease of the breast, atypical fibroxanthoma, leiomyosarcoma, and angiosarcoma, Lentigo Maligna, Lentigo Maligna Melanoma, Superficial Spreading Melanoma, Nodular Melanoma, Acral Lentiginous Melanoma, Desmoplastic Melanoma, and bladder cancer.

In some embodiments, the subject has cancer that is resistant (has been demonstrated to be resistant) to one or more anti-cancer therapies. In some embodiments, resistance to anti-cancer therapy includes recurrence of cancer or refractory cancer. Recurrence may refer to the reappearance of cancer, in the original site or a new site, after treatment. In some embodiments, resistance to anti-cancer therapy includes progression of the cancer during treatment with the anti-cancer therapy. In some embodiments, the cancer is at

early stage or at late stage. The subject may have a cancer that expresses (has been shown to express e.g., in a diagnostic test) PD-L1 biomarker. In some embodiments, the patient's cancer expresses low PD-L1 biomarker. In some embodiments, the patient's cancer expresses high PD-L1 biomarker. The PD-L1 biomarker can be detected in the sample using a method selected from the group consisting of FACS, Western blot, ELISA, immunoprecipitation, immunohistochemistry, immunofluorescence, radioimmunoassay, dot blotting, immunodetection methods, HPLC, surface plasmon resonance, optical spectroscopy, mass spectrometry, HPLC, qPCR, RT-qPCR, multiplex qPCR or RT-qPCR, RNA-seq, microarray analysis, SAGE, MassARRAY technique, and FISH, and combinations thereof.

In some embodiments, the cancer has low levels of T cell infiltration. In some embodiments, the cancer has no detectable T cell infiltrate. In some embodiments, the cancer is a non-immunogenic cancer (e.g., non-immunogenic colorectal cancer and/or ovarian cancer).

For example, a therapeutically effective or sufficient amount of the immune checkpoint inhibitor, such as an antibody, is administered to a human will be in the range of about 0.01 to about 50 mg/kg of patient body weight whether by one or more administrations. In some embodiments, the antibody used is about 0.01 to about 45 mg/kg, about 0.01 to about 40 mg/kg, about 0.01 to about 35 mg/kg, about 0.01 to about 30 mg/kg, about 0.01 to about 25 mg/kg, about 0.01 to about 20 mg/kg, about 0.01 to about 15 mg/kg, about 0.01 to about 10 mg/kg, about 0.01 to about 5 mg/kg, or about 0.01 to about 1 mg/kg administered daily, for example. In some embodiments, the antibody is administered at 15 mg/kg. However, other dosage regimens may be useful. In one embodiment, an anti-PD-L1 antibody described herein is administered to a human at a dose of about 100 mg, about 200 mg, about 300 mg, about 400 mg, about 500 mg, about 600 mg, about 700 mg, about 800 mg, about 900 mg, about 1000 mg, about 1100 mg, about 1200 mg, about 1300 mg or about 1400 mg on day 1 of 21-day cycles. The dose may be administered as a single dose or as multiple doses (e.g., 2 or 3 doses), such as infusions. The progress of this therapy is easily monitored by conventional techniques.

Anti-Cancer and Other Therapies

In some embodiments, the immune checkpoint inhibitor may be administered in combination with at least one additional therapeutic. The additional therapy may be a cancer therapy such as radiation therapy, surgery, chemotherapy, gene therapy, DNA therapy, viral therapy, RNA therapy, immunotherapy, bone marrow transplantation, nanotherapy, monoclonal antibody therapy, or a combination of the foregoing. The additional therapy may be in the form of adjuvant or neoadjuvant therapy.

In an example, the therapy of the cancer (whether with or without administration of an immune checkpoint inhibitor) or any other disease (eg, viral infection or autoimmune disease) may radiation therapy, surgery, chemotherapy, gene therapy, DNA therapy, viral therapy, RNA therapy, immunotherapy, bone marrow transplantation, nanotherapy or monoclonal antibody therapy. The therapy may be a combination of the foregoing. An additional therapy may be administered

In some embodiments, the therapy (or the additional cancer therapy) is the administration of a small molecule enzymatic inhibitor or anti-metastatic agent. In some embodiments, the additional therapy is the administration of

side-effect limiting agents (e.g., agents intended to lessen the occurrence and/or severity of side effects of treatment, such as anti-nausea agents, etc.).

In some embodiments, the therapy (or the additional cancer therapy) is radiation therapy. In some embodiments, the therapy (or the additional cancer therapy) is surgery. In some embodiments, the therapy (or the additional cancer therapy) is a combination of radiation therapy and surgery. In some embodiments, the therapy (or the additional cancer therapy) is gamma irradiation. In some embodiments, the therapy (or the additional cancer therapy) is therapy targeting PBK/AKT/mTOR pathway, HSP90 inhibitor, tubulin inhibitor, apoptosis inhibitor, and/or chemopreventative agent. The therapy (or the additional cancer therapy) may be one or more of the chemotherapeutic agents known in the art.

Administration of any compound or therapy of the present embodiments to a patient will follow general protocols for the administration of such compounds, taking into account the toxicity, if any, of the agents. Therefore, in some embodiments there is a step of monitoring toxicity that is attributable to combination therapy.

The therapy can comprise or consist of administration to the subject of any of the following: —

1. Chemotherapy

A wide variety of chemotherapeutic agents may be used in accordance with the present embodiments. The term "chemotherapy" refers to the use of drugs to treat cancer. A "chemotherapeutic agent" is used to connote a compound or composition that is administered in the treatment of cancer. These agents or drugs are categorized by their mode of activity within a cell, for example, whether and at what stage they affect the cell cycle. Alternatively, an agent may be characterized based on its ability to directly cross-link DNA, to intercalate into DNA, or to induce chromosomal and mitotic aberrations by affecting nucleic acid synthesis.

Examples of chemotherapeutic agents include alkylating agents, such as thiotepa and cyclophosphamide; alkyl sulfonates, such as busulfan, improsulfan, and piposulfan; aziridines, such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines, including altretamine, triethylenemelamine, triethylenephosphoramide, triethylthiophosphoramide, and trimethylolmelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatins; callistatin; CC-1065 (eg, its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (eg, the synthetic analogues, KW-2189 and CB 1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards, such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, and uracil mustard; nitrosureas, such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics, such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gammall and calicheamicin omegall); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores, aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin, chromomycin, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (eg, morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-

doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, pofthromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, and zorubicin; anti-metabolites, such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues, such as denopterin, pteropterin, and trimetrexate; purine analogs, such as fludarabine, 6-mercaptopurine, thiamiprine, and thioguanine; pyrimidine analogs, such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, and floxuridine; androgens, such as calusterone, dromostanolone propionate, epitioestanol, mepitiostane, and testolactone; anti-adrenals, such as mitotane and trilostane; folic acid replenisher, such as frolic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids, such as maytansine and ansamitocins; mitoguanzone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK polysaccharide complex; razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (eg, T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; taxoids, e.g., paclitaxel and docetaxel gemcitabine; 6-thioguanine; mercaptopurine; platinum coordination complexes, such as cisplatin, oxaliplatin, and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (e.g., CPT-11); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids, such as retinoic acid; capecitabine; carboplatin, procarbazine, plicomycin, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatin, and pharmaceutically acceptable salts, acids, or derivatives of any of the above

2. Radiotherapy

Other factors that cause DNA damage and have been used extensively include what are commonly known as γ -rays, X-rays, and/or the directed delivery of radioisotopes to tumour cells. Other forms of DNA damaging factors are also contemplated, such as microwaves, proton beam irradiation (U.S. Pat. Nos. 5,760,395 and 4,870,287), and UV-irradiation. It is most likely that all of these factors affect a broad range of damage on DNA, on the precursors of DNA, on the replication and repair of DNA, and on the assembly and maintenance of chromosomes. Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 weeks), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

3. Immunotherapy

The skilled artisan will understand that immunotherapies may be used in combination or in conjunction with the methods described herein. In the context of cancer treatment, immunotherapeutics generally rely on the use of immune effector cells and molecules to target and destroy cancer cells. Rituximab (RITUXAN®) is an example of an immunotherapy. The immune effector may be, for example, an antibody specific for a marker on the surface of a tumour

cell. The antibody alone may serve as an effector of therapy or it may recruit other cells to actually effect cell killing. The antibody also may be conjugated to a drug or toxin (chemo therapeutic, radionuclide, ricin A chain, cholera toxin, pertussis toxin, etc.) and serve as a targeting agent. Alternatively, the effector may be a lymphocyte carrying a surface molecule that interacts, either directly or indirectly, with a tumour cell target. Various effector cells include cytotoxic T cells and NK cells.

In an example, the immunotherapy comprises adoptive cell therapy, such as CAR-T administration, eg, anti-CD19 or CD20 CAR-T administration.

In an example, the immunotherapy comprises or consists of administration of an IL-2 (eg, a truncated IL-2 or pegylated IL-2 or Fc-fused IL-2).

Antibody-drug conjugates have emerged as a breakthrough approach to the development of cancer therapeutics. Antibody-drug conjugates (ADCs) comprise monoclonal antibodies (MAbs) that are covalently linked to cell-killing drugs. This approach combines the high specificity of MAbs against their antigen targets with highly potent cytotoxic drugs, resulting in "armed" MAbs that deliver the payload (drug) to tumour cells with enriched levels of the antigen. Targeted delivery of the drug also minimizes its exposure in normal tissues, resulting in decreased toxicity and improved therapeutic index. The approval of two ADC drugs, ADCE-TRIS® (brentuximab vedotin) in 2011 and KADCYLA® (trastuzumab emtansine or T-DM1) in 2013 by FDA validated the approach. There are currently more than 30 ADC drug candidates in various stages of clinical trials for cancer treatment. As antibody engineering and linker-payload optimization are becoming more and more mature, the discovery and development of new ADCs are increasingly dependent on the identification and validation of new targets that are suitable to this approach and the generation of targeting MAbs. Two criteria for ADC targets are upregulated/high levels of expression in tumour cells and robust internalization.

In one aspect of immunotherapy, the tumour cell must bear some marker that is amenable to targeting, i.e., is not present on the majority of other cells. Many tumour markers exist and any of these may be suitable for targeting in the context of the present embodiments. Common tumour markers include CD20, carcinoembryonic antigen, tyrosinase (p97), gp68, TAG-72, HMFG, Sialyl Lewis Antigen, MucA, MucB, PLAP, laminin receptor, erb B, and pi 55. An alternative aspect of immunotherapy is to combine anticancer effects with immune stimulatory effects. Immune stimulating molecules also exist including: cytokines, such as IL-2, IL-4, IL-12, GM-CSF, gamma-IFN, chemokines, such as MIP-1, MCP-1, IL-8, and growth factors, such as FLT3 ligand.

4. Surgery

The cancer or other disease or condition may be treated by surgery in the invention.

Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative, and palliative surgery. Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed and may be used in conjunction with other therapies, such as the treatment of the present embodiments, chemotherapy, radiotherapy, hormonal therapy, gene therapy, immunotherapy, and/or alternative therapies. Tumour resection refers to physical removal of at least part of a tumour. In addition to tumour resection, treatment by surgery includes laser sur-

gery, cryosurgery, electro surgery, and microscopically-controlled surgery (Mohs' surgery).

Upon excision of part or all of cancerous cells, tissue, or tumour, a cavity may be formed in the body. Treatment may be accomplished by perfusion, direct injection, or local application of the area with an additional anti-cancer therapy. Such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. These treatments may be of varying dosages as well.

5. Bacterial Transplants

In an embodiment, the therapy comprises administering to the subject a bacterial transplant, eg, a fecal microbial transplant, comprising defined bacteria. For example, the transplant is any composition disclosed in WO2018064165, the disclosure of which (especially the compositions therein) are incorporated herein by reference in its entirety for possible application in the present invention. For example, the transplant is according to any of the following Paragraphs (tables and sequence numbers referring to the tables and sequences in WO2018064165, which are explicitly incorporated herein for possible use in the Claims): —

1. A composition comprising at least one isolated or purified population of bacteria belonging to one or more of the families Ruminococcaceae, Clostridiaceae, Lachnospiraceae, Micrococcaceae, and/or Veilonellaceae.

2. A composition comprising at least two isolated or purified populations of bacteria belonging to one or more of the families Ruminococcaceae, Clostridiaceae, Lachnospiraceae, Micrococcaceae, and/or Veilonellaceae.

3. The composition of Paragraph 1 or Paragraph 2, wherein each of the populations of bacteria is present in the composition at a concentration of at least 10^3 CFU.

4. The composition of Paragraph 1 or Paragraph 2, wherein the composition is a live bacterial product or a live biotherapeutic product.

5. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria are provided as bacterial spores.

6. The composition of Paragraph 1 or Paragraph 2, wherein the at least one population of bacteria or the at least two isolated or purified populations of bacteria belong to Clostridiales Family XII and/or Clostridiales Family XIII.

7. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria belong to the family Ruminococcaceae and/or of the family Clostridiaceae.

8. The composition of Paragraph 1 or Paragraph 2, wherein the population of bacteria belonging to the family Ruminococcaceae is further defined as a population of bacteria belonging to the genus *Ruminococcus*.

9. The composition of Paragraph 8, wherein the population of bacteria belonging to the genus *Ruminococcus* is further defined as a population of bacteria belonging to the species *Ruminococcus bromii*.

10. The composition of Paragraph 1 or Paragraph 2, wherein the population of bacteria belonging to the family Ruminococcaceae is further defined as a population of bacteria belonging to the genus *Faecalibacterium*.

11. The composition of Paragraph 10, wherein the population of bacteria belonging to the genus *Faecalibacterium* is further defined as a population of bacteria belonging to the species *Faecalibacterium prausnitzii*.

12. The composition of Paragraph 1 or Paragraph 2, wherein the population of bacteria belonging to the family Micro-

coccaceae is further defined as a population of bacteria belonging to the genus *Rothia*.

13. The composition of Paragraph 1 or Paragraph 2, wherein the composition further comprises a population of bacteria belonging to the species *Porphyromonas pasteri*, the species *Clostridium hungatei*, the species *Phascolarctobacterium faecium*, the genus *Peptoniphilus*, and/or the class Mollicutes.

14. The composition of Paragraph 1 or Paragraph 2, wherein the composition is essentially free of populations of bacteria belonging to the order Bacteroidales.

15. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria belongs to one or more of the species, subspecies or bacterial strains selected from the group consisting of the species in Table 1 with an enrichment index (ei) greater than 0.5.

16. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria are selected from the group consisting of the species in Table 1 with an "ei" equal to 1.

17. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria comprise a 16S ribosomal RNA (rRNA) nucleotide sequence that is at least 90% identical (eg, at least 91, 92, 93, 94, 95, 96, 97, 98 or 99% identical) to the 16S rRNA nucleotide sequence of bacteria identified by NCBI Taxonomy IDs selected from the group consisting of NCBI Taxonomy ID: 717959, 587, 758823, 649756, 44749, 671218, 1264, 1122135, 853, 484018, 46503, 54565, 290052, 216931, 575978, 433321, 1796646, 213810, 228924, 290054, 1509, 1462919, 29375, 337097, 1298596, 487174, 642492, 1735, 1297424, 742766, 46680, 132925, 411467, 1318465, 1852367, 1841857, 169679, 1175296, 259063, 172901, 39488, 57172, 28118, 166486, 28133, 1529, 694434, 1007096, 84030, 56774, 102148, 626947, 216933, 1348613, 1472417, 100176, 824, 1471761, 1297617, 288966, 1317125, 28197, 358743, 264639, 1265, 1335, 66219, 69473, 115117, 341220, 1732, 873513, 396504, 1796619, 45851, 2741, 105841, 86332, 1349822, 84037, 180311, 54291, 1217282, 762984, 1185412, 154046, 663278, 1543, 398512, 69825, 1841867, 1535, 1510, 84026, 1502, 1619234, 39497, 1544, 29343, 649762, 332095, 536633, 1033731, 574930, 742818, 177412, 1121308, 419208, 1673717, 55779, 28117, 626937, 180332, 1776382, 40519, 34062, 40518, 74426, 1216062, 293826, 850, 645466, 474960, 36835, 115544, 1515, 88431, 216932, 1417852, 39492, 1583, 420247, 118967, 169435, 37658, 138595, 31971, 100886, 1197717, 234908, 537007, 319644, 168384, 915173, 95159, 1816678, 626940, 501571, 1796620, 888727, 1147123, 376806, 1274356, 1267, 39495, 404403, 1348, 253314, 258515, 33033, 1118061, 357276, 214851, 320502, 217731, 246787, 29371, 649764, 901, 29374, 33043, 39778, 682400, 871665, 160404, 745368, 408, 1584, 333367, 47246, 1096246, 53342, 438033, 351091, 1796622, 1776384, 817, 48256, 720554, 500632, 36849, 301302, 879970, 655811, 264463, 1532, 285, 995, 242750, 29539, 1432052, 622312, 1796636, 1337051, 328814, 28446, 1492, 820, 39496, 52786, 1549, 1796618, 582, 46507, 109327, 1531, 1382, 33039, 311460, 230143, 216935, 539, 35519, 1681, 328813, 214853, 89014, 1121115, 1585974, 29466, 1363, 292800, 270498, 214856, 142877, 133926, 209880, 179628, 1121102, 105612, 1796615, 39777, 29353, 1579, 163665, 53443, 261299, 1302, 1150298, 938289, 358742, 471875, 938278, 1796613,

1118057, 1077144, 1737, 218205, 1121298, 684066, 433659, 52699, 204516, 706562, 253257, 328812, 1280, 147802, 58134, 1335613, 891, 585394, 1582, 235931, 308994, 1589, 1682, 1736, 28129, 178001, 551788, 2051, 856, 118562, 101070, 515619, 40215, 187979, 82979, 29363, 1776391, 1285191, 84112, 157688, 38304, 36850, 341694, 287, 75612, 818, 371674, 338188, 88164, 588581, 676965, 546271, 1236512, 178338, 862517, 157687, 158, 51048, 1583331, 529, 888745, 394340, 40545, 855, 553973, 938293, 93063, 708634, 179995, 1351, 476652, 1464038, 555088, 237576, 879566, 1852371, 742727, 1377, 35830, 997353, 218538, 83771, 1605, 28111, 131109, 46609, 690567, 46206, 155615, 51616, 40542, 203, 294, 1034346, 156456, 80866, 554406, 796942, 1002367, 29347, 796944, 61592, 487175, 1050201, 762948, 137732, 1211819, 1019, 272548, 1717, 384636, 216940, 2087, 45634, 466107, 1689, 47678, 575, 979627, 840, 1660, 1236517, 617123, 546, 28135, 82171, 483, 501496, 99656, 1379, 84032, 39483, 1107316, 584, 28124, 1033744, 657309, 536441, 76123, 1118060, 89152, 76122, 303, 1541, 507751, 515620, 38302, 53419, 726, 40324, 1796610, 988946, 1852370, 1017, 1168289, 76936, 94869, 1161098, 215580, 1125779, 327575, 549, 1450648 and 478.

18. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population of bacteria or the at least two isolated or purified populations of bacteria are a species, subspecies or bacterial strains comprising a 16S rRNA gene sequence at least 80% identical (eg, at least 85, 90, 95 or 98% identical) to any one of the sequences of SEQ ID NOs: 1-876 in WO2018064165.

19. The composition of Paragraph 1 or Paragraph 2, wherein the at least one isolated or purified population bacteria or the at least two isolated or purified populations of bacteria belong to the species, subspecies or bacterial strains selected from the group consisting of *Bacteroides coagulans*, *Clostridium aldenense*, *Clostridium aldrichii*, *Clostridium alkalicellulosi*, *Clostridium amygdalinum*, *Clostridium asparagiforme*, *Clostridium cellulosi*, *Clostridium citroniae*, *Clostridium clariflavum* DSM 19732, *Clostridium clostridioforme*, *Clostridium colinum*, *Clostridium fimetarium*, *Clostridium hiranonis*, *Clostridium hungatei*, *Clostridium hylemonae* DSM 15053, *Clostridium indolis*, *Clostridium lactatijementans*, *Clostridium leptum*, *Clostridium methylpentosum*, *Clostridium oroticum*, *Clostridium papyrosolvans* DSM 2782, *Clostridium populeti*, *Clostridium propionicum*, *Clostridium saccharolyticum*, *Clostridium scindens*, *Clostridium sporosphaeroides*, *Clostridium stercorarium*, *Clostridium straminisolvans*, *Clostridium sufflavum*, *Clostridium tennitidis*, *Clostridium thermosuccino* genes, *Clostridium viride*, *Clostridium xylanolyticum*, *Desulfotomaculum guttoideum*, *Eubacterium rectale* ATCC 33656, *Eubacterium dolichum*, *Eubacterium eligens* ATCC 27750, *Eubacterium hallii*, *Eubacterium infirmum*, *Eubacterium siraeum*, *Eubacterium tenue*, *Ruminococcus torques*, *Acetanaerobacterium elongatum*, *Acetatifactor muris*, *Acetivibrio cellulolyticus*, *Acetivibrio ethanolgignens*, *Acholeplasma brassicae* 0502, *Acholeplasma parvum*, *Acholeplasma vituli*, *Acinetobacter junii*, *Actinobacillus porcicus*, *Actinomyces bowdenii*, *Actinomyces dentalis*, *Actinomyces odontolyticus*, *Acutalibacter muris*, *Aerococcus viridans*, *Aeromicrobium fastidiosum*, *Alistipes finegoldii*, *Alistipes obesi*, *Alistipes onderdonkii*, *Alistipes putredinis*, *Alistipes shahii*, *Alistipes shahii* WAL 8301, *Alistipes timonensis* JC136, *Alkalibacter saccharofennentans*, *Alkaliphilus metalliredigens* QYMF, *Allisonella histaminiformans*, *Allobaculum stercoricans* DSM 13633, *Alloprevotella rava*, *Alloprevotella tannerae*, *Anaerobacterium chartisolvans*, *Anaerobspirillum thom-*

asii, *Anaerobium acetethylicum*, *Anaerococcus octavius* NCTC 9810, *Anaerococcus provenciensis*, *Anaerococcus vaginalis* ATCC 51170, *Anaerocolumna jejuensis*, *Anaerofilum agile*, *Anaerofustis stercorihominis*, *Anaeroglobus geminatus*, *Anaeromassilibacillus senegalensis*, *Anaeroplasmabactoclasticum*, *Anaerorhabdusfurcosa*, *Anaerosporeobacter mobilis*, *Anaerostipes butyraticus*, *Anaerostipes caccae*, *Anaerostipes hadrus*, *Anaerotruncus colihominis*, *Anaerovorax odorimutans*, *Anoxybacillus rupiensis*, *Aquabacterium limnoticum*, *Arcobacter butzleri*, *Arthrospira platensis*, *Asaccharobacter celatus*, *Atopobium parvulum*, *Bacteroides caccae*, *Bacteroides caecimuris*, *Bacteroides cellulosityticus*, *Bacteroides clarus* YIT 12056, *Bacteroides dorei*, *Bacteroides eggerthii*, *Bacteroides finegoldii*, *Bacteroides fragilis*, *Bacteroides gallinarum*, *Bacteroides massiliensis*, *Bacteroides oleiciplenus* YIT 12058, *Bacteroides plebeius* DSM 17135, *Bacteroides rodentium* JCM 16496, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*, *Bacteroides xylanisolvans* XB1A, *Bacteroides xylanolyticus*, *Barnesiella intestinihominis*, *Beduini massiliensis*, *Bifidobacterium bifidum*, *Bifidobacterium dentium*, *Bifidobacterium longum* subsp. *infantis*, *Blautia caecimuris*, *Blautia coccoides*, *Blautia faecis*, *Blautia gluceracea*, *Blautia hansenii* DSM 20583, *Blautia hydrogenotrophica*, *Blautia luti*, *Blautia luti* DSM 14534, *Blautia wexlerae* DSM 19850, *Budvicia aquatica*, *Butyricoccus pullicaecorum*, *Butyricomonas paravirosa*, *Butyriovibrio crossotus*, *Caldicoprobacter oshimai*, *Caloramator coolhaasii*, *Caloramatorproteoclasticus*, *Caloramator quimbayensis*, *Campylobacter gracilis*, *Campylobacter rectus*, *Campylobacter ureolyticus* DSM 20703, *Capnocytophaga gingivalis*, *Capnocytophaga leadbetteri*, *Capnocytophaga sputigena*, *Casaltella massiliensis*, *Catabacter hongkongensis*, *Catenibacterium mitsuokai*, *Christensenella minuta*, *Christensenella timonensis*, *Chryseobacterium taklimakanense*, *Citrobacterfreundii*, *Cloaciabacillus porcorum*, *Clostridioides difficile* ATCC 9689=DSM 1296, *Clostridium amylolyticum*, *Clostridium bowmanii*, *Clostridium butyricum*, *Clostridium cadaveris*, *Clostridium colicanis*, *Clostridium gasigenes*, *Clostridium lentocellum* DSM 5427, *Clostridium oceanicum*, *Clostridium oryzae*, *Clostridium paraputrificum*, *Clostridium pascui*, *Clostridium perfringens*, *Clostridium quini*, *Clostridium saccharobutylicum*, *Clostridium sporogenes*, *Clostridium ventriculi*, *Collinsella aerofaciens*, *Comamonas testosteroni*, *Coprobacter fastidiosus* NSB1, *Coprococcus eutactus*, *Corynebacterium diphtheriae*, *Corynebacterium durum*, *Corynebacterium mycetoides*, *Corynebacterium pyruviciproducens* ATCC BAA-1742, *Corynebacterium tuberculostearicum*, *Culturomica massiliensis*, *Cuneatibacter caecimuris*, *Deftuviitalea saccharophila*, *Delftia acidovorans*, *Desulfobacterium chlororespirans*, *Desulfobacterium metallireducens*, *Desulfosporosinus acididurans*, *Desulfotomaculum halophilum*, *Desulfotomaculum intricatum*, *Desulfotomaculum tongense*, *Desulfovibrio desulfuricans* subsp. *desulfuricans*, *Desulfovibrio idahonensis*, *Desulfovibrio litoralis*, *Desulfovibrio piger*, *Desulfovibrio simplex*, *Desulfovibrio zosteriae*, *Desulfuromonas acetoxidans*, *Dethiobacter alkaliphilus* AHT 1, *Dethiosulfatibacter aminovorans*, *Dialister invisus*, *Dialister propionificiens*, *Dielma fastidiosa*, *Dietzia alimentaria* 72, *Dorea longicatena*, *Dysgonomonas gadei* ATCC BAA-286, *Dysgonomonas mossii*, *Eggerthella lenta*, *Eikenella corrodens*, *Eisenbergiella tayi*, *Emergencia timonensis*, *Enorma massiliensis phi*, *Enterococcus faecalis*, *Enterorhabdus muris*, *Ethanoligenens harbinense* YUAN-3, *Eubacterium coprostanoligenes*, *Eubacterium limosum*, *Eubacterium oxidoreducens*, *Eubacterium sulci* ATCC

35585, *Eubacterium uniforme*, *Eubacterium ventriosum*, *Eubacterium xylanophilum*, *Extibacter muris*, *Ezakiella peruensis*, *Faecalibacterium prausnitzii*, *Faecalicoccus acidiformans*, *Faecalitalea cylindroides*, *Filifactor villosus*, *Flavonifractor plautii*, *Flintibacter butyricus*, *Frisingicoccus caecimuris*, *Fucophilus fucoidanolyticus*, *Fusicatenibacter saccharivorans*, *Fusobacterium mortiferum*, *Fusobacterium nucleatum* subsp. *vincentii*, *Fusobacterium simiae*, *Fusobacterium varium*, *Garciella nitratireducens*, *Gemella haemolysans*, *Gemmiger formicilis*, *Gordoniabacter urolithinifaciens*, *Gracilibacter thermotolerans* JW/YJL-S1, *Granulicatella elegans*, *Guggenheimella bovis*, *Haemophilus haemolyticus*, *Helicobacter typhlonius*, *Hespellia stercorisuis*, *Holdemanella biformis*, *Holdemania massiliensis* AP2, *Howardella ureilytica*, *Hungatella effluvii*, *Hungatella hathewayi*, *Hydrogenoanaerobacterium saccharovorans*, *Ihubacter massiliensis*, *Intestinibacter bartlettii*, *Intestinimonas butyriciproducens*, *Irregularibacter muris*, *Kiloniella laminariae* DSM 19542, *Kroppenstedtia guangzhouensis*, *Lachnoanaerobaculum orale*, *Lachnoanaerobaculum umeanense*, *Lachnoclostridium phytofermentans*, *Lactobacillus acidophilus*, *Lactobacillus algidus*, *Lactobacillus animalis*, *Lactobacillus casei*, *Lactobacillus delbrueckii*, *Lactobacillus formicilis*, *Lactobacillus iners*, *Lactobacillus pentosus*, *Lactobacillus rogosae*, *Lactococcus garvieae*, *Lactonifactor longoviformis*, *Leptotrichia buccalis*, *Leptotrichia hofstadii*, *Leptotrichia hongkongensis*, *Leptotrichia wadei*, *Leuconostoc inhae*, *Levyella massiliensis*, *Loriellopsis cavernicola*, *Lutispora thermophila*, *Marinilabilia salmonicolor* JCM 21150, *Marvinbryantia formatexigens*, *Mesoplasma photuris*, *Methanobrevibacter smithii* ATCC 35061, *Methanomassiliicoccus luminyensis* BIO, *Methylobacterium extorquens*, *Mitsuokella jalaludinii*, *Mobilitalea sibirica*, *Mobiluncus curtisii*, *Mogibacterium pumilum*, *Mogibacterium timidum*, *Moorella glycerini*, *Moorella humiferrea*, *Moraxella nonliquefaciens*, *Moraxella osloensis*, *Morganella morgani*, *Moryella indoligenes*, *Muribaculum intestinale*, *Murimonas intestini*, *Natranaerovirga pectinivora*, *Neglecta timonensis*, *Neisseria cinerea*, *Neisseria oralis*, *Nocardioides mesophilus*, *Novibacillus thermophilus*, *Ochrobactrum anthropi*, *Odoribacter splanchnicus*, *Olsenella profusa*, *Olsenella uli*, *Oribacterium asaccharolyticum* ACB7, *Oribacterium sinus*, *Oscillibacter ruminantium* GH1, *Oscillibacter valericigenes*, *Oxobacter pfennigii*, *Pantoea agglomerans*, *Papillibacter cinnamivorans*, *Parabacteroides faecis*, *Parabacteroides goldsteinii*, *Parabacteroides gordonii*, *Parabacteroides merdae*, *Parasporobacterium paucivorans*, *Parasutterella excrementihominis*, *Parasutterella secunda*, *Parvimonas micra*, *Peptococcus niger*, *Peptoniphilus duerdenii* ATCC BAA-1640, *Peptoniphilus grossensis* ph5, *Peptoniphilus koenoenienae*, *Peptoniphilus senegalensis* JC140, *Peptostreptococcus stomatis*, *Phascolarctobacterium succinatutens*, *Phoecea massiliensis*, *Pontibacter indicus*, *Porphyromonas bennonis*, *Porphyromonas endodontalis*, *Porphyromonas pasteri*, *Prevotella bergensis*, *Prevotella buccae* ATCC 33574, *Prevotella denticola*, *Prevotella enoeca*, *Prevotella fusca* JCM 17724, *Prevotella loescheii*, *Prevotella nigrescens*, *Prevotella oris*, *Prevotella pollens* ATCC 700821, *Prevotella stercorea* DSM 18206, *P. rev. ote. llamas silia timonensis*, *Propionispira arcuata*, *Proteus mirabilis*, *Providencia rettgeri*, *Pseudobacteroides cellulossolvans* ATCC 35603=DSM 2933, *Pseudobutyrvibrio ruminis*, *Pseudoflavonifractor capillosus* ATCC 29799, *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Pseudomonas mandelii*, *Pseudomonas nitroreducens*, *Pseudomonas putida*, *Raoultella ornithinolytica*, *Raoultella planticola*, *Raoultibacter*

massiliensis, *Robinsoniella peoriensis*, *Romboutsia timonensis*, *Roseburia faecis*, *Roseburia hominis* A2-183, *Roseburia intestinalis*, *Roseburia inulinivorans* DSM 16841, *Rothia dentocariosa* ATCC 17931, *Ruminiclostridium thermocellum*, *Ruminococcus albus*, *Ruminococcus bromii*, *Ruminococcus callidus*, *Ruminococcus champanellensis* 18P13=JCM 17042, *Ruminococcus faecis* JCM 15917, *Ruminococcus flavefaciens*, *Ruminococcus gauvreauii*, *Ruminococcus lactaris* ATCC 29176, *Summeiliabacillus pycnus*, *Saccharofermentans acetigenes*, *Scardovia wiggisiae*, *Schlegelella thermodepolymerans*, *Sedimentibacter hongkongensis*, *Selenomonas sputigena* ATCC 35185, *Slackia exigua* ATCC 700122, *Slackia piriformis* YIT 12062, *Solitalea canadensis*, *Solobacterium moorei*, *Sphingomonas aquatilis*, *Spiroplasma alleghenense*, *Spiroplasma chinense*, *Spiroplasma chrysopicola*, *Spiroplasma culicicola*, *Spiroplasma lamproyridicola*, *Sporobacter termitidis*, *Staphylococcus aureus*, *Stenotrophomonas maltophilia*, *Stomatobaculum longum*, *Streptococcus agalactiae* ATCC 13813, *Streptococcus cristatus*, *Streptococcus equinus*, *Streptococcus gordonii*, *Streptococcus lactarius*, *Streptococcus parauberis*, *Subdoligranulum variabile*, *Succinivibrio dextrinosolvans*, *Sutterella stercoricanis*, *Sutterella wadsworthensis*, *Syntrophococcus sucromutans*, *Syntrophomonas zehnderi* OL-4, *Terrisporobacter mayombeii*, *Thermoleophilum album*, *Treponema denticola*, *Treponema socranskii*, *Tyzzera nexilis* DSM 1787, *Vallitalea guaymasensis*, *Vallitalea pronyensis*, *Vampirovibrio chlorellavorus*, *Veillonella atypica*, *Veillonella denticariosi*, *Veillonella dispar*, *Veillonella parvula*, *Victivallis vadensis*, *Vulcanibacillus modesticaldus* and *Weissella confusa*.

In an example, the transplant comprises or consists of SER-109 or SER-262 (and optionally the condition is a *C difficile* infection); VE202 or SER-287 (and optionally the disease is ulcerative colitis); SER-301 (and optionally the disease is IBD); SER-401 (and optionally the condition is a cancer; eg, wherein the therapy further comprises administration of an anti-PD-1 axis antibody, eg, an anti-PD-1 antibody); VE800 or SER-155 (and optionally the therapy further comprises the administration of a transplant, eg, a haematopoietic stem cell or solid organ transplant); EDP1066 or EDP1815 (and optionally the disease is an inflammatory condition, eg, colitis, Crohn's disease, asthma, rheumatoid arthritis (RA), psoriasis, dermatitis (eg, atopic dermatitis) or IBD); or EDP1503 (and the disease is a cancer, eg, colorectal cancer, renal cell carcinoma, melanoma or a PD-1 relapsed cancer). In an example, the therapy comprises the administration of SGM-1019, SG-2-0776 or EB8018 (and optionally the disease or condition is NASH or IBD or an inflammatory condition, eg, colitis, Crohn's disease, asthma, rheumatoid arthritis (RA), psoriasis and dermatitis (eg, atopic dermatitis)). Those starting "VE" are developed by Vadanta Biosciences, SER are developed by Seres Therapeutics, EDP are developed by Evelo Biosciences, SG are developed by Second Genome and EB are developed by Enterome.

In an example, the disease or condition herein is an inflammatory condition, eg, colitis, Crohn's disease, asthma, rheumatoid arthritis (RA), psoriasis, dermatitis (eg, atopic dermatitis) or IBD.

6. Other Agents

It is contemplated that other agents may be used in combination with certain aspects of the present embodiments to improve the therapeutic efficacy of treatment. These additional agents include agents that affect the upregulation of cell surface receptors and GAP junctions, cytostatic and differentiation agents, inhibitors of cell adhe-

sion, agents that increase the sensitivity of the hyperproliferative cells to apoptotic inducers, or other biological agents. Increases in intercellular signalling by elevating the number of GAP junctions would increase the anti-hyperproliferative effects on the neighbouring hyperproliferative cell population. In other embodiments, cytostatic or differentiation agents can be used in combination with certain aspects of the present embodiments to improve the anti-hyperproliferative efficacy of the treatments. Inhibitors of cell adhesion are contemplated to improve the efficacy of the present embodiments. Examples of cell adhesion inhibitors are focal adhesion kinase (FAKs) inhibitors and Lovastatin. It is further contemplated that other agents that increase the sensitivity of a hyperproliferative cell to apoptosis, such as the antibody c225, could be used in combination with certain aspects of the present embodiments to improve the treatment efficacy.

Diseases and Conditions

Optionally, the disease or condition is selected from

- (a) A neurodegenerative disease or condition;
- (b) A brain disease or condition;
- (c) A CNS disease or condition;
- (d) Memory loss or impairment;
- (e) A heart or cardiovascular disease or condition, eg, heart attack, stroke or atrial fibrillation;
- (f) A liver disease or condition;
- (g) A kidney disease or condition, eg, chronic kidney disease (CKD);
- (h) A pancreas disease or condition;
- (i) A lung disease or condition, eg, cystic fibrosis or COPD;
- (j) A gastrointestinal disease or condition;
- (k) A throat or oral cavity disease or condition;
- (l) An ocular disease or condition;
- (m) A genital disease or condition, eg, a vaginal, labial, penile or scrotal disease or condition;
- (n) A sexually-transmissible disease or condition, eg, gonorrhoea, HIV infection, syphilis or *Chlamydia* infection;
- (o) An ear disease or condition;
- (p) A skin disease or condition;
- (q) A heart disease or condition;
- (r) A nasal disease or condition
- (s) A haematological disease or condition, eg, anemia, eg, anemia of chronic disease or cancer;
- (t) A viral infection;
- (u) A pathogenic bacterial infection;
- (v) A cancer;
- (w) An autoimmune disease or condition, eg, SLE;
- (x) An inflammatory disease or condition, eg, rheumatoid arthritis, psoriasis, eczema, asthma, ulcerative colitis, colitis, Crohn's disease or IBD;
- (y) Autism;
- (z) ADHD;
- (aa) Bipolar disorder;
- (bb) ALS [Amyotrophic Lateral Sclerosis];
- (cc) Osteoarthritis;
- (dd) A congenital or development defect or condition;
- (ee) Miscarriage;
- (ff) A blood clotting condition;
- (gg) Bronchitis;
- (hh) Dry or wet AMD;
- (ii) Neovascularisation (eg, of a tumour or in the eye);
- (jj) Common cold;
- (kk) Epilepsy;
- (ll) Fibrosis, eg, liver or lung fibrosis;
- (mm) A fungal disease or condition, eg, thrush;
- (nn) A metabolic disease or condition, eg, obesity, anorexia, diabetes, Type I or Type II diabetes.
- (oo) Ulcer(s), eg, gastric ulceration or skin ulceration;

- (pp) Dry skin;
- (qq) Sjogren's syndrome;
- (rr) Cytokine storm;
- (ss) Deafness, hearing loss or impairment;
- 5 (tt) Slow or fast metabolism (ie, slower or faster than average for the weight, sex and age of the subject);
- (uu) Conception disorder, eg, infertility or low fertility;
- (vv) Jaundice;
- 10 (ww) Skin rash;
- (xx) Kawasaki Disease;
- (yy) Lyme Disease;
- (zz) An allergy, eg, a nut, grass, pollen, dust mite, cat or dog fur or dander allergy;
- 15 (aaa) Malaria, typhoid fever, tuberculosis or cholera;
- (bbb) Depression;
- (ccc) Mental retardation;
- (ddd) Microcephaly;
- 20 (eee) Malnutrition;
- (fff) Conjunctivitis;
- (ggg) Pneumonia;
- (hhh) Pulmonary embolism;
- (iii) Pulmonary hypertension;
- 25 (jjj) A bone disorder;
- (kkk) Sepsis or septic shock;
- (lll) Sinusitis;
- (mmm) Stress (eg, occupational stress);
- (nnn) Thalassaemia, anaemia, von Willebrand Disease, or haemophilia;
- 30 (ooo) Shingles or cold sore;
- (ppp) Menstruation;
- (qqq) Low sperm count.

Neurodegenerative or CNS Diseases or Conditions for Treatment or Prevention by the Method

In an example, the neurodegenerative or CNS disease or condition is selected from the group consisting of Alzheimer disease, geriopsychosis, Down syndrome, Parkinson's disease, Creutzfeldt-jakob disease, diabetic neuropathy, Parkinson syndrome, Huntington's disease, Machado-Joseph disease, amyotrophic lateral sclerosis, diabetic neuropathy, and Creutzfeldt Creutzfeldt-Jakob disease. For example, the disease is Alzheimer disease. For example, the disease is Parkinson syndrome.

45 In an example, wherein the method of the invention is practised on a human or animal subject for treating a CNS or neurodegenerative disease or condition, the method causes downregulation of Treg cells in the subject, thereby promoting entry of systemic monocyte-derived macrophages and/or Treg cells across the choroid plexus into the brain of the subject, whereby the disease or condition (eg, Alzheimer's disease) is treated, prevented or progression thereof is reduced. In an embodiment the method causes an increase of IFN-gamma in the CNS system (eg, in the brain and/or CSF) of the subject. In an example, the method restores nerve fibre and/or reduces the progression of nerve fibre damage. In an example, the method restores nerve myelin and/or reduces the progression of nerve myelin damage. In an example, the method of the invention treats or prevents a disease or condition disclosed in WO2015136541 and/or the method can be used with any method disclosed in WO2015136541 (the disclosure of this document is incorporated by reference herein in its entirety, eg, for providing disclosure of such methods, diseases, conditions and potential therapeutic agents that can be administered to the subject for effecting treatment and/or prevention of CNS and neurodegenerative diseases and conditions, eg, agents such as

immune checkpoint inhibitors, eg, anti-PD-1, anti-PD-L1, anti-TIM3 or other antibodies disclosed therein).

Cancers for Treatment or Prevention by the Method

Cancers that may be treated include tumours that are not vascularized, or not substantially vascularized, as well as vascularized tumours. The cancers may comprise non-solid tumours (such as haematological tumours, for example, leukaemias and lymphomas) or may comprise solid tumours. Types of cancers to be treated with the invention include, but are not limited to, carcinoma, blastoma, and sarcoma, and certain leukaemia or lymphoid malignancies, benign and malignant tumours, and malignancies e.g., sarcomas, carcinomas, and melanomas. Adult tumours/cancers and pediatric tumours/cancers are also included.

Hematologic cancers are cancers of the blood or bone marrow. Examples of haematological (or hematogenous) cancers include leukaemias, including acute leukaemias (such as acute lymphocytic leukaemia, acute myelocytic leukaemia, acute myelogenous leukaemia and myeloblasts, promyelocytic, myelomonocytic, monocytic and erythro-leukemia), chronic leukaemias (such as chronic myelocytic (granulocytic) leukaemia, chronic myelogenous leukaemia, and chronic lymphocytic leukaemia), polycythemia vera, lymphoma, Hodgkin's disease, non-Hodgkin's lymphoma (indolent and high grade forms), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, myelodysplastic syndrome, hairy cell leukaemia and myelodysplasia.

Solid tumours are abnormal masses of tissue that usually do not contain cysts or liquid areas. Solid tumours can be benign or malignant. Different types of solid tumours are named for the type of cells that form them (such as sarcomas, carcinomas, and lymphomas). Examples of solid tumours, such as sarcomas and carcinomas, include fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteosarcoma, and other sarcomas, synovioma, mesothelioma, Ewing's tumour, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, lymphoid malignancy, pancreatic cancer, breast cancer, lung cancers, ovarian cancer, prostate cancer, hepatocellular carcinoma, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, medullary thyroid carcinoma, papillary thyroid carcinoma, pheochromocytomas sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, Wilms' tumour, cervical cancer, testicular tumour, seminoma, bladder carcinoma, melanoma, and CNS tumours (such as a glioma (such as brainstem glioma and mixed gliomas), glioblastoma (also known as glioblastoma multiforme) astrocytoma, CNS lymphoma, germinoma, medulloblastoma, Schwannoma craniopharyngioma, ependymoma, pineaioma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, neuroblastoma, retinoblastoma and brain metastases).

In an example, the cancer is a haematological cancer. In an example, the cancer is NSCLC. In an example, the cancer is renal cell carcinoma. In an example, the cancer is urothelial carcinoma. In an example, the cancer is melanoma.

Autoimmune Diseases for Treatment or Prevention by the Method

Acute Disseminated Encephalomyelitis (ADEM)
Acute necrotizing hemorrhagic leukoencephalitis
Addison's disease
Agammaglobulinemia
Alopecia areata
Amyloidosis

Ankylosing spondylitis
Anti-GBM/Anti-TBM nephritis
Antiphospholipid syndrome (APS)
Autoimmune angioedema
Autoimmune aplastic anemia
Autoimmune dysautonomia
Autoimmune hepatitis
Autoimmune hyperlipidemia
Autoimmune immunodeficiency
Autoimmune inner ear disease (AIED)
Autoimmune myocarditis
Autoimmune oophoritis
Autoimmune pancreatitis
Autoimmune retinopathy
Autoimmune thrombocytopenic purpura (ATP)
Autoimmune thyroid disease
Autoimmune urticaria
Axonal & neuronal neuropathies
Balo disease
Behcet's disease
Bullous pemphigoid
Cardiomyopathy
Castleman disease
Celiac disease
Chagas disease
Chronic fatigue syndrome
Chronic inflammatory demyelinating polyneuropathy (CIDP)
Chronic recurrent multifocal osteomyelitis (CRMO)
Churg-Strauss syndrome
Cicatricial pemphigoid/benign mucosal pemphigoid
Crohn's disease
Cogans syndrome
Cold agglutinin disease
Congenital heart block
Coxsackie myocarditis
CREST disease
Essential mixed cryoglobulinemia
Demyelinating neuropathies
Dermatitis herpetiformis
Dermatomyositis
Devic's disease (neuromyelitis optica)
Discoid lupus
Dressler's syndrome
Endometriosis
Eosinophilic esophagitis
Eosinophilic fasciitis
Erythema nodosum
Experimental allergic encephalomyelitis
Evans syndrome
Fibromyalgia
Fibrosing alveolitis
Giant cell arteritis (temporal arteritis)
Giant cell myocarditis
Glomerulonephritis
Goodpasture's syndrome
Granulomatosis with Polyangiitis (GPA) (formerly called Wegener's Granulomatosis)
Graves' disease
Guillain-Barre syndrome
Hashimoto's encephalitis
Hashimoto's thyroiditis
Hemolytic anemia
Henoch-Schonlein purpura
Herpes gestationis
Hypogammaglobulinemia
Idiopathic thrombocytopenic purpura (ITP)

IgA nephropathy
 IgG4-related sclerosing disease
 Immunoregulatory lipoproteins
 Inclusion body myositis
 Interstitial cystitis
 Juvenile arthritis
 Juvenile diabetes (Type 1 diabetes)
 Juvenile myositis
 Kawasaki syndrome
 Lambert-Eaton syndrome
 Leukocytoclastic vasculitis
 Lichen planus
 Lichen sclerosus
 Ligneous conjunctivitis
 Linear IgA disease (LAD)
 Lupus (SLE)
 Lyme disease, chronic
 Meniere's disease
 Microscopic polyangiitis
 Mixed connective tissue disease (MCTD)
 Mooren's ulcer
 Mucha-Habermann disease
 Multiple sclerosis
 Myasthenia gravis
 Myositis
 Narcolepsy
 Neuromyelitis optica (Devic's)
 Neutropenia
 Ocular cicatricial pemphigoid
 Optic neuritis
 Palindromic rheumatism
 PANDAS (Pediatric Autoimmune Neuropsychiatric Disorders Associated with *Streptococcus*)
 Paraneoplastic cerebellar degeneration
 Paroxysmal nocturnal hemoglobinuria (PNH)
 Parry Romberg syndrome
 Parsonnage-Turner syndrome
 Pars planitis (peripheral uveitis)
 Pemphigus
 Peripheral neuropathy
 Perivenous encephalomyelitis
 Pernicious anemia
 POEMS syndrome
 Polyarteritis *nodosa*
 Type I, II, & III autoimmune polyglandular syndromes
 Polymyalgia rheumatica
 Polymyositis
 Postmyocardial infarction syndrome
 Postpericardiotomy syndrome
 Progesterone dermatitis
 Primary biliary cirrhosis
 Primary sclerosing cholangitis
 Psoriasis
 Psoriatic arthritis
 Idiopathic pulmonary fibrosis
 Pyoderma gangrenosum
 Pure red cell aplasia
 Raynauds phenomenon
 Reactive Arthritis
 Reflex sympathetic dystrophy
 Reiter's syndrome
 Relapsing polychondritis
 Restless legs syndrome
 Retroperitoneal fibrosis
 Rheumatic fever
 Rheumatoid arthritis
 Sarcoidosis

Schmidt syndrome
 Scleritis
 Scleroderma
 Sjogren's syndrome
 5 Sperm & testicular autoimmunity
 Stiff person syndrome
 Subacute bacterial endocarditis (SBE)
 Susac's syndrome
 Sympathetic ophthalmia
 10 Takayasu's arteritis
 Temporal arteritis/Giant cell arteritis
 Thrombocytopenic purpura (TTP)
 Tolosa-Hunt syndrome
 Transverse myelitis
 15 Type 1 diabetes
 Ulcerative colitis
 Undifferentiated connective tissue disease (UCTD)
 Uveitis
 Vasculitis
 20 Vesiculobullous dermatosis
 Vitiligo
 Wegener's granulomatosis (now termed Granulomatosis with Polyangiitis (GPA)).
 Inflammatory Diseases for Treatment or Prevention by the
 25 Method
 Alzheimer's
 ankylosing spondylitis
 arthritis (osteoarthritis, rheumatoid arthritis (RA), psoriatic arthritis)
 30 asthma
 atherosclerosis
 Crohn's disease
 colitis
 dermatitis
 35 diverticulitis
 fibromyalgia
 hepatitis
 irritable bowel syndrome (IBS)
 systemic lupus erythematosus (SLE)
 40 nephritis
 Parkinson's disease
 ulcerative colitis.
 Concepts: —
 The invention provides the following Concepts.
 1. A programmable nuclease for use in a method of treating an acute microbial infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the subject is treated.
 50
 2. A programmable nuclease for use in a method of durably treating a microbial (eg, bacterial) infection of a subject, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the subject, whereby microbes of the first species or strain are durably killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nucle-
 65

- ase is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and microbial infection of the subject is durably treated.
3. The nuclease of Concept 2, wherein the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1.
 4. The nuclease of any preceding Concept, wherein the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
 5. The nuclease of any preceding Concept, wherein the method comprises maintaining reduction of the infection by at least 100-fold for at least 60 minutes (eg, at least 120 minutes) after exposing the subject to the programmed nuclease.
 6. The nuclease of any preceding Concept, wherein the method comprises reducing the infection such that the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.
 7. The nuclease of any preceding Concept, wherein the method comprises administering to the subject a RNA or a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject.
 8. The nuclease of Concept 7, wherein the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject.
 9. The nuclease of Concept 7, wherein the subject comprises the nuclease prior to administration of the RNA or nucleic acid to the subject.
 10. The nuclease of any one of Concepts 7 to 9, wherein a plurality of viruses (eg, phage) are administered to the subject, wherein each virus comprises a copy of the nucleic acid, wherein the viruses infect the microbes comprised by the subject to deliver thereto the nucleic acid.
 11. The nuclease of Concept 10, wherein the ratio of administered viruses: microbes comprised by the subject is from 10 to 150.
 12. The nuclease according to any preceding Concept, wherein the subject is a human or animal, optionally wherein the subject is a human over 65 years of age or is a pediatric patient.
 13. The nuclease according to Concept 12, wherein the infection is an infection of the lungs, abdomen or urinary tract; or wherein the subject has undergone surgery, is on an immunosuppressant medication and/or is suffering from a chronic disease.
 14. The nuclease according to any preceding Concept, wherein the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
 15. The nuclease according to any preceding Concept, wherein the method comprises reducing the infection at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject to the programmed nuclease.
 16. The nuclease according to any one of Concepts 12 to 15, wherein the method treats or prevents septicemia and/or sepsis (eg, septic shock) in the subject.

17. The nuclease of Concept 16, wherein at the start of the treatment, the subject (eg, a human) has a temperature of $<36^{\circ}\text{C}$. or $>38^{\circ}\text{C}$.; a heart rate of $>90/\text{min}$, a respiratory rate of >20 breaths/min or $\text{PaCO}_2 < 4.3\text{ kPa}$; and white blood cell count of $<4000/\text{mm}^3$ or $>12,000/\text{mm}^3$.
18. The nuclease of Concept 16 or 17, wherein at the start of the treatment, the subject (eg, a human) has presence of two or more of the following: abnormal body temperature, abnormal heart rate, abnormal respiratory rate, abnormal blood gas and abnormal white blood cell count.
19. The nuclease of any preceding Concept, wherein the subject is a human or animal and the microbes are bacteria (eg, *E coli* or *C difficile*), wherein blood infection of the subject by the bacteria is reduced at least 100- or 1000-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
20. The nuclease of any one of Concepts 12 to 19, wherein the blood of the subject is infected with from 10^7 to 10^{12} CFU/ml of the bacteria immediately before the treatment.
21. The nuclease according to any one of Concepts 1 to 11, wherein the subject is a plant.
22. The nuclease according to any preceding Concept, wherein the microbes are bacteria.
23. The nuclease according to Concept 22, wherein the bacteria are gram positive bacteria.
24. The nuclease according to Concept 22 or 23, wherein the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).
25. The nuclease according to any preceding Concept, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.
26. A plurality of viruses (eg, phage or phagemids for producing phage) for use with the nuclease of any preceding Concept in the method of treatment, wherein each virus comprises a copy of a nucleic acid as defined in any one of Concepts 7 to 9, wherein the viruses are capable of infecting microbes comprised by the subject to deliver thereto the nucleic acid.
27. A composition comprising a plurality of nucleic acids for programming the nuclease of any one of Concepts 1 to 25 in the method of treatment, wherein each nucleic acid is a nucleic acid as defined in any one of Concepts 7 to 9.
28. A CRISPR/Cas system comprising a nuclease according to any preceding Concept for use in the method of treatment, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9) and the system comprises one or more guide RNAs or DNA encoding one or more guide RNAs, wherein each guide RNA is capable of programming the Cas nuclease to cut a target site comprised by the genomes of the microbes.
29. A guide RNA or a DNA encoding a guide RNA for use in the system of Concept 28 for use in the method of treating an acute microbial infection in the subject, eg, septicemia or sepsis.
30. A nucleic acid vector comprising the guide RNA or DNA recited in Concept 27 or 29.

31. The vector of Concept 30 wherein the vector is a phage, phagemid, virophage, virus, plasmid (eg, conjugative plasmid) or transposon.
32. An anti-sepsis or anti-septicaemia composition for administration to a human or animal for treating sepsis or septicaemia, the composition comprising a plurality of vectors, wherein each vector is according to Concept 30 or 31.
33. A method of treating an acute microbial infection of a subject, wherein the method is as defined by any preceding Concept.
34. Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of any one of Concepts 1 to 26 and 28 to 30, in the manufacture of a composition for carrying out a method of treatment as defined by any preceding Concept, wherein the subject is an organism other than a human or animal.
35. Use of a nuclease, plurality of viruses, system, guide RNA, DNA or vector of any one of Concepts 1 to 26 and 28 to 30, in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.
36. Use of a programmable nuclease in the manufacture of a composition for carrying out an ex vivo method of treatment of a microbial infection of a substrate, wherein the microbial infection is caused by microbes of a first species or strain and the nuclease is programmable to cut a target site comprised by the genomes of microbes that have infected the substrate, whereby microbes of the first species or strain are killed, or growth or proliferation of the microbes is reduced, the treatment method comprising exposing the subject to the nuclease wherein the nuclease is programmed to cut the target site, whereby genomes of the microbes comprised by the subject are cut and acute microbial infection of the substrate is treated.
37. The use of Concept 34, 35 or 36, wherein the nuclease (eg, programmed nuclease) and/or a nucleic acid that programs the nuclease to recognise and cut the target site is administered to the subject or substrate at a first time (T1) and at a second time (T2) wherein T2 is at least 1 hour after T1.
38. The use of any one of Concepts 34 to 37, wherein the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
39. The use of any one of Concepts 34 to 38, wherein the reduction of the infection is maintained by at least 100-fold for at least 60 minutes (eg, at least 120 minutes) after exposing the subject to the programmed nuclease.
40. The use of any one of Concepts 34 to 39, wherein the reduction in infection persists for 30 minutes immediately after the first 30 minutes of the treatment.
41. The use of any one of Concepts 34 to 40, wherein the method comprises administering to the subject or substrate a RNA or a nucleic acid that encodes an RNA for

- expression of the RNA in or on the subject or substrate, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in microbes comprised by the subject or substrate.
42. The use of Concept 41, wherein the nuclease is administered simultaneously or sequentially with the RNA or nucleic acid to the subject or substrate.
43. The use of Concept 41, wherein the subject or substrate comprises the nuclease prior to administration of the RNA or nucleic acid.
44. The use of any one of Concepts 41 to 43, wherein a plurality of viruses (eg, phage) are administered to the subject or substrate, wherein each virus comprises a copy of the nucleic acid, wherein the viruses infect the microbes comprised by the subject or substrate to deliver thereto the nucleic acid.
45. The use of Concept 44, wherein the ratio of administered viruses: microbes is from 10 to 150.
46. The use of any one of Concepts 34 to 45, wherein the infection is reduced by at least 90% for 1 hour or more, optionally by the first 30 minutes (eg, by the first 15 minutes) of the treatment.
47. The use of any one of Concepts 34 to 46, wherein the infection is reduced at least 100-fold by the first 30 minutes (eg, by the first 15 minutes) of the treatment; and wherein reduction of the infection by at least 100-fold is maintained for at least 60 minutes (eg, at least 120, 145 or 180 minutes) after exposing the subject or substrate to the programmed nuclease.
48. The use of any one of Concepts 34 to 47, wherein the subject is a plant; or wherein the substrate is a metallic, plastic, concrete, stone, wood, glass or ceramic substrate.
49. The use of any one of Concepts 34 to 48, wherein the microbes are bacteria.
50. The use according to Concept 49, wherein the bacteria are gram positive bacteria.
51. The use according to Concept 49 or 50, wherein the bacteria are *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Legionella*, *Haemophilus*, *Ghonnorhea*, *Acinetobacter*, *Escherichia*, *Klebsiella*, *Pseudomonas* or *Stenotrophomonas* bacteria (eg, *E coli* (eg, EHEC *E coli*), *C difficile*, *V cholera*, *Staphylococcus* (eg, *S aureus* or MRSA), *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* bacteria).
52. The use of any one of Concepts 34 to 51, wherein the nuclease is a Cas nuclease (eg, a Cas 3 or 9), a meganuclease, a TALEN (Transcription activator-like effector nuclease) or zinc finger nuclease.

EMBODIMENTS

1. A method for treating a pathogenic bacterial infection in a human or animal subject caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a programmable nuclease that is programmed to cut the target site, wherein the subject is suffering from a further disease or condition other than the pathogenic bacterial infection and the method comprises administering a therapy to the subject for treating or preventing the further disease or condition, wherein the nuclease treats

- the infection and the therapy is efficacious in the presence of the programmed nuclease to treat or prevent the disease or condition.
2. The method of Embodiment 1, wherein the subject is a cancer patient and the therapy comprises administration of a haematopoietic stem cell transplant, chemotherapeutic agent, immune checkpoint inhibitor, immune checkpoint agonist or an immune cell enhancer; adoptive cell therapy; radiation or surgery.
 3. The method of Embodiment 2, wherein the therapy is an immune checkpoint inhibitor antibody, or an antibody selected from ipilimumab (or YERVOY®), tremelimumab, nivolumab (or OPDIVO®), pembrolizumab (or KEYTRUDA®), pidilizumab, BMS-936559, durvalumab (or IMFINZI®) and atezolizumab (or TECENTRIQ®).
 4. The method of Embodiment 1, wherein the therapy is a tissue, organ or cell transplant.
 5. The method of Embodiment 1, wherein the treatment of the bacterial infection is carried out simultaneously with the administration of the therapy to the subject.
 6. The method of Embodiment 1, wherein the treatment of the bacterial infection is carried out immediately before or after administering the therapy to the subject.
 7. The method of Embodiment 1, wherein the method comprises administering to the subject a or a nucleic acid that encodes an RNA for expression of the RNA in the subject, wherein the RNA complexes with the nuclease to program the nuclease to cut the target site in first bacteria comprised by the subject, thereby killing the first bacteria.
 8. The method of Embodiment 1, comprising administering a nucleic acid vector to the subject, wherein the vector encodes the programmable nuclease.
 9. The method of Embodiment 1, wherein the programmable nuclease is an endogenous nuclease of the first cells.
 10. The method of Embodiment 1, wherein the efficacy of the therapy in the presence of the programmed nuclease is greater than the efficacy of the therapy in the presence of a broad-spectrum antibiotic.
 11. The method of Embodiment 1, wherein the efficacy of the therapy in the presence of the programmed nuclease is greater than the efficacy of the therapy in the presence of an antibiotic selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbapristin; teicoplanin; cephalosporin; carbapenem; fluoroquinolone; aminoglycoside; colistin; erythromycin; clindamycin; beta-lactam; macrolide; amoxicillin; azithromycin; penicillin; ceftriaxone; azithromycin; ciprofloxacin; isoniazid (INH); rifampicin (RMP); amikacin; kanamycin; capreomycin; trimethoprim; itrofurantoin; cefalexin; amoxicillin; metronidazole (MTZ); cefixime; tetracycline; and meropenem.
 12. The method of Embodiment 1, wherein the first bacteria is selected from (i) *Staphylococcus aureus* that is resistant to an antibiotic selected from methicillin, vancomycin, linezolid, daptomycin, quinupristin, dalbapristin and teicoplanin; (ii) *Pseudomonas aeruginosa* that is resistant to an antibiotic selected from cephalosporins, carbapenems, fluoroquinolones, aminoglycosides and colistin; (iii) *Klebsiella* species that is resistant to carbapenem; (iv) *Streptococcus* species that is resistant to an antibiotic selected from erythromycin, clindamycin, beta-lactam, macrolide, amoxicillin, azithromycin and penicillin; (v) *Salmonella* species that is resistant to an antibiotic selected from ceftriaxone, azithromycin and ciprofloxacin; (vi) *Shigella* species that is resistant to ciprofloxacin or azithromycin; (vii) *Mycobacterium tuberculosis* that is resistant to an antibiotic selected from Resistance to

- isoniazid (INH), rifampicin (RMP), fluoroquinolone, amikacin, kanamycin, capreomycin and azithromycin; (viii) *Enterococcus* species that is resistant to vancomycin; (ix) Enterobacteriaceae species that is resistant to an antibiotic selected from cephalosporin and carbapenem; (x) *E coli* that is resistant to an antibiotic selected from trimethoprim, itrofurantoin, cefalexin and amoxicillin; (xi) *Clostridium* species that is resistant to metronidazole (MTZ), fluoroquinolone or carbapenem; (xii) *Neisseria gonorrhoea* that is resistant to an antibiotic selected from cefixime, ceftriaxone, azithromycin and tetracycline; (xiii) *Acinetobacter baumannii* that is resistant to an antibiotic selected from beta-lactam, meropenem and carbapenem; and (xiv) *Campylobacter* species that is resistant to ciprofloxacin or azithromycin.
13. The method of Embodiment 1, wherein the treatment of the infection treats or prevents in the subject a condition selected from vaginosis, meningitis, pneumonia, urinary tract infection, cystitis, nephritis, gastroenteritis, a skin infection, impetigo, erysipelas, cellulitis, septicaemia or sepsis in the subject.
 14. The method of Embodiment 1, wherein the further disease or condition is a cancer; autoimmune disease or condition; or GI tract disease or condition.
 15. The method of Embodiment 1, wherein the subject comprises bacteria (second bacteria) of one or more strains or species that are different to the first strain or species, wherein the genomes of the second bacteria do not comprise the target site, wherein the genomes of the second bacteria are not cut by the programmed nuclease in the subject, whereby second bacteria survive in the presence of the programmed nuclease in the patient; and wherein the therapy is efficacious in the presence of the second bacteria.
 16. The method of Embodiment 15, wherein reduction in the second bacteria in patients is associated with reduced efficacy of the therapy.
 17. The method of Embodiment 15, wherein the second bacteria are selected from the group consisting of *Akkermansia*, *Alistipes*, *Bacteroides*, *Barnesiella*, *Bifidobacterium*, *Clostridium*, *Collinsella*, *Enterococcus*, *Fusobacterium*, *Lactobacillus*, *Propionibacterium*, *Ruminococcus*, *Segmented filamentous bacteria* (SFB); *Veillonella*, *Prevotella*, *Escherichia* and *Streptococcus bacteria*.
 18. The method of Embodiment 1, wherein the first bacteria are selected from the group consisting of *E coli*, *C difficile*, *V cholera*, *Staphylococcus*, *Streptococcus pyogenes*, *Acinetobacter baumannii*, *Legionella*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* bacteria.
 19. The method of Embodiment 1, wherein the nuclease is a Cas nuclease, a meganuclease, a Transcription activator-like effector nuclease (TALEN) or zinc finger nuclease.
 20. A method for treating a pathogenic bacterial infection in a cancer patient caused by bacteria (first bacteria) of a first species or strain, the method comprising selectively killing first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by guide RNA to cut the target site, wherein the method comprises administering an immunotherapy to the subject for treating cancer in the patient, wherein the nuclease treats the infection and the immunotherapy is efficacious in the presence of the programmed nuclease to treat the cancer.
- It will be understood that particular embodiments described herein are shown by way of illustration and not as limitations of the invention. The principal features of this

invention can be employed in various embodiments without departing from the scope of the invention. Those skilled in the art will recognize, or be able to ascertain using no more than routine study, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of this invention and are covered by the claims. All publications and patent applications mentioned in the specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications and all US equivalent patent applications and patents are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference. The use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.” The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” Throughout this application, the term “about” is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

As used in this specification and claim(s), the words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, unrecited elements or method steps

The term “or combinations thereof” or similar as used herein refers to all permutations and combinations of the listed items preceding the term. For example, “A, B, C, or combinations thereof” is intended to include at least one of: A, B, C, AB, AC, BC, or ABC, and if order is important in a particular context, also BA, CA, CB, CBA, BCA, ACB, BAC, or CAB. Continuing with this example, expressly included are combinations that contain repeats of one or more item or term, such as BB, AAA, MB, BBC, AAABCCCC, CBBAAA, CABABB, and so forth. The skilled artisan will understand that typically there is no limit on the number of items or terms in any combination, unless otherwise apparent from the context.

Any part of this disclosure may be read in combination with any other part of the disclosure, unless otherwise apparent from the context.

All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

The present invention is described in more detail in the following non limiting Examples.

EXAMPLES

Precision Fast Bacteria Killing with Programmable Nucleases

The examples provide a method for fast and precision killing of *Escherichia coli* and *Clostridium difficile* strains. As a model programmable nuclease system, we used a CRISPR guided vector (CGV™) system to specifically target enterohemorrhagic *E. coli* (EHEC) and probiotic *E. coli* Nissle.

Example 1. Precision Killing of Target Strain Enterohemorrhagic *E. coli* (EHEC)

1.1. Design, Construction and Delivery of CRISPR Guided Vector (CGV) System Targeting *E. coli* (EHEC) ATCC43888.

The invention provides a CGV system to specifically target enterohemorrhagic *E. coli* (EHEC) ATCC43888 (a human fecal isolate obtained from the American Type Culture Collection). The CGV system comprises two vectors: (a) a vector containing a tracrRNA and the Cas9 protein from *Streptococcus pyogenes* (SpCas); (b) a vector containing a guide RNA (gRNA) that comprises a nucleotide sequence capable of hybridizing to a target sequence in the host cells to guide SpCas9 to the target sequence. To enable specific killing of *E. coli* (EHEC) ATCC43888, a particular sequence from the genome of this strain was chosen to target. Specifically, the sequence contains 20 nucleotides from the 23S ribosomal RNA gene from *E. coli* (EHEC) ATCC43888. Additionally, the 5'-NGG protospacer adjacent motif (PAM) was located adjacent to the selected target sequence. The selected target sequence in the 23S rRNA gene can be found in Table 3.

1.2 Characterization of the CGV System Targeting *E. coli* (EHEC) ATCC43888.

To establish the CGV system functionality in mediating sequence-specific killing in *E. coli* (EHEC) ATCC43888, the system was transformed into *E. coli* (EHEC) ATCC43888 cells. Overnight cultures were diluted 1:100 in fresh lysogeny broth (LB) and grown to mid-exponential phase OD600 ~0.6. The CRISPR system was induced by adding theophylline and arabinose (2 mM theophylline and 1% arabinose), and survival of the strain was followed over time by plating the cultures in serial dilutions every 15 minutes, for 1 h (FIG. 1B). CRISPR induction in *E. coli* (EHEC) surprisingly triggered a rapid killing of the cells, achieving 99.98% killing within 30 minutes of induction (FIG. 1A).

Example 2. In Vivo CRISPR Killing of Target Strain Enterohemorrhagic *E. coli* (EHEC) in *Galleria mellonella* Larvae In Vivo Infection Model

2.1. CRISPR Efficacy Against *E. coli* (EHEC) ATCC43888 Infections in *Galleria mellonella*

CRISPR killing of target strain *E. coli* (EHEC) ATCC43888 was tested in *G. mellonella* in vivo infection model. To this aim, *G. mellonella* larvae were delivered injections of bacteria 10⁸ CFU *E. coli* (EHEC) ATCC43888 behind the final left proleg. Approximately 1 h after the injection, CRISPR inducers (2 mM theophylline and 1% arabinose) were administered behind the final right proleg. Larvae were incubated at 37° C. and they were sacrificed

after 1 and 2 h after induction. As shown in FIGS. 2 and 3, CRISPR induction killed 99% of the population after 2 h, as compared to the off-target control.

2.2. Survival Curves of *G. mellonella* Larvae Infected with Enterohemorrhagic *E. coli* (EHEC).

G. mellonella larvae were delivered injections of bacteria (8×10^4 CFU *E. coli* ATCC43888) behind the final left proleg. Approximately 1 h after the injection, CRISPR inducers (2 mM theophylline and 1% arabinose) were administered behind the final right proleg. Larvae were incubated at 37° C. and survival was monitored for 115 h, with death indicated by lack of movement and unresponsiveness to touch. CRISPR killing of target strain *E. coli* (EHEC) ATCC43888 in *G. mellonella* larvae significantly improved survival of the larvae compared to the off-target control (FIG. 4) (log-rank test, $P < 0.03$).

Example 3. Precision Killing of Target Strain Probiotic *E. coli* Nissle 1917

3.1. Design, Construction and Delivery of CRISPR Guided Vector (CGV) System Targeting *E. coli* Nissle 1917.

The invention provides a CGV system to specifically target *E. coli* Nissle 1917. The CGV system comprises two vectors: (a) a vector containing a tracrRNA and the Cas9 from *Streptococcus pyogenes* (SpCas9); (b) a vector containing a guide RNA (gRNA) that comprises a nucleotide sequence capable of hybridizing to a target sequence in the host cells to guide SpCas9 to the target sequence. To enable specific killing of *E. coli* Nissle 1917, a specific sequence from the genome of this strain was chosen to target. Specifically, the sequence contains 20 nucleotides from the *pks* gene from *E. coli* Nissle 1917. Additionally, the 5'-NGG protospacer adjacent motif (PAM) was located adjacent to the selected target sequence. The selected target sequence in the *pks* gene can be found in Table 3.

Furthermore, a different genome target was selected to specifically kill *E. coli* Nissle 1917. The sequence contains 20 nucleotides from the *yapH* gene. Additionally, the 5'-NGG protospacer adjacent motif (PAM) was located adjacent to the selected target sequence. The selected target sequence in *yapH* gene can be found in Table 3.

3.2. Construction and Delivery of CRISPR Guided Vectors (CGV) Targeting *E. coli* Nissle 1917

To establish CGVs functionality in mediating sequence-specific killing in *E. coli* Nissle 1917, the CGV system was transformed into *E. coli* Nissle 1917 cells. Overnight cultures were diluted 1:100 in fresh lysogeny broth (LB) and grown to mid-exponential phase OD₆₀₀ ~0.6. The CRISPR system was induced by adding theophylline and arabinose (2 mM theophylline and 1% arabinose), and survival of the strain was followed over time by plating the cultures in serial dilutions every 15 minutes, for 3 h (FIGS. 5B and 6B). FIGS. 5B and 6A show CRISPR killing assay in *E. coli* Nissle 1917, targeting *pks* gene and *yapH* gene, respectively. In both cases, CRISPR induction triggers a rapid killing of *E. coli* Nissle 1917 cells, achieving 99.98% killing within only 15 minutes of induction.

Example 4. In Vitro CRISPR Killing of *Clostridium difficile* by Conjugative Plasmid Vectors and Cas3

This experiment involves the precision killing of *Clostridium difficile* using a gRNA-encoding CRISPR array that is delivered from a probiotic carrier bacterial species by conjugative plasmids as vectors (which we call CRISPR

guided vectors (CGV™)). A carrier bacterium (*E. coli* donor strain containing the CRISPR guided vector (CGV™)) was mated with *Clostridium difficile* which was killed upon delivery of the CGV™ containing the designed array. This CGV™ harnessed the endogenous Cas3 machinery of *Clostridium difficile* 630Aerm. A 100% killing of *Clostridium difficile* cells was achieved.

Introduction

Clostridium difficile (*C. difficile*) is a spore-forming human opportunistic pathogen that can asymptotically colonize the intestine of healthy individuals. The two main risk factors for contracting *C. difficile*-associated diseases, such as nosocomial diarrhea, are age and antibiotic treatment and can have fatal consequences. *C. difficile* 630Aerm, the subject of our study, is a well-characterized strain and is widely used for the generation of mutant specimens.

Study Objectives

Objective 1: Delivery of CGVs by Conjugation.

A CRISPR guided vector (CGV) containing an array to specifically target and kill *C. difficile* was designed and assembled. The same CGV lacking the array was assembled to use as a control for conjugation efficiency. Both CGVs were transformed into the carrier strain *Escherichia coli* CA434, which was used as a donor strain to conjugate the plasmid into our strain of interest *C. difficile* 630Aerm.

Objective 2: Harnessing *Clostridium difficile* Endogenous Cas3 Machinery.

Upon transcription of the delivered CRISPR array in the recipient target strain *C. difficile*, the endogenous Cas3 was guided to cut its own DNA; leading to bacterial death.

Objective 3: Eradication of *Clostridium difficile* 630Aerm.

Achievement of efficient killing of transconjugant *C. difficile* cells using designed CGVs.

Materials and Methods

Bacterial Strains and Growth Conditions

E. coli strain CA434 was acquired from Chain Biotech. It was cultured on nutrient-rich media (2×YT) and grown overnight at 37° C. and 250 rpm. Medium was supplemented with 12.5 µg/mL of thiamphenicol when required to maintain the CGVs.

Clostridium difficile 630Aerm was grown on BHI agar supplemented with 5 g/L of yeast extract, 0.03% L-cysteine, 250 µg/ml D-cycloserine and 8 µg/ml of cefoxitin (BHIS+CC). *C. difficile* was grown overnight in a Coy vinyl anaerobic cabinet in an atmosphere of 92% N₂, 6% CO₂ and 2% H₂ at 37° C. The mating of the donor CA434 and *C. difficile* was grown on plain BHI agar to allow for growth of the donor strain. Thiamphenicol was added to BHIS+CC plates to a final concentration of 12.5 µg/mL for selection of transconjugants after mating. All plates were dried for 1.5 hours and transferred, along with the broth version of this medium, to the anaerobic chamber at least 3 hours before use.

CGV Transfer Procedures

Carrier cells of *E. coli* CA434 were obtained by electroporation of either of our CGVs (control vector pMTL84151-FJ797649 and CRISPR vector pMTL84151-cdCRISPR1). In order to do that, overnight cultures of *E. coli* CA434 were diluted 1:100 in fresh 2× YT medium without selection and grown to OD₆₀₀ ~0.5. Then, they were made electrocompetent by standard procedures (Sharan et al., 2009). Electrocompetent cells were transformed with either plasmid pMTL84151-FJ797649 or pMTL84151-cdCRISPR1 and recovered in 2×YT for 1 h at 37° C. with shaking (250 rpm). Finally, they were plated on LB agar supplemented with 12.5 µg/mL thiamphenicol for selection of transformants. Transformants were grown in

liquid 2xYT supplemented with 12.5 gg/mL thiamphenicol at 37° C. and 250 rpm for mating with *C. difficile*. 1 ml of donor cells was centrifuged at 4000xg for 2 minutes, supernatant removed and carefully washed with 400 l of PBS. After a second centrifugation cycle the pellet was transferred to the anaerobic chamber for mating with *C. difficile* in BHI non-selective plates. *C. difficile* was prepared for mating following a modified protocol (Des Purdy et al., 2002). *C. difficile* 630Δerm was incubated overnight in selective BHIS+CC plates, from which, a scrape was inoculated overnight in 1 ml of non-selective BHI and incubated overnight for mating. 200 μl of that culture was used to resuspend the pelleted donor cells and mixed culture was plated in 20 μl spots on top of non-selective BHI plates. The mating was incubated 24h to allow for conjugation. After incubation, the whole plate was thoroughly scraped with a sterile inoculation loop, resuspended in BHI and serial dilutions were plated on BHI+CC plates to prevent growth of donor *E. coli* and on BHI+CC supplemented with thiamphenicol for additional selection of transconjugants. Single colonies were counted after 48 hours.

Results

Replicates of BHI+CC+Thiamphenicol plates, selecting for *C. difficile* transconjugants carrying the control CGV, showed a consistent number of colonies resulting in about ~600-750 CFUs per mating experiment. For the

mating of *C. difficile* with *E. coli* CA434 carrying the CGV with the CRISPR array the plates were empty, no colonies were observed. This translates into 100% killing of transconjugant *C. difficile* 630Δerm cells receiving the CRISPR array (see FIG. 7: Killing of transconjugant *C. difficile* 630Δerm).

DISCUSSION AND CONCLUSIONS

The results of this experiment show that we could successfully conjugate CGVs containing the desired CRISPR arrays into *C. difficile* 630Δerm from an *E. coli* carrier bacterium. We could also successfully harness *C. difficile* endogenous Cas3 machinery for very efficient CRISPR killing.

REFERENCES

- Purdy D, O'Keeffe T A, Elmore M, Herbert M, McLeod A, Bokori-Brown M, Ostrowski A, Minton N P. (2002) Conjugative transfer of clostridial shuttle vectors from *Escherichia coli* to *Clostridium difficile* through circumvention of the restriction barrier. *Molec. Microbiology* 46(2), 439-452
- Sharan, S. K., Thomason, L. C., Kuznetsov, S. G., and Court, D. L. (2009) Recombineering: a homologous recombination-based method of genetic engineering. *Nat. Protoc.* 4, 206-223

TABLE 1

Example Bacteria
Optionally, the bacteria are selected from this Table.

<i>Abiotrophia</i>	<i>Acidocella</i>	<i>Actinomyces</i>	<i>Alkalilimnicola</i>	<i>Aquaspirillum</i>
<i>Abiotrophia defectiva</i>	<i>Acidocella aminolytica</i>	<i>Actinomyces bovis</i>	<i>Alkalilimnicola ehrlichii</i>	<i>Aquaspirillum polymorphum</i>
<i>Acaricomes</i>	<i>Acidocella facilis</i>	<i>Actinomyces denticolens</i>	<i>Alkaliphilus</i>	<i>Aquaspirillum</i>
<i>Acaricomes phytoseveli</i>	<i>Acidomonas</i>	<i>Actinomyces europaeus</i>	<i>Alkaliphilus oremlandii</i>	<i>putridiconchylum</i>
<i>Acetitomaculum</i>	<i>Acidomonas methanolica</i>	<i>Actinomyces georgiae</i>	<i>Alkaliphilus transvaalensis</i>	<i>Aquaspirillum serpens</i>
<i>Acetitomaculum ruminis</i>	<i>Acidothermus</i>	<i>Actinomyces gerencseriae</i>	<i>Allochromatium</i>	<i>Aquimarina</i>
<i>Acetivibrio</i>	<i>Acidothermus cellulolyticus</i>	<i>Actinomyces</i>	<i>Allochromatium vinosum</i>	<i>Aquimarina latercula</i>
<i>Acetivibrio cellulolyticus</i>	<i>Acidovorax</i>	<i>hordeovulneris</i>	<i>Alloiooccus</i>	<i>Arcanobacterium</i>
<i>Acetivibrio ethanolgignens</i>	<i>Acidovorax anthurii</i>	<i>Actinomyces howellii</i>	<i>Alloiooccus otitis</i>	<i>Arcanobacterium</i>
<i>Acetivibrio multivorans</i>	<i>Acidovorax caeni</i>	<i>Actinomyces hyovaginalis</i>	<i>Allokutzneria</i>	<i>haemolyticum</i>
<i>Acetoanaerobium</i>	<i>Acidovorax cattleyae</i>	<i>Actinomyces israelii</i>	<i>Allokutzneria alбата</i>	<i>Arcanobacterium pyogenes</i>
<i>Acetoanaerobium noterae</i>	<i>Acidovorax citrulli</i>	<i>Actinomyces johnsonii</i>	<i>Altererythrobacter</i>	<i>Archangium</i>
<i>Acetobacter</i>	<i>Acidovorax defluvii</i>	<i>Actinomyces meyeri</i>	<i>Altererythrobacter</i>	<i>Archangium gephyra</i>
<i>Acetobacter acetii</i>	<i>Acidovorax delafieldii</i>	<i>Actinomyces naeslundii</i>	<i>ishigakiensis</i>	<i>Arcobacter</i>
<i>Acetobacter cerevisiae</i>	<i>Acidovorax facilis</i>	<i>Actinomyces neuui</i>	<i>Altermonas</i>	<i>Arcobacter butzleri</i>
<i>Acetobacter cibinongensis</i>	<i>Acidovorax konjaci</i>	<i>Actinomyces odontolyticus</i>	<i>Altermonas haloplanktis</i>	<i>Arcobacter cryaerophilus</i>
<i>Acetobacter estunensis</i>	<i>Acidovorax temperans</i>	<i>Actinomyces oris</i>	<i>Altermonas macleodii</i>	<i>Arcobacter halophilus</i>
<i>Acetobacter fabarum</i>	<i>Acidovorax valerianellae</i>	<i>Actinomyces radingae</i>	<i>Alysiella</i>	<i>Arcobacter nitrofigilis</i>
<i>Acetobacter ghanensis</i>	<i>Acinetobacter</i>	<i>Actinomyces slackii</i>	<i>Alysiella crassa</i>	<i>Arcobacter skirrowii</i>
<i>Acetobacter indonesiensis</i>	<i>Acinetobacter baumannii</i>	<i>Actinomyces turicensis</i>	<i>Alysiella filiformis</i>	<i>Arhodomonas</i>
<i>Acetobacter lovaniensis</i>	<i>Acinetobacter baylyi</i>	<i>Actinomyces viscosus</i>	<i>Aminobacter</i>	<i>Arhodomonas aquaeolei</i>
<i>Acetobacter malorum</i>	<i>Acinetobacter bouvetii</i>	<i>Actinoplanes</i>	<i>Aminobacter aganoensis</i>	<i>Arsenophonus</i>
<i>Acetobacter nitrogenifigens</i>	<i>Acinetobacter calcoaceticus</i>	<i>Actinoplanes auranticolor</i>	<i>Aminobacter aminovorans</i>	<i>Arsenophonus nasoniae</i>
<i>Acetobacter oeni</i>	<i>Acinetobacter gernerii</i>	<i>Actinoplanes brasiliensis</i>	<i>Aminobacter niigataensis</i>	<i>Arthrobacter</i>
<i>Acetobacter orientalis</i>	<i>Acinetobacter haemolyticus</i>	<i>Actinoplanes consettenis</i>	<i>Aminobacterium</i>	<i>Arthrobacter agilis</i>
<i>Acetobacter orleanensis</i>	<i>Acinetobacter johnsonii</i>	<i>Actinoplanes deccanensis</i>	<i>Aminobacterium mobile</i>	<i>Arthrobacter albus</i>
<i>Acetobacter pasteurianus</i>	<i>Acinetobacter junii</i>	<i>Actinoplanes derwentensis</i>	<i>Aminomonas</i>	<i>Arthrobacter aureescens</i>
<i>Acetobacter pomorum</i>	<i>Acinetobacter lwoffi</i>	<i>Actinoplanes digitatis</i>	<i>Aminomonas paucivorans</i>	<i>Arthrobacter</i>
<i>Acetobacter senegalensis</i>	<i>Acinetobacter parvus</i>	<i>Actinoplanes dirhamensis</i>	<i>Ammoniphilus</i>	<i>chlorophenolicus</i>
<i>Acetobacter xylinus</i>	<i>Acinetobacter radioresistens</i>	<i>Actinoplanes ferrugineus</i>	<i>Ammoniphilus oxalaticus</i>	<i>Arthrobacter citreus</i>
<i>Acetobacterium</i>	<i>Acinetobacter schindleri</i>	<i>Actinoplanes globisporus</i>	<i>Ammoniphilus oxalivorans</i>	<i>Arthrobacter crystallopietes</i>
<i>Acetobacterium bakkii</i>	<i>Acinetobacter soli</i>	<i>Actinoplanes humidus</i>	<i>Amphibacillus</i>	<i>Arthrobacter cummingsii</i>
<i>Acetobacterium carbinolicum</i>	<i>Acinetobacter tandooii</i>	<i>Actinoplanes italicus</i>	<i>Amphibacillus xylanus</i>	<i>Arthrobacter globiformis</i>
<i>Acetobacterium dehalogenans</i>	<i>Acinetobacter tjembergiae</i>	<i>Actinoplanes liguricus</i>	<i>Amphritea</i>	<i>Arthrobacter</i>
<i>Acetobacterium fimetarium</i>	<i>Acinetobacter townneri</i>	<i>Actinoplanes lobatus</i>	<i>Amphritea balanae</i>	<i>histidinovorans</i>
<i>Acetobacterium malicum</i>	<i>Acinetobacter ursingii</i>	<i>Actinoplanes missouriensis</i>	<i>Amphritea japonica</i>	<i>Arthrobacter ilicis</i>
<i>Acetobacterium paludosum</i>	<i>Acinetobacter venetianus</i>	<i>Actinoplanes palleronii</i>	<i>Amycolatopsis</i>	<i>Arthrobacter luteus</i>
<i>Acetobacterium tumdrae</i>	<i>Acrocarpospora</i>	<i>Actinoplanes philippinensis</i>	<i>Amycolatopsis alba</i>	<i>Arthrobacter methylotrophus</i>
<i>Acetobacterium wieringae</i>	<i>Acrocarpospora corrugata</i>	<i>Actinoplanes rectilineatus</i>	<i>Amycolatopsis albidoflavus</i>	<i>Arthrobacter mysorens</i>
<i>Acetobacterium woodii</i>	<i>Acrocarpospora</i>	<i>Actinoplanes regularis</i>	<i>Amycolatopsis azurea</i>	<i>Arthrobacter nicotianae</i>
<i>Acetofilamentum</i>	<i>macrocephala</i>	<i>Actinoplanes</i>	<i>Amycolatopsis coloradensis</i>	<i>Arthrobacter nicotinovorans</i>
<i>Acetofilamentum rigidum</i>	<i>Acrocarpospora</i>	<i>teichomyceticus</i>	<i>Amycolatopsis lurida</i>	<i>Arthrobacter oxydans</i>

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>Acetohalobium</i>	<i>pleiomorpha</i>	<i>Actinoplanes utahensis</i>	<i>Amycolatopsis mediterranei</i>	<i>Arthrobacter pascens</i>
<i>Acetohalobium arabaticum</i>	<i>Actibacter</i>	<i>Actinopolyspora</i>	<i>Amycolatopsis rifamycinica</i>	<i>Arthrobacter</i>
<i>Acetomicrobium</i>	<i>Actibacter sediminis</i>	<i>Actinopolyspora halophila</i>	<i>Amycolatopsis rubida</i>	<i>phenanthrenivorans</i>
<i>Acetomicrobium faecale</i>	<i>Actinoalloteichus</i>	<i>Actinopolyspora</i>	<i>Amycolatopsis sulphurea</i>	<i>Arthrobacter</i>
<i>Acetomicrobium flavidum</i>	<i>Actinoalloteichus</i>	<i>mortivallis</i>	<i>Amycolatopsis tolypomycina</i>	<i>polychromogenes</i>
<i>Acetonema</i>	<i>cyanogriseus</i>	<i>Actinosynnema</i>	<i>Anabaena</i>	<i>Arthrobacter protophormiae</i>
<i>Acetonema longum</i>	<i>Actinoalloteichus</i>	<i>Actinosynnema mirum</i>	<i>Anabaena cylindrica</i>	<i>Arthrobacter</i>
<i>Acetothermus</i>	<i>hymeniacidonis</i>	<i>Actinotalea</i>	<i>Anabaena flos-aquae</i>	<i>psychroactophilus</i>
<i>Acetothermus paucivorans</i>	<i>Actinoalloteichus spitiensis</i>	<i>Actinotalea fermentans</i>	<i>Anabaena variabilis</i>	<i>Arthrobacter ramosus</i>
<i>Acholeplasma</i>	<i>Actinobacillus</i>	<i>Aerococcus</i>	<i>Anaerococcus</i>	<i>Arthrobacter sulfonivorans</i>
<i>Acholeplasma axanthum</i>	<i>Actinobacillus capsulatus</i>	<i>Aerococcus sanguinicola</i>	<i>Anaerococcus burkinensis</i>	<i>Arthrobacter sulfureus</i>
<i>Acholeplasma brassicae</i>	<i>Actinobacillus delphinicola</i>	<i>Aerococcus urinae</i>	<i>Anaerobaculum</i>	<i>Arthrobacter uratoxydans</i>
<i>Acholeplasma cavigenitalium</i>	<i>Actinobacillus hominis</i>	<i>Aerococcus urinaequi</i>	<i>Anaerobaculum mobile</i>	<i>Arthrobacter ureafaciens</i>
<i>Acholeplasma equifetale</i>	<i>Actinobacillus indolicus</i>	<i>Aerococcus urinaehominis</i>	<i>Anaerobiospirillum</i>	<i>Arthrobacter viscosus</i>
<i>Acholeplasma granularum</i>	<i>Actinobacillus lignieresii</i>	<i>Aerococcus viridans</i>	<i>Anaerobiospirillum</i>	<i>Arthrobacter woluwensis</i>
<i>Acholeplasma hippikon</i>	<i>Actinobacillus minor</i>	<i>Aeromicrobium</i>	<i>succiniciproducens</i>	<i>Asaia</i>
<i>Acholeplasma laidlawii</i>	<i>Actinobacillus muris</i>	<i>Aeromicrobium erythreum</i>	<i>Anaerobiospirillum thomasi</i>	<i>Asaia bogorensis</i>
<i>Acholeplasma modicum</i>	<i>Actinobacillus</i>	<i>Aeromonas</i>	<i>Anaerococcus</i>	<i>Asanoa</i>
<i>Acholeplasma morum</i>	<i>pleuropneumoniae</i>	<i>Aeromonas</i>	<i>Anaerococcus hydrogenalis</i>	<i>Asanoa ferruginea</i>
<i>Acholeplasma multilocale</i>	<i>Actinobacillus porcinus</i>	<i>allosaccharophila</i>	<i>Anaerococcus lactolyticus</i>	<i>Asticcacaulis</i>
<i>Acholeplasma oculi</i>	<i>Actinobacillus rossii</i>	<i>Aeromonas bestiarum</i>	<i>Anaerococcus prevotii</i>	<i>Asticcacaulis biprosthecium</i>
<i>Acholeplasma palmae</i>	<i>Actinobacillus scotiae</i>	<i>Aeromonas caviae</i>	<i>Anaerococcus tetradus</i>	<i>Asticcacaulis excentricus</i>
<i>Acholeplasma parvum</i>	<i>Actinobacillus seminis</i>	<i>Aeromonas encheleia</i>	<i>Anaerococcus vaginalis</i>	<i>Atopobacter</i>
<i>Acholeplasma pleciae</i>	<i>Actinobacillus</i>	<i>Aeromonas</i>	<i>Anaerofustis</i>	<i>Atopobacter phocae</i>
	<i>succinogenes</i>	<i>enteropelogenes</i>		
<i>Acholeplasma vituli</i>	<i>Actinobacillus suis</i>	<i>Aeromonas eucrenophila</i>	<i>Anaerofustis stercorihominis</i>	<i>Atopobium</i>
<i>Achromobacter</i>	<i>Actinobacillus ureae</i>	<i>Aeromonas ichthiosmia</i>	<i>Anaeromusa</i>	<i>Atopobium fossor</i>
<i>Achromobacter denitrificans</i>	<i>Actinobaculum</i>	<i>Aeromonas jandaei</i>	<i>Anaeromusa acidaminophila</i>	<i>Atopobium minutum</i>
<i>Achromobacter insolitus</i>	<i>Actinobaculum massiliense</i>	<i>Aeromonas media</i>	<i>Anaeromyxobacter</i>	<i>Atopobium parvulum</i>
<i>Achromobacter piechaudii</i>	<i>Actinobaculum schaalii</i>	<i>Aeromonas popoffii</i>	<i>Anaeromyxobacter</i>	<i>Atopobium rimae</i>
<i>Achromobacter ruhlandii</i>	<i>Actinobaculum suis</i>	<i>Aeromonas sobria</i>	<i>dehalogenans</i>	<i>Atopobium vaginae</i>
<i>Achromobacter spanius</i>	<i>Actinomyces urinale</i>	<i>Aeromonas veronii</i>	<i>Anaerorhabdus</i>	<i>Aureobacterium</i>
<i>Acidaminobacter</i>	<i>Actinocatenispora</i>	<i>Agrobacterium</i>	<i>Anaerorhabdus furcosa</i>	<i>Aureobacterium barkeri</i>
<i>Acidaminobacter</i>	<i>Actinocatenispora rupis</i>	<i>Agrobacterium</i>	<i>Anaerostinus</i>	<i>Aureobacterium</i>
<i>hydrogenoformans</i>	<i>Actinocatenispora</i>	<i>gelatinovorum</i>	<i>Anaerostinus glycerini</i>	<i>Aureobacterium liquefaciens</i>
<i>Acidaminococcus</i>	<i>thailandica</i>	<i>Agrococcus</i>	<i>Anaerovirgula</i>	<i>Avibacterium</i>
<i>Acidaminococcus fermentans</i>	<i>Actinocatenispora sera</i>	<i>Agrococcus citreus</i>	<i>Anaerovirgula multivorans</i>	<i>Avibacterium avium</i>
<i>Acidaminococcus intestini</i>	<i>Actinocorallia</i>	<i>Agrococcus jenensis</i>	<i>Ancalomicrobium</i>	<i>Avibacterium gallinarum</i>
<i>Acidicaldus</i>	<i>Actinocorallia aurantiaca</i>	<i>Agromonas</i>	<i>Ancalomicrobium adetum</i>	<i>Avibacterium paragallinarum</i>
<i>Acidicaldus organivorans</i>	<i>Actinocorallia aurea</i>	<i>Agromonas oligotrophica</i>	<i>Ancylobacter</i>	<i>Avibacterium volantium</i>
<i>Acidimicrobium</i>	<i>Actinocorallia cavernae</i>	<i>Agromyces</i>	<i>Ancylobacter aquaticus</i>	<i>Azoarcus</i>
<i>Acidimicrobium ferrooxidans</i>	<i>Actinocorallia glomerata</i>	<i>Agromyces fucosus</i>	<i>Aneurinibacillus</i>	<i>Azoarcus indigenus</i>
<i>Acidiphilium</i>	<i>Actinocorallia herbida</i>	<i>Agromyces hippuratus</i>	<i>Aneurinibacillus</i>	<i>Azoarcus toluyliticus</i>
<i>Acidiphilium acidophilum</i>	<i>Actinocorallia libanotica</i>	<i>Agromyces luteolus</i>	<i>aneuriniyticus</i>	<i>Azoarcus toluvorans</i>
<i>Acidiphilium angustum</i>	<i>Actinocorallia longicatena</i>	<i>Agromyces mediolanus</i>	<i>Aneurinibacillus migulanus</i>	<i>Azohydromonas</i>
<i>Acidiphilium cryptum</i>	<i>Actinomadura</i>	<i>Agromyces ramosus</i>	<i>Aneurinibacillus</i>	<i>Azohydromonas australica</i>
<i>Acidiphilium multivorum</i>	<i>Actinomadura alba</i>	<i>Agromyces rhizosphaerae</i>	<i>thermoaerophilus</i>	<i>Azohydromonas lata</i>
<i>Acidiphilium organovorum</i>	<i>Actinomadura atramentaria</i>	<i>Akkermansia</i>	<i>Angiococcus</i>	<i>Azomonas</i>
<i>Acidiphilium rubrum</i>	<i>Actinomadura</i>	<i>Akkermansia muciniphila</i>	<i>Angiococcus disciformis</i>	<i>Azomonas agilis</i>
<i>Acidisoma</i>	<i>bangladeshensis</i>	<i>Albidiferax</i>	<i>Angulomicrobium</i>	<i>Azomonas insignis</i>
<i>Acidisoma sibiricum</i>	<i>Actinomadura</i>		<i>Angulomicrobium tetraedrale</i>	<i>Azomonas macrocytogenes</i>
	<i>catellatispora</i>			
<i>Acidisoma tundrae</i>	<i>Actinomadura chibensis</i>	<i>Albidiferax ferrereducens</i>	<i>Anoxybacillus</i>	<i>Azorhizobium</i>
<i>Acidisphaera</i>	<i>Actinomadura chokoriensis</i>	<i>Albidovulum</i>	<i>Anoxybacillus pushchinoensis</i>	<i>Azorhizobium caulinodans</i>
<i>Acidisphaera rubrifaciens</i>	<i>Actinomadura citrea</i>	<i>Albidovulum inexpectatum</i>	<i>Aquabacterium</i>	<i>Azorhizophilus</i>
<i>Acidithiobacillus</i>	<i>Actinomadura coerulea</i>	<i>Alcaligenes</i>	<i>Aquabacterium commune</i>	<i>Azorhizophilus paspali</i>
<i>Acidithiobacillus albertensis</i>	<i>Actinomadura echinospora</i>	<i>Alcaligenes denitrificans</i>	<i>Aquabacterium parvum</i>	<i>Azospirillum</i>
<i>Acidithiobacillus caldus</i>	<i>Actinomadura fibrosa</i>	<i>Alcaligenes faecalis</i>		<i>Azospirillum brasilense</i>
<i>Acidithiobacillus ferrooxidans</i>	<i>Actinomadura formosensis</i>	<i>Alcanivorax</i>		<i>Azospirillum halopraeferens</i>
<i>Acidithiobacillus thiooxidans</i>	<i>Actinomadura hibisca</i>	<i>Alcanivorax borkumensis</i>		<i>Azospirillum irakense</i>
<i>Acidobacterium</i>	<i>Actinomadura kijaniata</i>	<i>Alcanivorax jadensis</i>		<i>Azotobacter</i>
<i>Acidobacterium capsulatum</i>	<i>Actinomadura latina</i>	<i>Algicola</i>		<i>Azotobacter beijerinckii</i>
	<i>Actinomadura livida</i>	<i>Algicola bacteriolytica</i>		<i>Azotobacter chroococcum</i>
	<i>Actinomadura</i>	<i>Alicyclobacillus</i>		<i>Azotobacter nigricans</i>
	<i>luteofluorescens</i>	<i>Alicyclobacillus</i>		<i>Azotobacter salmestris</i>
	<i>Actinomadura macra</i>	<i>disulfidooxidans</i>		<i>Azotobacter vinelandii</i>
	<i>Actinomadura madurae</i>	<i>Alicyclobacillus</i>		
	<i>Actinomadura oligospora</i>	<i>sendaiensis</i>		
	<i>Actinomadura pelletieri</i>	<i>Alicyclobacillus vulcanalis</i>		
	<i>Actinomadura rubrobrunea</i>	<i>Alisewanella</i>		
	<i>Actinomadura</i>	<i>Alisewanella fetalis</i>		
	<i>rugatobispora</i>			
	<i>Actinomadura umbrina</i>	<i>Alkalibacillus</i>		
	<i>Actinomadura</i>	<i>Alkalibacillus</i>		
	<i>verrucosospora</i>	<i>haloalkaliphilus</i>		
	<i>Actinomadura vinacea</i>			
	<i>Actinomadura viridilutea</i>			

TABLE 1-continued

Example Bacteria			
Optionally, the bacteria are selected from this Table.			
<i>Actinomadura viridis</i>			
<i>Actinomadura yumaensis</i>			
<i>Bacillus</i>			
[see below]			
<i>Bacteriovorax</i>			
<i>Bacteriovorax stolpii</i>			
<i>Bacteroides</i>	<i>Bibersteinia</i>	<i>Borrelia</i>	<i>Brevinema</i>
<i>Bacteroides caccae</i>	<i>Bibersteinia trehalosi</i>	<i>Borrelia afzelii</i>	<i>Brevinema andersonii</i>
<i>Bacteroides coagulans</i>	<i>Bifidobacterium</i>	<i>Borrelia americana</i>	<i>Brevundimonas</i>
<i>Bacteroides eggerthii</i>	<i>Bifidobacterium</i>	<i>Borrelia burgdorferi</i>	<i>Brevundimonas alba</i>
	<i>adolescentis</i>		
<i>Bacteroides fragilis</i>	<i>Bifidobacterium angulatum</i>	<i>Borrelia carolinensis</i>	<i>Brevundimonas aurantiaca</i>
<i>Bacteroides galacturonicus</i>	<i>Bifidobacterium animalis</i>	<i>Borrelia coriacea</i>	<i>Brevundimonas diminuta</i>
<i>Bacteroides helcogenes</i>	<i>Bifidobacterium asteroides</i>	<i>Borrelia garinii</i>	<i>Brevundimonas intermedia</i>
<i>Bacteroides ovatus</i>	<i>Bifidobacterium bifidum</i>	<i>Borrelia japonica</i>	<i>Brevundimonas subvibrioides</i>
<i>Bacteroides pectinophilus</i>	<i>Bifidobacterium boum</i>	<i>Bosea</i>	<i>Brevundimonas vancanneytii</i>
<i>Bacteroides pyogenes</i>	<i>Bifidobacterium breve</i>	<i>Bosea minatitlanensis</i>	<i>Brevundimonas variabilis</i>
<i>Bacteroides salyersiae</i>	<i>Bifidobacterium</i>	<i>Bosea thiooxidans</i>	<i>Brevundimonas vesicularis</i>
	<i>catenulatum</i>		
<i>Bacteroides stercoris</i>	<i>Bifidobacterium choerinum</i>	<i>Brachybacterium</i>	<i>Brochothrix</i>
<i>Bacteroides suis</i>	<i>Bifidobacterium</i>	<i>Brachybacterium</i>	<i>Brochothrix campestris</i>
	<i>coryneforme</i>		
<i>Bacteroides tectus</i>	<i>Bifidobacterium cuniculi</i>	<i>alimentarium</i>	<i>Brochothrix thermosphacta</i>
<i>Bacteroides</i>	<i>Bifidobacterium dentium</i>	<i>Brachybacterium faecium</i>	<i>Bruceella</i>
<i>thetaiotaomicron</i>			
<i>Bacteroides uniformis</i>	<i>Bifidobacterium gallicum</i>	<i>Brachybacterium</i>	<i>Bruceella canis</i>
<i>Bacteroides ureolyticus</i>	<i>Bifidobacterium gallinarum</i>	<i>paraconglomeratum</i>	<i>Bruceella neotomae</i>
<i>Bacteroides vulgatus</i>	<i>Bifidobacterium indicum</i>	<i>Brachybacterium rhamnosum</i>	<i>Bryobacter</i>
<i>Balnearium</i>	<i>Bifidobacterium longum</i>	<i>Brachybacterium</i>	<i>Bryobacter aggregatus</i>
<i>Balnearium lithotrophicum</i>	<i>Bifidobacterium</i>	<i>tyrofermentans</i>	<i>Burkholderia</i>
<i>Balneatrix</i>	<i>magnumBifidobacterium</i>	<i>Brachyspira</i>	<i>Burkholderia ambifaria</i>
<i>Balneatrix alpica</i>	<i>merycicum</i>	<i>Brachyspira alvinipulli</i>	<i>Burkholderia andropogonis</i>
<i>Balneola</i>	<i>Bifidobacterium minimum</i>	<i>Brachyspira hyodysenteriae</i>	<i>Burkholderia anthina</i>
<i>Balneola vulgaris</i>	<i>Bifidobacterium</i>	<i>Brachyspira imocens</i>	<i>Burkholderia caledonica</i>
<i>Barnesiella</i>	<i>pseudocatenulatum</i>	<i>Brachyspira murdochii</i>	<i>Burkholderia caryophylli</i>
<i>Barnesiella viscericola</i>	<i>Bifidobacterium</i>	<i>Brachyspira pilosicoli</i>	<i>Burkholderia cenocepacia</i>
<i>Bartonella</i>	<i>pseudolongum</i>	<i>Bradyrhizobium</i>	<i>Burkholderia cepacia</i>
<i>Bartonella alsatica</i>	<i>Bifidobacterium pullorum</i>	<i>Bradyrhizobium canariense</i>	<i>Burkholderia cocovenenans</i>
<i>Bartonella bacilliformis</i>	<i>Bifidobacterium</i>	<i>Bradyrhizobium elkamii</i>	<i>Burkholderia dolosa</i>
	<i>ruminantium</i>		
<i>Bartonella clarridgeiae</i>	<i>Bifidobacterium saeculare</i>	<i>Bradyrhizobium japonicum</i>	<i>Burkholderia fungorum</i>
<i>Bartonella doshaiae</i>	<i>Bifidobacterium subtile</i>	<i>Bradyrhizobium liaoningense</i>	<i>Burkholderia glathei</i>
<i>Bartonella elizabethae</i>	<i>Bifidobacterium</i>	<i>Brenneria</i>	<i>Burkholderia glumae</i>
<i>Bartonella grahamii</i>	<i>thermophilum</i>	<i>Brenneria alni</i>	<i>Burkholderia graminis</i>
<i>Bartonella henselae</i>	<i>Bifidobacterium</i>	<i>Brenneria nigrifluens</i>	<i>Burkholderia kururuiensis</i>
<i>Bartonella rochalimae</i>	<i>Bifidobacterium</i>	<i>Brenneria quercina</i>	<i>Burkholderia multivorans</i>
<i>Bartonella vinsonii</i>	<i>Biostraticola</i>	<i>Brenneria quercina</i>	<i>Burkholderia phenazinium</i>
<i>Bavariococcus</i>	<i>Biostraticola tofi</i>	<i>Brenneria salicis</i>	<i>Burkholderia plantarii</i>
<i>Bavariococcus seileri</i>	<i>Bizonia</i>	<i>Brevibacillus</i>	<i>Burkholderia pyrocinia</i>
<i>Bdellovibrio</i>	<i>Bizonia argentinensis</i>	<i>Brevibacillus agri</i>	<i>Burkholderia silvatlantica</i>
<i>Bdellovibrio bacteriovorus</i>	<i>Blastobacter</i>	<i>Brevibacillus borstelensis</i>	<i>Burkholderia stabilis</i>
<i>Bdellovibrio exovorax</i>	<i>Blastobacter capsulatus</i>	<i>Brevibacillus brevis</i>	<i>Burkholderia thailandensis</i>
<i>Beggiatoa</i>	<i>Blastobacter denitrificans</i>	<i>Brevibacillus centrosporus</i>	<i>Burkholderia tropica</i>
<i>Beggiatoa alba</i>	<i>Blastococcus</i>	<i>Brevibacillus choshinensis</i>	<i>Burkholderia unamae</i>
<i>Beijerinckia</i>	<i>Blastococcus aggregatus</i>	<i>Brevibacillus invocatus</i>	<i>Burkholderia vietnamiensis</i>
<i>Beijerinckia dextrii</i>	<i>Blastococcus saxosidens</i>	<i>Brevibacillus laterosporus</i>	<i>Buttiauxella</i>
<i>Beijerinckia fluminensis</i>	<i>Blastochloris</i>	<i>Brevibacillus parabrevis</i>	<i>Buttiauxella agrestis</i>
<i>Beijerinckia indica</i>	<i>Blastochloris viridis</i>	<i>Brevibacillus reuszeri</i>	<i>Buttiauxella brennerae</i>
<i>Beijerinckia mobilis</i>	<i>Blastomonas</i>	<i>Brevibacterium</i>	<i>Buttiauxella ferrugutiae</i>
<i>Belliella</i>	<i>Blastomonas natoria</i>	<i>Brevibacterium abidum</i>	<i>Buttiauxella gavintiae</i>
<i>Belliella baltica</i>	<i>Blastopirellula</i>	<i>Brevibacterium album</i>	<i>Buttiauxella izardii</i>
<i>Belliella</i>	<i>Blastopirellula marina</i>	<i>Brevibacterium aurantiacum</i>	<i>Buttiauxella noackiae</i>
<i>Belliella caldifistulae</i>	<i>Blautia</i>	<i>Brevibacterium celere</i>	<i>Buttiauxella warmboldiae</i>
<i>Belnapia</i>	<i>Blautia coccoides</i>	<i>Brevibacterium epidermidis</i>	<i>Butyrivibrio</i>
<i>Belnapia moabensis</i>	<i>Blautia hansenii</i>	<i>Brevibacterium</i>	<i>Butyrivibrio fibrisolvens</i>
<i>Bergeriella</i>	<i>Blautia producta</i>	<i>frigoritolerans</i>	<i>Butyrivibrio hungatei</i>
<i>Bergeriella denitrificans</i>	<i>Blautia wexlerae</i>	<i>Brevibacterium halotolerans</i>	<i>Butyrivibrio proteoclasticus</i>
<i>Beutenbergia</i>	<i>Bogoriella</i>	<i>Brevibacterium iodinum</i>	
<i>Beutenbergia cavernae</i>	<i>Bogoriella caseilytica</i>	<i>Brevibacterium linens</i>	
	<i>Bordetella</i>	<i>Brevibacterium lyticum</i>	
	<i>Bordetella avium</i>	<i>Brevibacterium mcbrellneri</i>	
	<i>Bordetella bronchiseptica</i>	<i>Brevibacterium otitidis</i>	
	<i>Bordetella hinzii</i>	<i>Brevibacterium oxydans</i>	
	<i>Bordetella holmesii</i>	<i>Brevibacterium paucivorans</i>	
	<i>Bordetella parapertussis</i>	<i>Brevibacterium stationis</i>	
	<i>Bordetella pertussis</i>		
	<i>Bordetella petrii</i>		
	<i>Bordetella trematum</i>		

TABLE 1-continued

Example Bacteria
Optionally, the bacteria are selected from this Table.

<i>Bacillus</i>				
<i>B. acidicer</i>	<i>B. aminovorans</i>	<i>B. glucanolyticus</i>	<i>B. taeanensis</i>	<i>B. lautus</i>
<i>B. acidicola</i>	<i>B. amylolyticus</i>	<i>B. gordonae</i>	<i>B. tequilensis</i>	<i>B. lehensis</i>
<i>B. acidiproducens</i>	<i>B. andreesenii</i>	<i>B. gottheilii</i>	<i>B. thermantarcticus</i>	<i>B. lentimorbus</i>
<i>B. acidocaldarius</i>	<i>B. aneurinilyticus</i>	<i>B. graminis</i>	<i>B. thermoaerophilus</i>	<i>B. lentus</i>
<i>B. acidoterrestris</i>	<i>B. anthracis</i>	<i>B. halmapalus</i>	<i>B. thermoamylovorans</i>	<i>B. licheniformis</i>
<i>B. aëolius</i>	<i>B. aquimaris</i>	<i>B. haloalkaliphilus</i>	<i>B. thermocatenulatus</i>	<i>B. ligniniphilus</i>
<i>B. aerius</i>	<i>B. arenosi</i>	<i>B. halochares</i>	<i>B. thermocloacae</i>	<i>B. litoralis</i>
<i>B. aerophilus</i>	<i>B. arseniciselenatis</i>	<i>B. halodenitrificans</i>	<i>B. thermocopriæ</i>	<i>B. localis</i>
<i>B. agaradhaerens</i>	<i>B. arsenicus</i>	<i>B. halodurans</i>	<i>B. thermodenitrificans</i>	<i>B. luciflavus</i>
<i>B. agri</i>	<i>B. aurantiacus</i>	<i>B. halophilus</i>	<i>B. thermoglucosidasius</i>	<i>B. luteolus</i>
<i>B. attingensis</i>	<i>B. arvi</i>	<i>B. halosaccharovorans</i>	<i>B. thermolactis</i>	<i>B. luteus</i>
<i>B. akibai</i>	<i>B. aryabhatai</i>	<i>B. hemicellulosilyticus</i>	<i>B. thermoleovorans</i>	<i>B. macauensis</i>
<i>B. alcalophilus</i>	<i>B. asahii</i>	<i>B. hemicentroti</i>	<i>B. thermophilus</i>	<i>B. macerans</i>
<i>B. algicola</i>	<i>B. atrophaeus</i>	<i>B. herbersteinensis</i>	<i>B. thermoruber</i>	<i>B. macquariensis</i>
<i>B. alginolyticus</i>	<i>B. axarquiensis</i>	<i>B. hemicellulosilyticus</i>	<i>B. thermosphaericus</i>	<i>B. macyae</i>
<i>B. alkalidiazotrophicus</i>	<i>B. azotofixans</i>	<i>B. horneckiae</i>	<i>B. thiaminolyticus</i>	<i>B. malacitensis</i>
<i>B. alkalinitrilicus</i>	<i>B. azotoformans</i>	<i>B. horti</i>	<i>B. thioparans</i>	<i>B. mannanilyticus</i>
<i>B. alkalisediminis</i>	<i>B. badius</i>	<i>B. huizhouensis</i>	<i>B. thuringiensis</i>	<i>B. marisflavi</i>
<i>B. alkalitelluris</i>	<i>B. barbaricus</i>	<i>B. humi</i>	<i>B. tianshenii</i>	<i>B. marismortui</i>
<i>B. altitudinis</i>	<i>B. bataviensis</i>	<i>B. hwajinpoensis</i>	<i>B. tryoxylicola</i>	<i>B. marmarensis</i>
<i>B. alveayuensis</i>	<i>B. beijingensis</i>	<i>B. idriensis</i>	<i>B. tusciae</i>	<i>B. massiliensis</i>
<i>B. alvei</i>	<i>B. benzoovorans</i>	<i>B. indicus</i>	<i>B. validus</i>	<i>B. megaterium</i>
<i>B. amyloliquefaciens</i>	<i>B. beringensis</i>	<i>B. infantis</i>	<i>B. vallismortis</i>	<i>B. mesonae</i>
<i>B.</i>	<i>B. berkeleyi</i>	<i>B. infernus</i>	<i>B. vedderi</i>	<i>B. methanolicus</i>
<i>a. subsp. amyloliquefaciens</i>	<i>B. beveridgei</i>	<i>B. insolitus</i>	<i>B. velezensis</i>	<i>B. methylotrophicus</i>
<i>B. a. subsp. plantarum</i>	<i>B. bogoriensis</i>	<i>B. invictae</i>	<i>B. viamamensis</i>	<i>B. migulanus</i>
<i>B. diposauri</i>	<i>B. boroniphilus</i>	<i>B. iranensis</i>	<i>B. vireti</i>	<i>B. mojavenis</i>
<i>B. drementis</i>	<i>B. borstelensis</i>	<i>B. isabellae</i>	<i>B. vulcani</i>	<i>B. mucilaginosus</i>
<i>B. edaphicus</i>	<i>B. brevis</i> Migula	<i>B. isronensis</i>	<i>B. wakoensis</i>	<i>B. muralis</i>
<i>B. ehimensis</i>	<i>B. butanolivorans</i>	<i>B. jeotgali</i>	<i>B. weihenstephanensis</i>	<i>B. murimartini</i>
<i>B. eiseniae</i>	<i>B. canaveralius</i>	<i>B. kaustophilus</i>	<i>B. xiamenensis</i>	<i>B. mycoideis</i>
<i>B. enclensis</i>	<i>B. carboniphilus</i>	<i>B. kobensis</i>	<i>B. xiaoxiensis</i>	<i>B. naganoensis</i>
<i>B. endophyticus</i>	<i>B. cecembensis</i>	<i>B. kochii</i>	<i>B. zhanjiangensis</i>	<i>B. nanhaiensis</i>
<i>B. endoradicis</i>	<i>B. cellulosityticus</i>	<i>B. kokeshiiformis</i>	<i>B. peoriae</i>	<i>B. nanhaiisediminis</i>
<i>B. farraginis</i>	<i>B. centrosporus</i>	<i>B. korensis</i>	<i>B. persepolensis</i>	<i>B. nealsonii</i>
<i>B. fastidiosus</i>	<i>B. cereus</i>	<i>B. korlensis</i>	<i>B. persicus</i>	<i>B. neidei</i>
<i>B. fengqiensis</i>	<i>B. chagammovensis</i>	<i>B. kribbensis</i>	<i>B. pervagus</i>	<i>B. neizhouensis</i>
<i>B. firmus</i>	<i>B. chitinolyticus</i>	<i>B. krulwichiae</i>	<i>B. plakortidis</i>	<i>B. niabensis</i>
<i>B. flexus</i>	<i>B. chondroitinus</i>	<i>B. laevolacticus</i>	<i>B. pocheonensis</i>	<i>B. niacini</i>
<i>B. foraminis</i>	<i>B. choshinensis</i>	<i>B. larvae</i>	<i>B. polygoni</i>	<i>B. novalis</i>
<i>B. fordii</i>	<i>B. chungangensis</i>	<i>B. laterosporus</i>	<i>B. polymyxa</i>	<i>B. oceanisediminis</i>
<i>B. formosus</i>	<i>B. cibi</i>	<i>B. salixigens</i>	<i>B. popilliae</i>	<i>B. odysssei</i>
<i>B. fortis</i>	<i>B. circulans</i>	<i>B. saliphilus</i>	<i>B. pseudocalophilus</i>	<i>B. okhensis</i>
<i>B. fumarioli</i>	<i>B. clarkii</i>	<i>B. schlegelii</i>	<i>B. pseudofirmus</i>	<i>B. okuhidensis</i>
<i>B. funiculus</i>	<i>B. clausii</i>	<i>B. sediminis</i>	<i>B. pseudomycoideis</i>	<i>B. oleronius</i>
<i>B. fusiformis</i>	<i>B. coagulans</i>	<i>B. selenatarsenatis</i>	<i>B. psychrodurans</i>	<i>B. oryzacorticis</i>
<i>B. galactophilus</i>	<i>B. coahuilensis</i>	<i>B. selenitireducens</i>	<i>B. psychrophilus</i>	<i>B. oshimensis</i>
<i>B. galactosidilyticus</i>	<i>B. cohmii</i>	<i>B. seohaeanensis</i>	<i>B. psychrosaccharolyticus</i>	<i>B. pabuli</i>
<i>B. galliensi</i>	<i>B. composti</i>	<i>B. shacheensis</i>	<i>B. psychrotolerans</i>	<i>B. pakistanensis</i>
<i>B. gelatini</i>	<i>B. curdlanolyticus</i>	<i>B. shackletonii</i>	<i>B. pulvifaciens</i>	<i>B. pallidus</i>
<i>B. gibsonii</i>	<i>B. cycloheptanicus</i>	<i>B. siamensis</i>	<i>B. pumilus</i>	<i>B. pallidus</i>
<i>B. ginsengi</i>	<i>B. cytotoxicus</i>	<i>B. silvestris</i>	<i>B. purgationiresistens</i>	<i>B. panacisoli</i>
<i>B. ginsengihumi</i>	<i>B. daliensis</i>	<i>B. simplex</i>	<i>B. pycnus</i>	<i>B. panaciterrae</i>
<i>B. ginsengisoli</i>	<i>B. decisifrondis</i>	<i>B. siralis</i>	<i>B. qingdaonensis</i>	<i>B. pantothenicus</i>
<i>B. globisporus</i> (eg. <i>B.</i>	<i>B. decolorationis</i>	<i>B. smithii</i>	<i>B. qingshengii</i>	<i>B. parabrevis</i>
<i>g. subsp. Globisporus</i> ; or <i>B.</i>	<i>B. deserti</i>	<i>B. soli</i>	<i>B. reuszeri</i>	<i>B. paraflexus</i>
<i>g. subsp. Marinus</i>)		<i>B. solimangrovi</i>	<i>B. rhizosphaerae</i>	<i>B. pasteurii</i>
		<i>B. solisalsi</i>	<i>B. rigui</i>	<i>B. patagoniensis</i>
		<i>B. songklensis</i>	<i>B. ruris</i>	
		<i>B. sonorensis</i>	<i>B. safensis</i>	
		<i>B. sphaericus</i>	<i>B. salarius</i>	
		<i>B. sporothermodurans</i>		
		<i>B. stearothermophilus</i>		
		<i>B. stratosphericus</i>		
		<i>B. subterraneus</i>		
		<i>B. subtilis</i> (eg. <i>B.</i>		
		<i>s. subsp. Inaquosorum</i> ;		
		or <i>B. s. subsp. Spizizeni</i> ;		
		or <i>B. s. subsp. Subtilis</i>)		
<i>Caenimonas</i>	<i>Campylobacter</i>	<i>Cardiobacterium</i>	<i>Catenuloplanes</i>	<i>Curtobacterium</i>
<i>Caenimonas korensis</i>	<i>Campylobacter coli</i>	<i>Cardiobacterium hominis</i>	<i>Catenuloplanes atrovinosus</i>	<i>Curtobacterium</i>
<i>Caldalkalibacillus</i>	<i>Campylobacter concisus</i>	<i>Caenimonas</i>	<i>Catenuloplanes castaneus</i>	<i>albidum</i>
<i>Caldalkalibacillus uzonensis</i>	<i>Campylobacter curvus</i>	<i>Caenimonas nigrificans</i>	<i>Catenuloplanes crispus</i>	<i>Curtobacterium citreus</i>
<i>Caldanaerobacter</i>	<i>Campylobacter fetus</i>	<i>Carnobacterium</i>	<i>Catenuloplanes indicus</i>	
<i>Caldanaerobacter</i>	<i>Campylobacter gracilis</i>	<i>Carnobacterium</i>	<i>Catenuloplanes japonicus</i>	
<i>subterraneus</i>				

TABLE 1-continued

Example Bacteria			
Optionally, the bacteria are selected from this Table.			
<i>Caldanaerobius</i>	<i>Campylobacter helveticus</i>	<i>alterfunditum</i>	<i>Catenuloplanes nepalensis</i>
<i>Caldanaerobius fijiensis</i>	<i>Campylobacter hominis</i>	<i>Carnobacterium divergens</i>	<i>Catenuloplanes niger</i>
<i>Caldanaerobius</i>	<i>Campylobacter</i>	<i>Carnobacterium funditum</i>	<i>Chryseobacterium</i>
	<i>hyointestinalis</i>		
<i>polysaccharolyticus</i>	<i>Campylobacter jejuni</i>	<i>Carnobacterium gallinarum</i>	<i>Chryseobacterium</i>
<i>Caldanaerobius zeae</i>	<i>Campylobacter lari</i>	<i>Carnobacterium</i>	<i>balustinum</i>
<i>Caldanaerovirga</i>	<i>Campylobacter mucosalis</i>	<i>maltaromaticum</i>	<i>Citrobacter</i>
<i>Caldanaerovirga acetigignens</i>	<i>Campylobacter rectus</i>	<i>Carnobacterium mobile</i>	<i>C. amalonaticus</i>
<i>Caldicellulosiruptor</i>	<i>Campylobacter showae</i>	<i>Carnobacterium viridans</i>	<i>C. braakii</i>
<i>Caldicellulosiruptor bescii</i>	<i>Campylobacter sputorum</i>	<i>Caryophanon</i>	<i>C. diversus</i>
<i>Caldicellulosiruptor kristjanssonii</i>	<i>Campylobacter upsaliensis</i>	<i>Caryophanon latum</i>	<i>C. farmeri</i>
<i>Caldicellulosiruptor owensensis</i>	<i>Capnocytophaga</i>	<i>Caryophanon tenue</i>	<i>C. freundii</i>
	<i>Capnocytophaga</i>	<i>Catellatospora</i>	<i>C. gillenii</i>
	<i>canimorsus</i>		
	<i>Capnocytophaga</i>	<i>Catellatospora citrea</i>	<i>C. koseri</i>
	<i>cynodegmi</i>		
	<i>Capnocytophaga gingivalis</i>	<i>Catellatospora</i>	<i>C. murlinae</i>
	<i>Capnocytophaga granulosa</i>	<i>methionotrophica</i>	<i>C. pasteurii</i> ¹¹
	<i>Capnocytophaga</i>	<i>Catenococcus</i>	<i>C. rodentium</i>
	<i>haemolytica</i>		
	<i>Capnocytophaga ochracea</i>	<i>Catenococcus thiocycli</i>	<i>C. sedlakii</i>
	<i>Capnocytophaga sputigena</i>		<i>C. werkmanii</i>
			<i>C. youngae</i>
			<i>Clostridium</i>
			(see below)
			<i>Coccochloris</i>
			<i>Coccochloris elabens</i>
			<i>Corynebacterium</i>
			<i>Corynebacterium flavescens</i>
			<i>Corynebacterium variabile</i>
<i>Clostridium</i>			
<i>Clostridium absonum</i> , <i>Clostridium aceticum</i> , <i>Clostridium acetireducens</i> , <i>Clostridium acetobutylicum</i> , <i>Clostridium acidisoli</i> , <i>Clostridium aciditolerans</i> , <i>Clostridium acidurici</i> , <i>Clostridium aerotolerans</i> , <i>Clostridium aestuarii</i> , <i>Clostridium akagii</i> , <i>Clostridium aldenense</i> , <i>Clostridium aldrichii</i> , <i>Clostridium algidicarni</i> , <i>Clostridium algidixylanolyticum</i> , <i>Clostridium algifaecis</i> , <i>Clostridium algoriphilum</i> , <i>Clostridium alkalicellulosi</i> , <i>Clostridium aminophilum</i> , <i>Clostridium aminovalericum</i> , <i>Clostridium amygdalinum</i> , <i>Clostridium amylolyticum</i> , <i>Clostridium arbuti</i> , <i>Clostridium arcticum</i> , <i>Clostridium argentinense</i> , <i>Clostridium asparagiforme</i> , <i>Clostridium aurantibutyricum</i> , <i>Clostridium autoethanogenum</i> , <i>Clostridium baratii</i> , <i>Clostridium barkeri</i> , <i>Clostridium bartlettii</i> , <i>Clostridium beijerinckii</i> , <i>Clostridium bifementans</i> , <i>Clostridium boltae</i> , <i>Clostridium bornimense</i> , <i>Clostridium botulinum</i> , <i>Clostridium bowmanii</i> , <i>Clostridium bryantii</i> , <i>Clostridium butyricum</i> , <i>Clostridium cadaveris</i> , <i>Clostridium caenicola</i> , <i>Clostridium caminitermale</i> , <i>Clostridium carboxidivorans</i> , <i>Clostridium carnis</i> , <i>Clostridium cavendishii</i> , <i>Clostridium celatum</i> , <i>Clostridium celerecrescens</i> , <i>Clostridium cellobioparum</i> , <i>Clostridium cellulofementans</i> , <i>Clostridium cellulolyticum</i> , <i>Clostridium cellulosi</i> , <i>Clostridium cellulovorans</i> , <i>Clostridium chartatabidum</i> , <i>Clostridium chauvoei</i> , <i>Clostridium chromiireducens</i> , <i>Clostridium citroniae</i> , <i>Clostridium clariflavum</i> , <i>Clostridium clostridioforme</i> , <i>Clostridium coccoides</i> , <i>Clostridium cochlearium</i> , <i>Clostridium colletant</i> , <i>Clostridium colicanis</i> , <i>Clostridium colinum</i> , <i>Clostridium collagenovorans</i> , <i>Clostridium cylindrosporium</i> , <i>Clostridium difficile</i> , <i>Clostridium diolis</i> , <i>Clostridium disporicum</i> , <i>Clostridium drakei</i> , <i>Clostridium durum</i> , <i>Clostridium estertheticum</i> , <i>Clostridium estertheticum estertheticum</i> , <i>Clostridium estertheticum laramiense</i> , <i>Clostridium fallax</i> , <i>Clostridium felsineum</i> , <i>Clostridium fervidum</i> , <i>Clostridium fimetarium</i> , <i>Clostridium formicaceticum</i> , <i>Clostridium frigidicarnis</i> , <i>Clostridium frigroris</i> , <i>Clostridium ganghwense</i> , <i>Clostridium gasigenes</i> , <i>Clostridium ghonii</i> , <i>Clostridium glycolicum</i> , <i>Clostridium glycyrrhizinilyticum</i> , <i>Clostridium grantii</i> , <i>Clostridium haemolyticum</i> , <i>Clostridium kalophilum</i> , <i>Clostridium hastiforme</i> , <i>Clostridium hathewayi</i> , <i>Clostridium herbivorans</i> , <i>Clostridium hiranonis</i> , <i>Clostridium histolyticum</i> , <i>Clostridium homopropionicum</i> , <i>Clostridium huakuii</i> , <i>Clostridium hungatei</i> , <i>Clostridium hydrogeniformans</i> , <i>Clostridium hydroxybenzoicum</i> , <i>Clostridium hylemonae</i> , <i>Clostridium jejuense</i> , <i>Clostridium indolis</i> , <i>Clostridium innocuum</i> , <i>Clostridium intestinale</i> , <i>Clostridium irregulare</i> , <i>Clostridium isatidis</i> , <i>Clostridium josui</i> , <i>Clostridium klayveri</i> , <i>Clostridium lactatifermentans</i> , <i>Clostridium lacusfryxellense</i> , <i>Clostridium laramiense</i> , <i>Clostridium lavalense</i> , <i>Clostridium lentocellum</i> , <i>Clostridium lentoputrescens</i> , <i>Clostridium leptum</i> , <i>Clostridium limosum</i> , <i>Clostridium litorale</i> , <i>Clostridium liusebunense</i> , <i>Clostridium ljungdahlii</i> , <i>Clostridium lortetii</i> , <i>Clostridium lundense</i> , <i>Clostridium magnum</i> , <i>Clostridium malenominatum</i> , <i>Clostridium mangelonii</i> , <i>Clostridium mayombeii</i> , <i>Clostridium methoxybenzovorans</i> , <i>Clostridium methylpentosum</i> , <i>Clostridium neopropionicum</i> , <i>Clostridium nexile</i> , <i>Clostridium nitrophenolicum</i> , <i>Clostridium novyi</i> , <i>Clostridium oceanicum</i> , <i>Clostridium orbiscindens</i> , <i>Clostridium oroticum</i> , <i>Clostridium oxalicum</i> , <i>Clostridium papyrosolvans</i> , <i>Clostridium paradoxum</i> , <i>Clostridium paraperfringens</i> (Alias: <i>C. welchii</i>), <i>Clostridium paraputrificum</i> , <i>Clostridium pascui</i> , <i>Clostridium pasteurianum</i> , <i>Clostridium peptidivorans</i> , <i>Clostridium perenne</i> , <i>Clostridium perfringens</i> , <i>Clostridium pfennigii</i> , <i>Clostridium phytofermentans</i> , <i>Clostridium piliforme</i> , <i>Clostridium polysaccharolyticum</i> , <i>Clostridium populei</i> , <i>Clostridium propionicum</i> , <i>Clostridium proteoclasticum</i> , <i>Clostridium proteolyticum</i> , <i>Clostridium psychrophilum</i> , <i>Clostridium puniceum</i> , <i>Clostridium purinilyticum</i> , <i>Clostridium putrefaciens</i> , <i>Clostridium putrificum</i> , <i>Clostridium quercicolum</i> , <i>Clostridium quinii</i> , <i>Clostridium ramosum</i> , <i>Clostridium rectum</i> , <i>Clostridium roseum</i> , <i>Clostridium saccharobutylicum</i> , <i>Clostridium saccharogumia</i> , <i>Clostridium saccharolyticum</i> , <i>Clostridium saccharoperbutylacetonicum</i> , <i>Clostridium sardiniense</i> , <i>Clostridium sartagoforme</i> , <i>Clostridium scatologenes</i> , <i>Clostridium schirmacherense</i> , <i>Clostridium scindens</i> , <i>Clostridium septicum</i> , <i>Clostridium sordellii</i> , <i>Clostridium sphenoides</i> , <i>Clostridium spiroforme</i> , <i>Clostridium sporogenes</i> , <i>Clostridium sporosphaeroides</i> , <i>Clostridium stercorarium</i> , <i>Clostridium stercorarium leptospartum</i> , <i>Clostridium stercorarium stercorarium</i> , <i>Clostridium stercorarium thermolacticum</i> , <i>Clostridium sticklandii</i> , <i>Clostridium straminisolvans</i> , <i>Clostridium subterminale</i> , <i>Clostridium sufflavum</i> , <i>Clostridium sulfidigenes</i> , <i>Clostridium symbiosum</i> , <i>Clostridium tagluense</i> , <i>Clostridium tepidiprofundii</i> , <i>Clostridium terminidis</i> , <i>Clostridium tertium</i> , <i>Clostridium tetani</i> , <i>Clostridium tetanomorphum</i> , <i>Clostridium thermaceticum</i> , <i>Clostridium thermautotrophicum</i> , <i>Clostridium thermoalcaliphilum</i> , <i>Clostridium thermobutyricum</i> , <i>Clostridium thermocellum</i> , <i>Clostridium thermocopriae</i> , <i>Clostridium thermohydrosulfuricum</i> , <i>Clostridium thermolacticum</i> , <i>Clostridium thermopalmarium</i> , <i>Clostridium thermopapulyticum</i> , <i>Clostridium thermosaccharolyticum</i> , <i>Clostridium thermosuccinogenes</i> , <i>Clostridium thermosulfurigenes</i> , <i>Clostridium thiosulfatireducens</i> , <i>Clostridium tyrobutyricum</i> , <i>Clostridium uliginosum</i> , <i>Clostridium ultunense</i> , <i>Clostridium villosum</i> , <i>Clostridium vincentii</i> , <i>Clostridium viride</i> , <i>Clostridium xylanolyticum</i> , <i>Clostridium xylanovorans</i>			
<i>Dactylosporangium</i>	<i>Deinococcus</i>	<i>Delftia</i>	<i>Echmicola</i>
<i>Dactylosporangium aurantiacum</i>	<i>Deinococcus aerius</i>	<i>Delftia acidovorans</i>	<i>Echmicola pacifica</i>

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>Dactylosporangium fulvum</i>	<i>Deinococcus apachenis</i>	<i>Desulfovibrio</i>	<i>Echinicola vietnamensis</i>	
<i>Dactylosporangium matsuzakiense</i>	<i>Deinococcus aquaticus</i>	<i>Desulfovibrio desulfuricans</i>		
<i>Dactylosporangium roseum</i>	<i>Deinococcus aquatilis</i>	<i>Diplococcus</i>		
<i>Dactylosporangium thailandense</i>	<i>Deinococcus caeni</i>	<i>Diplococcus pneumoniae</i>		
<i>Dactylosporangium vinaceum</i>	<i>Deinococcus radiodurans</i>			
	<i>Deinococcus radiophilus</i>			
<i>Enterobacter</i>	<i>Enterobacter kobei</i>	<i>Faecalibacterium</i>	<i>Flavobacterium</i>	
<i>E. aerogenes</i>	<i>E. ludwigii</i>	<i>Faecalibacterium prausnitzii</i>	<i>Flavobacterium antarcticum</i>	
<i>E. amnigenus</i>	<i>E. mori</i>	<i>Fangia</i>	<i>Flavobacterium aquatile</i>	
<i>E. agglomerans</i>	<i>E. nimipressuralis</i>	<i>Fangia hongkongensis</i>	<i>Flavobacterium aquidurens</i>	
<i>E. arachidis</i>	<i>E. oryzae</i>	<i>Fastidiosipila</i>	<i>Flavobacterium balustinum</i>	
<i>E. asburiae</i>	<i>E. pulveris</i>	<i>Fastidiosipila sanguinis</i>	<i>Flavobacterium croceum</i>	
<i>E. cancerogenus</i>	<i>E. pyrinus</i>	<i>Fusobacterium</i>	<i>Flavobacterium cucumis</i>	
<i>E. cloacae</i>	<i>E. radicincitans</i>	<i>Fusobacterium nucleatum</i>	<i>Flavobacterium daejeonense</i>	
<i>E. cowanii</i>	<i>E. taylorae</i>		<i>Flavobacterium defluvii</i>	
<i>E. dissolvens</i>	<i>E. turicensis</i>			
<i>E. gergoviae</i>	<i>E. sakazakii</i>			
	<i>Enterobacter soli</i>			
<i>E. helveticus</i>	<i>Enterococcus</i>		<i>Flavobacterium degerlachei</i>	
<i>E. hormaechei</i>	<i>Enterococcus durans</i>		<i>Flavobacterium denitrificans</i>	
<i>E. intermedius</i>	<i>Enterococcus faecalis</i>		<i>Flavobacterium filum</i>	
	<i>Enterococcus faecium</i>		<i>Flavobacterium flevense</i>	
	<i>Erwinia</i>		<i>Flavobacterium frigidarium</i>	
	<i>Erwinia hapontici</i>		<i>Flavobacterium mizutaii</i>	
	<i>Escherichia</i>		<i>Flavobacterium okeanokoites</i>	
	<i>Escherichia coli</i>		<i>Jamibacter</i>	
<i>Gaetbulibacter</i>	<i>Haemophilus</i>	<i>Ideonella</i>	<i>Jamibacter anophelis</i>	
<i>Gaetbulibacter saemankumensis</i>	<i>Haemophilus aegyptius</i>	<i>Ideonella azotifigens</i>		
<i>Gallibacterium</i>	<i>Haemophilus aphrophilus</i>	<i>Idiomarina</i>	<i>Jamibacter corallicola</i>	
<i>Gallibacterium anatis</i>	<i>Haemophilus felis</i>	<i>Idiomarina abyssalis</i>	<i>Jamibacter limosus</i>	
<i>Gallicola</i>	<i>Haemophilus gallinarum</i>	<i>Idiomarina baltica</i>	<i>Jamibacter melonis</i>	
<i>Gallicola barnesae</i>	<i>Haemophilus haemolyticus</i>	<i>Idiomarina fontislapidosi</i>	<i>Jamibacter terrae</i>	
<i>Garciella</i>	<i>Haemophilus influenzae</i>	<i>Idiomarina loihensis</i>	<i>Jannaschia</i>	
<i>Garciella nitratireducens</i>	<i>Haemophilus paracuniculus</i>	<i>Idiomarina ramblicola</i>	<i>Jannaschia cystaugens</i>	
<i>Geobacillus</i>	<i>Haemophilus parahaemolyticus</i>	<i>Idiomarina seosinensis</i>	<i>Jannaschia helgolandensis</i>	
	<i>Haemophilus parainfluenzae</i>	<i>Idiomarina zobellii</i>	<i>Jannaschia pohangensis</i>	
<i>Geobacillus stearothermophilus</i>	<i>Haemophilus paraphrohaemolyticus</i>	<i>Ignatzschineria</i>	<i>Jannaschia rubra</i>	
<i>Geobacter</i>	<i>Haemophilus parasuis</i>	<i>Ignatzschineria larvae</i>	<i>Janthinobacterium</i>	
<i>Geobacter bemidjiensis</i>	<i>Haemophilus pittmaniae</i>	<i>Ignavigranum</i>	<i>Janthinobacterium agaricidamnorum</i>	
<i>Geobacter bremensis</i>	<i>Hafnia</i>	<i>Ignavigranum ruoffiae</i>	<i>Janthinobacterium lividum</i>	
<i>Geobacter chapellei</i>	<i>Hafnia alvei</i>	<i>Ilumatobacter</i>	<i>Jejuia</i>	
<i>Geobacter grbiciae</i>	<i>Hahella</i>	<i>Ilumatobacter fluminis</i>	<i>Jejuia pallidilutea</i>	
<i>Geobacter hydrogenophilus</i>	<i>Hahella ganghwensis</i>	<i>Ilyobacter</i>	<i>Jeotgalibacillus</i>	
<i>Geobacter lovleyi</i>	<i>Halalkalibacillus</i>	<i>Ilyobacter delafieldii</i>	<i>Jeotgalibacillus alimentarius</i>	
<i>Geobacter metallireducens</i>	<i>Halalkalibacillus halophilus</i>	<i>Ilyobacter insuetus</i>		
<i>Geobacter pelophilus</i>	<i>Helicobacter</i>	<i>Ilyobacter polytropus</i>		
	<i>Helicobacter pylori</i>	<i>Ilyobacter tartaricus</i>	<i>Jeotgalicoccus</i>	
<i>Geobacter pickeringii</i>			<i>Jeotgalicoccus halotolerans</i>	
<i>Geobacter sulfurreducens</i>				
<i>Geodermatophilus</i>				
<i>Geodermatophilus obscurus</i>				
<i>Gluconacetobacter</i>				
<i>Gluconacetobacter xylinus</i>				
<i>Gordonia</i>				
<i>Gordonia rubripertincta</i>				
<i>Kaistia</i>	<i>Labeledella</i>	<i>Listeria ivanovii</i>	<i>Micrococcus</i>	<i>Nesterenkonia</i>
<i>Kaistia adipata</i>	<i>Labeledella gwakjiensis</i>	<i>L. marthii</i>	<i>Micrococcus luteus</i>	<i>Nesterenkonia holobia</i>
<i>Kaistia soli</i>	<i>Labrenzia</i>	<i>L. monocytogenes</i>	<i>Micrococcus lylae</i>	<i>Nocardia</i>
<i>Kangiella</i>	<i>Labrenzia aggregata</i>	<i>L. newyorkensis</i>	<i>Moraxella</i>	<i>Nocardia argentinensis</i>
<i>Kangiella aquimarina</i>	<i>Labrenzia alba</i>	<i>L. riparia</i>	<i>Moraxella bovis</i>	<i>Nocardia corallina</i>
<i>Kangiella koreensis</i>	<i>Labrenzia alexandrii</i>	<i>L. rocourtiae</i>	<i>Moraxella nonliquefaciens</i>	<i>Nocardia</i>
<i>Kersteria</i>	<i>Labrenzia marina</i>	<i>L. seeligeri</i>	<i>Moraxella osloensis</i>	<i>otitidiscaeviarum</i>
<i>Kersteria gyiorum</i>	<i>Labrys</i>	<i>L. weihenstephanensis</i>	<i>Nakamurella</i>	
<i>Kiloniella</i>	<i>Labrys methylaminiphilus</i>	<i>L. welshimeri</i>	<i>Nakamurella multipartita</i>	
<i>Kiloniella laminariae</i>	<i>Labrys miyagiensis</i>	<i>Listonella</i>	<i>Nannocystis</i>	
<i>Klebsiella</i>	<i>Labrys monachus</i>	<i>Listonella anguillarum</i>	<i>Nannocystis pusilla</i>	
<i>K. granulomatis</i>	<i>Labrys okinawensis</i>	<i>Macrocococcus</i>	<i>Natranaerobius</i>	
<i>K. oxytoca</i>	<i>Labrys portucalensis</i>	<i>Macrocococcus bovicus</i>	<i>Natranaerobius thermophilus</i>	
<i>K. pneumoniae</i>	<i>Lactobacillus</i>	<i>Marinobacter</i>		

TABLE 1-continued

Example Bacteria			
Optionally, the bacteria are selected from this Table.			
<i>K. terrigena</i>	[see below]	<i>Marinobacter algicola</i>	<i>Natronaerobius trueperi</i>
<i>K. variicola</i>	<i>Laceyella</i>	<i>Marinobacter bryozoorum</i>	<i>Naxibacter</i>
<i>Kluyvera</i>	<i>Laceyella putida</i>	<i>Marinobacter flavimaris</i>	<i>Naxibacter alkalitolerans</i>
<i>Kluyvera ascorbata</i>	<i>Lechevalieria</i>	<i>Meiothermus</i>	<i>Neisseria</i>
<i>Kocuria</i>	<i>Lechevalieria</i>	<i>Meiothermus ruber</i>	<i>Neisseria cinerea</i>
	<i>aerocolonigenes</i>		
<i>Kocuria roasea</i>	<i>Legionella</i>	<i>Methylophilus</i>	<i>Neisseria denitrificans</i>
<i>Kocuria varians</i>	[see below]	<i>Methylophilus</i>	<i>Neisseria gonorrhoeae</i>
<i>Kurthia</i>	<i>Listeria</i>	<i>methylophilus</i>	<i>Neisseria lactamica</i>
<i>Kurthia zopfii</i>	<i>L. aquatica</i>	<i>Microbacterium</i>	<i>Neisseria mucosa</i>
	<i>L. booriae</i>	<i>Microbacterium</i>	<i>Neisseria sicca</i>
	<i>L. cornellensis</i>	<i>ammoniophilum</i>	<i>Neisseria subflava</i>
	<i>L. fleischmannii</i>	<i>Microbacterium</i>	<i>Neptunomonas</i>
		<i>arborescens</i>	
	<i>L. floridensis</i>	<i>Microbacterium</i>	<i>Neptunomonas japonica</i>
		<i>liquefaciens</i>	
	<i>L. grandensis</i>	<i>Microbacterium oxydans</i>	
	<i>L. grayi</i>		
	<i>L. innocua</i>		
<i>Lactobacillus</i>			
<i>L. acetotolerans</i>	<i>L. cateniformis</i>	<i>L. mali</i>	<i>L. parakefiri</i>
<i>L. acidifarinae</i>	<i>L. ceti</i>	<i>L. manihotivorans</i>	<i>L. paralimentarius</i>
<i>L. acidipiscis</i>	<i>L. coleohominis</i>	<i>L. mindensis</i>	<i>L. paraplanctarum</i>
<i>L. acidophilus</i>	<i>L. collinoides</i>	<i>L. mucosae</i>	<i>L. pentosus</i>
<i>Lactobacillus agilis</i>	<i>L. composti</i>	<i>L. murinus</i>	<i>L. perolens</i>
<i>L. aligidus</i>	<i>L. concavus</i>	<i>L. nagelii</i>	<i>L. plantarum</i>
<i>L. alimentarius</i>	<i>L. coryniformis</i>	<i>L. namurensis</i>	<i>L. pontis</i>
<i>L. amylolyticus</i>	<i>L. crispatus</i>	<i>L. nantensis</i>	<i>L. protectus</i>
<i>L. amylophilus</i>	<i>L. crustorum</i>	<i>L. oligofermentans</i>	<i>L. psittaci</i>
<i>L. amylotrophicus</i>	<i>L. curvatus</i>	<i>L. oris</i>	<i>L. rennini</i>
<i>L. amylovorus</i>	<i>L. delbrueckii</i> subsp.	<i>L. panis</i>	<i>L. reuteri</i>
<i>L. animalis</i>	<i>bulgaricus</i>	<i>L. pantheris</i>	<i>L. rhamnosus</i>
<i>L. antri</i>	<i>L. delbrueckii</i> subsp.	<i>L. parabrevis</i>	<i>L. rimae</i>
<i>L. apodemi</i>	<i>delbrueckii</i>	<i>L. parabuchneri</i>	<i>L. rogosae</i>
<i>L. aviarius</i>	<i>L. delbrueckii</i> subsp. <i>lactis</i>	<i>L. paracasei</i>	<i>L. rossiae</i>
<i>L. bifementans</i>	<i>L. dextrinicus</i>	<i>L. paracollinoides</i>	<i>L. ruminis</i>
<i>L. brevis</i>	<i>L. diolivorans</i>	<i>L. parafarraginis</i>	<i>L. saerimneri</i>
<i>L. buchneri</i>	<i>L. equi</i>	<i>L. homohiochii</i>	<i>L. jensenii</i>
<i>L. camelliae</i>	<i>L. equigenerosi</i>	<i>L. iners</i>	<i>L. johnsonii</i>
<i>L. casei</i>	<i>L. farraginis</i>	<i>L. ingluviei</i>	<i>L. kalixensis</i>
<i>L. kitasatonis</i>	<i>L. farciminis</i>	<i>L. intestinalis</i>	<i>L. kefranofermentans</i>
<i>L. kunkeei</i>	<i>L. fermentum</i>	<i>L. fuchuensis</i>	<i>L. kefiri</i>
<i>L. leichmannii</i>	<i>L. fornicalis</i>	<i>L. gallinarum</i>	<i>L. kimchii</i>
<i>L. lindneri</i>	<i>L. fructivorans</i>	<i>L. gasserii</i>	<i>L. helveticus</i>
<i>L. malefermentans</i>	<i>L. frumenti</i>		<i>L. hilgardii</i>
<i>Legionella</i>			
<i>Legionella adelaidensis</i>	<i>Legionella drancourtii</i>	<i>Candidatus Legionella jeonii</i>	<i>Legionella quinlivanii</i>
<i>Legionella anisa</i>	<i>Legionella dresdenensis</i>	<i>Legionella jordanis</i>	<i>Legionella rowbothamii</i>
<i>Legionella beliardensis</i>	<i>Legionella drozanskii</i>	<i>Legionella lansingensis</i>	<i>Legionella rubrilucens</i>
<i>Legionella birminghamensis</i>	<i>Legionella dumoffii</i>	<i>Legionella londiniensis</i>	<i>Legionella sainthelensii</i>
<i>Legionella bozemaniae</i>	<i>Legionella erythra</i>	<i>Legionella longbeachae</i>	<i>Legionella sancticrucis</i>
<i>Legionella brunensis</i>	<i>Legionella fairfieldensis</i>	<i>Legionella lytica</i>	<i>Legionella shakespearei</i>
<i>Legionella busanensis</i>	<i>Legionella fallonii</i>	<i>Legionella maceachernii</i>	<i>Legionella spiritensis</i>
<i>Legionella cardiaca</i>	<i>Legionella feeleii</i>	<i>Legionella massiliensis</i>	<i>Legionella steelei</i>
<i>Legionella cherrii</i>	<i>Legionella geestiana</i>	<i>Legionella micdadei</i>	<i>Legionella steigerwaltii</i>
<i>Legionella cincinmatensis</i>	<i>Legionella genomospecies</i>	<i>Legionella monrovia</i>	<i>Legionella taurinensis</i>
<i>Legionella clemsonensis</i>	<i>Legionella gormanii</i>	<i>Legionella moravica</i>	<i>Legionella tucsonensis</i>
<i>Legionella donaldsonii</i>	<i>Legionella gratiana</i>	<i>Legionella nagasakiensis</i>	<i>Legionella tumisensis</i>
	<i>Legionella gresilensis</i>	<i>Legionella nautarum</i>	<i>Legionella wadsworthii</i>
	<i>Legionella hackeliae</i>	<i>Legionella norrlandica</i>	<i>Legionella waltersii</i>
	<i>Legionella impletisoli</i>	<i>Legionella oakridgensis</i>	<i>Legionella worsteiensis</i>
	<i>Legionella israelensis</i>	<i>Legionella parisiensis</i>	<i>Legionella yabuuchiae</i>
	<i>Legionella jamestownensis</i>	<i>Legionella pittsburghensis</i>	
		<i>Legionella pneumophila</i>	
		<i>Legionella quateirensis</i>	
<i>Oceanibulbus</i>	<i>Paemibacillus</i>	<i>Prevotella</i>	<i>Quadrifera</i>
<i>Oceanibulbus indolifex</i>	<i>Paemibacillus</i>	<i>Prevotella albensis</i>	<i>Quadrifera granulorum</i>
	<i>thiaminolyticus</i>		
<i>Oceanicaulis</i>	<i>Pantoea</i>	<i>Prevotella amnii</i>	<i>Quatronicoccus</i>
<i>Oceanicaulis alexandrii</i>	<i>Pantoea agglomerans</i>	<i>Prevotella bergensis</i>	<i>Quatronicoccus</i>
<i>Oceanicola</i>	<i>Paracoccus</i>	<i>Prevotella bivia</i>	<i>australiensis</i>
<i>Oceanicola batsensis</i>	<i>Paracoccus alcaliphilus</i>	<i>Prevotella brevis</i>	<i>Quinella</i>
<i>Oceanicola granulosis</i>	<i>Paucimonas</i>	<i>Prevotella bryantii</i>	<i>Quinella ovalis</i>
<i>Oceanicola nanhaiensis</i>	<i>Paucimonas lemoignei</i>	<i>Prevotella buccae</i>	<i>Ralstonia</i>
<i>Oceanimonas</i>	<i>Pectobacterium</i>	<i>Prevotella buccalis</i>	<i>Ralstonia eutropha</i>
<i>Oceanimonas baumannii</i>	<i>Pectobacterium aroidearum</i>	<i>Prevotella copri</i>	<i>Ralstonia insidiosa</i>

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>Oceaniserpentilla</i>	<i>Pectobacterium atrosepticum</i>	<i>Prevotella dentalis</i>	<i>Ralstonia mammitolytica</i>	
<i>Oceaniserpentilla haliotis</i>	<i>Pectobacterium betavascularum</i>	<i>Prevotella denticola</i>	<i>Ralstonia pickettii</i>	
<i>Oceanisphaera</i>	<i>Pectobacterium cacticida</i>	<i>Prevotella disiens</i>	<i>Ralstonia pseudosolanacearum</i>	
<i>Oceanisphaera donghaensis</i>	<i>Pectobacterium carnegieana</i>	<i>Prevotella histicola</i>	<i>Ralstonia syzygii</i>	
<i>Oceanisphaera litoralis</i>	<i>Pectobacterium carotovorum</i>	<i>Prevotella intermedia</i>		
<i>Oceanithermus</i>	<i>Pectobacterium chrysanthemi</i>	<i>Prevotella maculosa</i>	<i>Ralstonia solanacearum</i>	
<i>Oceanithermus desulfurans</i>	<i>Pectobacterium cypripedii</i>	<i>Prevotella marshii</i>	<i>Ramlibacter</i>	
<i>Oceanithermus profundus</i>	<i>Pectobacterium rhapontici</i>	<i>Prevotella melaninogenica</i>	<i>Ramlibacter henchirensis</i>	
<i>Oceanobacillus</i>	<i>Pectobacterium rhabdovirus</i>	<i>Prevotella micans</i>	<i>Ramlibacter tataouinensis</i>	
<i>Oceanobacillus caeni</i>	<i>Pectobacterium wasabiae</i>	<i>Prevotella multififormis</i>	<i>Raoultella</i>	
<i>Oceanospirillum</i>	<i>Planococcus</i>	<i>Prevotella nigrescens</i>	<i>Raoultella ornithinolytica</i>	
<i>Oceanospirillum linum</i>	<i>Planococcus citreus</i>	<i>Prevotella oralis</i>	<i>Raoultella planticola</i>	
	<i>Planomicrobium</i>	<i>Prevotella oris</i>	<i>Raoultella terrigena</i>	
	<i>Planomicrobium okeanokoites</i>	<i>Prevotella oulorum</i>	<i>Rathayibacter</i>	
	<i>Plesiomonas</i>	<i>Prevotella pallens</i>	<i>Rathayibacter caricis</i>	
	<i>Plesiomonas shigelloides</i>	<i>Prevotella salivae</i>	<i>Rathayibacter festucae</i>	
	<i>Proteus</i>	<i>Prevotella stercorea</i>	<i>Rathayibacter iranicus</i>	
	<i>Proteus vulgaris</i>	<i>Prevotella tanneriae</i>	<i>Rathayibacter rathayi</i>	
		<i>Prevotella timonensis</i>	<i>Rathayibacter toxicus</i>	
		<i>Prevotella veroralis</i>	<i>Rathayibacter tritici</i>	
		<i>Providencia</i>	<i>Rhodobacter</i>	
		<i>Providencia stuartii</i>	<i>Rhodobacter sphaeroides</i>	
		<i>Pseudomonas</i>	<i>Ruegeria</i>	
		<i>Pseudomonas aeruginosa</i>	<i>Ruegeria gelatinovorans</i>	
		<i>Pseudomonas alcaligenes</i>		
		<i>Pseudomonas anguilliseptica</i>		
		<i>Pseudomonas fluorescens</i>		
		<i>Pseudoalteromonas haloplanktis</i>		
		<i>Pseudomonas mendocina</i>		
		<i>Pseudomonas pseudoalcaligenes</i>		
		<i>Pseudomonas putida</i>		
		<i>Pseudomonas tutzeri</i>		
		<i>Pseudomonas syringae</i>		
		<i>Psychrobacter</i>		
		<i>Psychrobacter faecalis</i>		
		<i>Psychrobacter phenylpyruvicus</i>		
<i>Saccharococcus</i>	<i>Sagittula</i>	<i>Sanguibacter</i>	<i>Stenotrophomonas</i>	<i>Tatlockia</i>
<i>Saccharococcus thermophilus</i>	<i>Sagittula stellata</i>	<i>Sanguibacter keddieii</i>	<i>Stenotrophomonas maltophilia</i>	<i>Tatlockia maceachernii</i>
<i>Saccharomonospora</i>	<i>Salagentibacter</i>	<i>Sanguibacter suarezii</i>	<i>Streptococcus</i>	<i>Tatlockia micdadei</i>
<i>Saccharomonospora azurea</i>	<i>Salagentibacter salegens</i>	<i>Saprosira</i>	[also see below]	<i>Tenacibaculum amylolyticum</i>
<i>Saccharomonospora cyanea</i>	<i>Salimicrobium</i>	<i>Saprosira grandis</i>	<i>Streptomyces</i>	<i>Tenacibaculum discolor</i>
<i>Saccharomonospora viridis</i>	<i>Salimicrobium album</i>	<i>Sarcina</i>	<i>Streptomyces achromogenes</i>	<i>Tenacibaculum gallaicum</i>
<i>Saccharophagus</i>	<i>Salinibacter</i>	<i>Sarcina maxima</i>	<i>Streptomyces cesalbus</i>	<i>Tenacibaculum lutimaris</i>
<i>Saccharophagus degradans</i>	<i>Salinibacter ruber</i>	<i>Sarcina ventriculi</i>	<i>Streptomyces cescaepitosus</i>	<i>Tenacibaculum mesophilum</i>
<i>Saccharopolyspora</i>	<i>Salinicoccus</i>	<i>Sebaldella</i>	<i>Streptomyces cesexfoliatus</i>	<i>Tenacibaculum</i>
<i>Saccharopolyspora erythraea</i>	<i>Salinicoccus alkaliphilus</i>	<i>Sebaldella termitidis</i>	<i>Streptomyces cesexfoliatus</i>	
<i>Saccharopolyspora gregorii</i>	<i>Salinicoccus hispanicus</i>	<i>Serratia</i>	<i>Streptomyces fimbriatus</i>	
<i>Saccharopolyspora hirsuta</i>	<i>Salinicoccus roseus</i>	<i>Serratia fonticola</i>	<i>Streptomyces fradiae</i>	
<i>Saccharopolyspora hordei</i>	<i>Salinivibrio</i>	<i>Serratia marcescens</i>		
<i>Saccharopolyspora rectivirgula</i>	<i>Salinispora arenicola</i>	<i>Sphaerotilus</i>		
<i>Saccharopolyspora spinosa</i>	<i>Salinispora tropica</i>	<i>Sphaerotilus natans</i>	<i>Streptomyces fulvissimus</i>	<i>skagerrakense</i>
<i>Saccharopolyspora taberi</i>	<i>Salinivibrio</i>	<i>Sphingobacterium</i>	<i>Streptomyces griseoruber</i>	<i>Tepidanaerobacter</i>
<i>Saccharothrix</i>	<i>Salinivibrio costicola</i>	<i>Sphingobacterium multivorum</i>	<i>Streptomyces griseus</i>	<i>Tepidanaerobacter</i>
<i>Saccharothrix australiensis</i>	<i>Salmonella</i>	<i>Staphylococcus</i>	<i>Streptomyces lavendulae</i>	<i>syntrophicus</i>
<i>Saccharothrix coeruleofusca</i>	<i>Salmonella bongori</i>	[see below]	<i>Streptomyces phaeochromogenes</i>	<i>Tepidibacter</i>
<i>Saccharothrix espanaensis</i>	<i>Salmonella enterica</i>		<i>Streptomyces formicigenes</i>	<i>Tepidibacter formicigenes</i>
<i>Saccharothrix longispora</i>	<i>Salmonella subterranea</i>		<i>Streptomyces thermodiastaticus</i>	<i>Tepidibacter thalassicus</i>
<i>Saccharothrix mutabilis</i>	<i>Salmonella typhi</i>		<i>Streptomyces tubercidicus</i>	<i>Thermus</i>
<i>Saccharothrix syringae</i>				<i>Thermus aquaticus</i>
<i>Saccharothrix tangerinus</i>				<i>Thermus filiformis</i>
<i>Saccharothrix texasensis</i>				<i>Thermus thermophilus</i>
<i>Staphylococcus</i>				
<i>S. arlettae</i>	<i>S. equorum</i>	<i>S. microti</i>	<i>S. schleiferi</i>	
<i>S. agnetis</i>	<i>S. felis</i>	<i>S. muscae</i>	<i>S. sciuri</i>	

TABLE 1-continued

Example Bacteria				
Optionally, the bacteria are selected from this Table.				
<i>S. aureus</i>	<i>S. fleurettii</i>	<i>S. nepalensis</i>	<i>S. simiae</i>	
<i>S. auricularis</i>	<i>S. gallinarum</i>	<i>S. pasteurii</i>	<i>S. simulans</i>	
<i>S. capitis</i>	<i>S. haemolyticus</i>	<i>S. petrasii</i>	<i>S. stepanovicii</i>	
<i>S. caprae</i>	<i>S. hominis</i>	<i>S. pettenkoferi</i>	<i>S. succinus</i>	
<i>S. carnosus</i>	<i>S. hyicus</i>	<i>S. piscifermentans</i>	<i>S. vitulinus</i>	
<i>S. caseolyticus</i>	<i>S. intermedius</i>	<i>S. pseudointermedius</i>	<i>S. warneri</i>	
<i>S. chromogenes</i>	<i>S. kloosii</i>	<i>S. pseudolugdunensis</i>	<i>S. xylosus</i>	
<i>S. cohnii</i>	<i>S. leei</i>	<i>S. pulvereri</i>		
<i>S. condimenti</i>	<i>S. lentus</i>	<i>S. rostri</i>		
<i>S. delphini</i>	<i>S. lugdunensis</i>	<i>S. saccharolyticus</i>		
<i>S. devriesei</i>	<i>S. lutrae</i>	<i>S. saprophyticus</i>		
<i>S. epidermidis</i>	<i>S. lyticans</i>			
	<i>S. massiliensis</i>			
<i>Streptococcus</i>				
<i>Streptococcus agalactiae</i>	<i>Streptococcus infantarius</i>	<i>Streptococcus orisratti</i>	<i>Streptococcus thermophilus</i>	
<i>Streptococcus anginosus</i>	<i>Streptococcus iniae</i>	<i>Streptococcus parasanguinis</i>	<i>Streptococcus sanguinis</i>	
<i>Streptococcus bovis</i>	<i>Streptococcus intermedius</i>	<i>Streptococcus peroris</i>	<i>Streptococcus sobrinus</i>	
<i>Streptococcus canis</i>	<i>Streptococcus lactarius</i>	<i>Streptococcus pneumoniae</i>	<i>Streptococcus suis</i>	
<i>Streptococcus constellatus</i>	<i>Streptococcus milleri</i>	<i>Streptococcus pseudopneumoniae</i>	<i>Streptococcus uberis</i>	
<i>Streptococcus downei</i>	<i>Streptococcus mitis</i>	<i>Streptococcus pyogenes</i>	<i>Streptococcus vestibularis</i>	
<i>Streptococcus dysgalactiae</i>	<i>Streptococcus mutans</i>	<i>Streptococcus rattii</i>	<i>Streptococcus viridans</i>	
<i>Streptococcus equines</i>	<i>Streptococcus oralis</i>	<i>Streptococcus salivarius</i>	<i>Streptococcus zooepidemicus</i>	
<i>Streptococcus faecalis</i>	<i>Streptococcus tigurinus</i>			
<i>Streptococcus ferus</i>				
<i>Uliginosibacterium</i>	<i>Vagococcus</i>	<i>Vibrio</i>	<i>Virgibacillus</i>	<i>Xanthobacter</i>
<i>Uliginosibacterium gangwonense</i>	<i>Vagococcus carniphilus</i>	<i>Vibrio aerogenes</i>	<i>Virgibacillus halodenitrificans</i>	<i>Xanthobacter aminoxidans</i>
<i>Ulvibacter</i>	<i>Vagococcus elongatus</i>	<i>Vibrio aestuarianus</i>	<i>Virgibacillus pantothenicus</i>	<i>Xanthobacter autotrophicus</i>
<i>Ulvibacter litoralis</i>	<i>Vagococcus fessus</i>	<i>Vibrio albensis</i>	<i>Weissella</i>	<i>Xanthobacter flavus</i>
<i>Umezawaea</i>	<i>Vagococcus fluvialis</i>	<i>Vibrio alginolyticus</i>	<i>Weissella cibaria</i>	<i>Xanthobacter tagetidis</i>
<i>Umezawaea tangerina</i>	<i>Vagococcus lutrae</i>	<i>Vibrio campbellii</i>	<i>Weissella hellenica</i>	<i>Xanthobacter viscosus</i>
<i>Undibacterium</i>	<i>Vagococcus salmoninarum</i>	<i>Vibrio cholerae</i>	<i>Weissella koreensis</i>	<i>Xanthomonas alfalfae</i>
<i>Undibacterium pigrum</i>	<i>Variovorax</i>	<i>Vibrio cincinnatiensis</i>	<i>Weissella minor</i>	<i>Xanthomonas axonopodis</i>
<i>Ureaplasma</i>	<i>Variovorax boronicummulans</i>	<i>Vibrio coralliilyticus</i>	<i>Weissella thailandensis</i>	<i>Xanthomonas campestris</i>
<i>Ureaplasma urealyticum</i>	<i>Variovorax dokdonensis</i>	<i>Vibrio cyclitrophicus</i>	<i>Weissella viridescens</i>	<i>Xanthomonas citri</i>
<i>Ureibacillus</i>	<i>Variovorax paradoxus</i>	<i>Vibrio diazotrophicus</i>	<i>Williamsia</i>	<i>Xanthomonas codiaei</i>
<i>Ureibacillus composti</i>	<i>Variovorax soli</i>	<i>Vibrio fluvialis</i>	<i>Williamsia maris</i>	<i>Xanthomonas cucurbitae</i>
<i>Ureibacillus suwonensis</i>	<i>Veillonella</i>	<i>Vibrio furnissii</i>	<i>Williamsia serinedens</i>	
<i>Ureibacillus terrenus</i>	<i>Veillonella atypica</i>	<i>Vibrio gazogenes</i>	<i>Winogradskyella</i>	
<i>Ureibacillus thermophilus</i>	<i>Veillonella caviae</i>	<i>Vibrio halotocoli</i>	<i>Winogradskyella thalassocola</i>	
<i>Ureibacillus thermosphaericus</i>	<i>Veillonella criceti</i>	<i>Vibrio harveyi</i>		
	<i>Veillonella dispar</i>	<i>Vibrio ichthyenteri</i>		
	<i>Veillonella montpellierensis</i>	<i>Vibrio mediterranei</i>		
	<i>Veillonella parvula</i>	<i>Vibrio metschnikovii</i>		
	<i>Veillonella ratti</i>	<i>Vibrio mytili</i>		
	<i>Veillonella rodentium</i>	<i>Vibrio natriegens</i>		
	<i>Venenivibrio</i>	<i>Vibrio navarrensis</i>		
	<i>Venenivibrio stagnispumantis</i>	<i>Vibrio nereis</i>		
	<i>Verminephrobacter</i>	<i>Vibrio nigrripulchritudo</i>		
	<i>Verminephrobacter eiseniae</i>	<i>Vibrio ordalii</i>		
	<i>Verrucomicrobium</i>			
	<i>Verrucomicrobium spinosum</i>	<i>Vibrio orientalis</i>	<i>Wolbachia</i>	<i>Xanthomonas fragariae</i>
		<i>Vibrio parahaemolyticus</i>	<i>Wolbachia persica</i>	<i>Xanthomonas fuscans</i>
		<i>Vibrio pectenida</i>	<i>Wolinella</i>	<i>Xanthomonas gardneri</i>
		<i>Vibrio penaeicida</i>	<i>Wolinella succinogenes</i>	<i>Xanthomonas hortorum</i>
		<i>Vibrio proteolyticus</i>	<i>Zobellia</i>	<i>Xanthomonas hyacinthi</i>
		<i>Vibrio shilonii</i>	<i>Zobellia galactanivorans</i>	<i>Xanthomonas perforans</i>
		<i>Vibrio splendidus</i>	<i>Zobellia uliginosa</i>	<i>Xanthomonas phaseoli</i>
		<i>Vibrio tubiashii</i>	<i>Zoogloea</i>	<i>Xanthomonas pisi</i>
		<i>Vibrio vulnificus</i>	<i>Zoogloea ramigera</i>	<i>Xanthomonas populi</i>
			<i>Zoogloea resiniphila</i>	<i>Xanthomonas theicola</i>
				<i>Xanthomonas translucens</i>
				<i>Xanthomonas vesicatoria</i>
				<i>Xylella</i>
				<i>Xylella fastidiosa</i>
				<i>Xylophilus</i>
				<i>Xylophilus ampelinus</i>
<i>Xenophilus</i>	<i>Yangia</i>	<i>Yersinia mollaretti</i>	<i>Zooshikella</i>	<i>Zobellia</i>
<i>Xenophilus azovorans</i>	<i>Yangia pacifica</i>	<i>Yersinia philomiragia</i>	<i>Zooshikella ganghwensis</i>	<i>Zobellia denitrificans</i>
<i>Xenorhabdus</i>	<i>Yaniella</i>	<i>Yersinia pestis</i>	<i>Zunongwangia</i>	<i>Zobellia taiwanensis</i>
<i>Xenorhabdus beddingii</i>	<i>Yaniella flava</i>	<i>Yersinia pseudotuberculosis</i>	<i>Zunongwangia profumada</i>	<i>Zeaxanthinibacter</i>
<i>Xenorhabdus bovienii</i>	<i>Yaniella halotolerans</i>	<i>Yersinia rohdei</i>	<i>Zymbobacter</i>	<i>Zeaxanthinibacter</i>

TABLE 1-continued

Example Bacteria
Optionally, the bacteria are selected from this Table.

<i>Xenorhabdus cabanillasii</i>	<i>Yeosuana</i>	<i>Yersinia ruckeri</i>	<i>Zymobacter palmae</i>	<i>enoshimensis</i>
<i>Xenorhabdus doucetiae</i>	<i>Yeosuana aromativorans</i>	<i>Yokenella</i>	<i>Zymomonas</i>	<i>Zhihengliuella</i>
<i>Xenorhabdus griffithiae</i>	<i>Yersinia</i>	<i>Yokenella regensburgei</i>	<i>Zymomonas mobilis</i>	<i>Zhihengliuella</i>
<i>Xenorhabdus hominickii</i>	<i>Yersinia aldovae</i>	<i>Yonghaparkia</i>	<i>Zymophilus</i>	<i>halotolerans</i>
<i>Xenorhabdus koppenhoeferi</i>	<i>Yersinia bercovieri</i>	<i>Yonghaparkia alkaliphila</i>	<i>Zymophilus paucivorans</i>	<i>Xylanibacterium</i>
<i>Xenorhabdus nematophila</i>	<i>Yersinia enterocolitica</i>	<i>Zavarzinia</i>	<i>Zymophilus raffinosivorans</i>	<i>Xylanibacterium ulmi</i>
<i>Xenorhabdus poinarii</i>	<i>Yersinia entomophaga</i>	<i>Zavarzinia compransoris</i>		
<i>Xylanibacter</i>	<i>Yersinia frederiksenii</i>			
<i>Xylanibacter oryzae</i>	<i>Yersinia intermedia</i>			
	<i>Yersinia kristensenii</i>			

TABLE 2

Systemic inflammatory response syndrome	
Finding	Value
Temperature	<36° C. (96.8° F.) or >38° C. (100.4° F.)
Heart rate	>90/min
Respiratory rate	>20/min or PaCO ₂ < 32 mmHg (4.3 kPa)
WBC	<4 × 10 ⁹ /L (<4000/mm ³), >12 × 10 ⁹ /L (>12,000/mm ³), or 10% bands

TABLE 3

Sequences of genomic targets to selectively CRISPR-kill *E. coli* strains. The 3-bp PAM are shown in bold type (ie, tgg, agg and cgg). Expected SpCas9-gRNA cleavage sites are indicated by vertical arrows (3 bp upstream of the PAM).

Target strain	Target gene	Genomic target sequence
<i>E. coli</i> (EHEC) ATCC43888	23S ribosomal RNA	taggtgaagtc ccctcgcgga tgg ↓ (SEQ ID NO: 1)
<i>E. coli</i> Nissle 1917	yapH	caccgcattac ctttatgcagg ↓ (SEQ ID NO: 2)
<i>E. coli</i> Nissle 1917	pks	gcgcggtgtg gttgcctcgg ↓ (SEQ ID NO: 3)

TABLE 4

Sequences of genomic targets & Cas to selectively CRISPR-kill *C. difficile*

(a) Overview of design spacers in pMTL84151-cdCRISPR1 targeting *C. difficile*

Gene	Name	PAM	target
tcdA	TA2	CCA	ACACCTTAACCCAGCCATAGAGTCTGATAA TAACCTC (SEQ ID NO: 4)
tcdB	TB2	CCA	GACTTATTTGAGTCTATAGAGAAACCTAGT TCAGTAA (SEQ ID NO: 5)
rRNA-1	TRR1	CCT	TCGACGACTTCTTCCAAAAGGTTAGATAAT CGGCTC (SEQ ID NO: 6)
rRNA-2	TRR2	CCA	GTACAGGATGGACCCGCGTCTGATTAGCTA GTTGGTA (SEQ ID NO: 7)

15

TABLE 4-continued

Sequences of genomic targets & Cas to selectively CRISPR-kill <i>C. difficile</i>				
gyrA	TAR1	CCA	TCCTCATGGAGATACTGCTGTTATTATGCTATGGTA (SEQ ID NO: 8)	20
(b) Cas3 sequence (SEQ ID NO: 9)				
Sequence				
TCAAATAAAATTTGGTCTATTTTCATTACTTATAAGCACACCTTTACCAAATAT				25
CTGTTTTGTATGCTTATTTTCGTATATATCATATTTATATAAAAGTATTTT				
TAAATCTTCAAGACCTTTTCACCTGTATGTCAATTACATTTTGTTTAGCTTT				
ATATATTTGGTAAATTTACAGTTTTTTTAAATATCTCTCTCTGCTTTTTT				30
TCTCTTAGACTTTTATTTTATTTATCAATTGTCTATCATCATTACTATATGC				
ATTTATAAGCTCTTGTCCCAACTCTTCATAATCTTTGATTAAAGTTTCTTC				
AATTTTATTATATATATCCTCTGGAATTACTGTATATCCTTCAATATTTCT				35
AAGTATATTTTGTGCATCTTTAGAACCTAAACTATATGGCCTTATAGTATC				
TAAATATTTAAAGCACTGGTAAATCTCTTTTCAAAGCTGTACCTTCTAG				
ACTTCTTTTAGAGTATAGTATCTTAACCATTTCTACCTTATATTTTCTTT				40
CATCTTCACACATTTCTTTCCATTTATAAAAGTTTTTCAGTAGCTCAAGTCC				
TTTTTCTACAATATCCTTATCATAAATACTACCTATTCTGTAGCTTTTTT				
TGTATATATAAATATATTTGGGCTATTTTCTTCACTACGACTTCTATA				45
ACATCTACCAAATCGTTGAAATAGACTGTCAAGTGTGAATTTTCTGTATG				
AAGCTCATCAAATCAATATCAAGGGATGCTTCCACTAATTGTGTAGTAAT				
CCAAATACCATTACTATCACTATCTGCAAATCTTTTATATATTTTCTAA				50
TTTTGCTCTATCTTCTTGTATATACATAGAATGTAAAAGATTTAAGTTGAC				
ATCTATCTCTTTGGCTTAACATTTCTTCTATTAACCTCATATTTTCTAC				
AGCACTTTTAACTGTATTTACTATTACAAGAACTTTTTTATTCACTCCACT				55
TTGTATTATTTTACCTAAGTTTTTCATCTATTGAATTTTCTACAATAGACAC				
ACAATGTCTTATTTTTCTGTATTACATGTCAATTCAGCTAAGTTACTATT				
CATTACACCTCTTTTTTTAATTCATCTATATATATGGTTGGCATAGTAGC				60
TGTCATTATCATAAATCTGCCACCTATCTTATGTATCATTTCTATACCTTT				
TACCAATACAGCTGTATTTCTGGTGAATATGCTTGTATCTCATCTATTAC				
TACTTTTGAATATGCTAATGTTGAGTACACTTTTTTACACCTCTATACAA				65
AAAAGGAAATTTAAATATTTGGTCTATTGTAGAAATGTCAAGTTTGTCAAGA				

TABLE 4-continued

Sequences of genomic targets & Cas to selectively CRISPR-kill *C. difficile*

TAATAACTTTGCTAAATCTACAATCTCACTTGAATTTTCTTGATTACTTTC
 TTCTAGATAATCTATTGTGTGAATGTAACAAACCTAAGAATGTATCATT
 TGATTCTCCAACCTCCAACCTATATTTTTGCTCTATCAAATAATGCATTTAT
 ACTTACTCTTAATGGAAGTGTAAAAATGCCTTGTCTTTATCTATCCTAAAT
 TAAGGCAGTTTCTGTTTTTCCATTCTCGTAGGTGCAATCAGTATTATATT
 CTTATTTCTATTAGATTAGCAAATGATTGAGCCTCTCTCAAACCTACCAAA
 CTCTTTTCATTAATAATTTTCTGTCTGTCTCTATATTTATAACATTATT
 GCATTCCACAACCTTATGAGCAGAAGCACTGTGGTCTAATCTATGTAGTAT
 TCCTTTTAAACATAATATAAATATAGTACTTGTGATTTTTATCTATTCT
 TTTTTCTACACTTTGTAGATATACTTTACTCAATTTTTCTGTTTTTATTGG

TABLE 4-continued

Sequences of genomic targets & Cas to selectively CRISPR-kill *C. difficile*

5 ATATCTAACTTTAAATTCATGCTGTAGTTCATAAACTTTATTTATTAATC
 TTCATCTAATATTTTTTGTATTAAATTTTTAAAACTTTATCTATAAAGAT
 ATCTCTTTTCATGATGATATACAATAAAGTGTATTAATACTGCTCTAAGTTC
 10 TTTATTTTTTTTCTGTCTATATAACTATAATCAATAAATGCAGGAGAAAG
 AATATTATGGCTACATTATTTCTAAATGAGTTACTATTTTAGGTTTACT
 TATTTTAGTATCCATTCTGCTTTTTTATTAACCTTGAAACGGTGAAAATGC
 15 TTTTCCAATATCATGAAATTCATAACAAAGTCAAGTAATTGCCAAAATAT
 CTCTCTTTCTAGAAAATCTAAGCTATTTATATTTTTTCCATAACTTTCTCT
 TAATACATTCATTTGTTTTAAAAGTTCATCAGTATGTTCTCTAAGTGTTTC
 20 CACTGGATTAGATTTAGCATATAACAT

TABLE 4

Sequences of Cas9 used to selectively kill *E. coli*.

SEQ ID NO: 10 (Cas9 nucleotide sequence)
 ATGGATAAGAAATACTCAATAGGCTTAGATATCGGCACAAATAGCGTCCGATGGGCGGT
 GATCACTGATGAATATAAGTTCCGCTCAAAAAGTTCAAGGTTCTGGGAAATACAGACC
 GCCACAGTATCAAAAAAATCTTATAGGGGCTCTTTATTTGACAGTGGAGAGACAGCG
 GAAGCGACTCGTCTCAAACGGACAGCTCGTAGAAGGTATACACGTCGGAAGAATCGTAT
 TTGTTATCTACAGGAGATTTTTCAAAATGAGATGGCGAAAGTATAGATAGTTTCTTTCA
 TCGACTTGAAGAGTCTTTTTTGGTGAAGAAGACAGAAGCATGAACCTTATCTCATT
 TGGAAATATAGTAGATGAAGTTGCTTATCATGAGAATATCCAACATCTATCATCTGCG
 AAAAAATTTGGTAGATTTCTACTGATAAAGCGGATTTGCGCTTAATCTATTTGGCCTTAGC
 GCATATGATTAAGTTTCGTGGTCAATTTTTGATTGAGGGAGATTTAAATCCTGATAATAGT
 GATGTGGACAAACTATTTATCCAGTTGGTACAAAACCTACAATCAATTTTGAAGAAAAAC
 CCTATTAAACGCAAGTGGAGTAGATGCTAAAGCGATTTCTTTCGACGATGAGTAAATCA
 AGACGATTAGAAAATCTCATTGCTCAGCTCCCGGTGAGAAGAAAATGGCTTATTGG
 GAATCTCATTGCTTTGTCTTTGGTTTGGACCCCTAATTTTAAATCAAAATTTGATTGGCA
 GAAGATGCTAAATACAGCTTTCAAAGATACCTACGATGATGATTAGATAATTTATTG
 CGCAAAATGGAGATCAATATGCTGATTTGTTTTTGGCAGCTAAGAAATTCATCTGAGT
 ATTTTACTTTTCAGATATCCTAAGAGTAAATCTGAATAACTAAGGCTCCCTATCAGCT
 TCAATGATTAACGCTACGATGAACATCATCAAGACTTGACTCTTTAAAAGCTTTAGTT
 CGACAACAACCTCCAGAAAAGTATAAAGAAATCTTTTTTGTCAATCAAAAACCGGATA
 TGCAGGTTATATTGATGGGGAGCTAGCCAAGAAGAAATTTATAAAATTTATCAAACCAAT
 TTTAGAAAAAATGGATGGTACTGAGGAATTTATGGTGAACATAAATCGTGAAGATTTGCT
 CGCAAGCAACGGACCTTTGACAACCGCTCTATTCCCATCAAAATTCATTTGGGTGAGCT
 GCATGCTATTTTGAAGAACAAGAAGACTTTTATCCATTTTTAAAAGCAATCGTGAGAA
 GATTGAAAAATCTGACTTTTCGAATTCCTTATATGTTGGTCCATTGGCCGTTGGCAAT
 AGTCGTTTTGCAATGGATGACTCGGAAGTCTGAAGAACAATTAACCGTCAAGATTTTGA
 GAAGTTGTCGATAAAGGTGCTTCAGCTCAATCATTATTGAAACGCATGACAAACTTTGAT
 AAAAACTTCCAAATGAAAAGTACTACCAAAACATAGTTTGTCTTATGAGTATTTTACG
 GTTTATAACGAATTGACAAAAGTCAAAATGTTACTGAAGGAATGCGAAAAACGAGCATT
 TCTTTCAAGTGAACAGAAGAAAGCCATTGTGATTTACTCTTCAAAAACAATCGAAAAGT
 AACCGTTAAGCAATTAAGAAGATTTTCAAAAAAATAGAATGTTTGTATAGTGTGTA
 AATTCAGGAGTTGAAGATAGATTTAATGCTTCATTAGGTACCTACCATGATTTGCTAAA
 AATTATTAAGATAAAGATTTTTGGATAATGAAGAAAAATGAAGATATCTTAGAGGATA
 TTGTTTTAACAATGACCTTATTTGAAGATAGGAGATGATTGAGGAAGACTTAAAACAT
 ATGCTCACCTCTTTGATGATAAAGGTGATGAAACAGCTTAAACGTCGCGTTATACTGGTT
 GGGACGTTTGTCTCGAAAATGATTAATGGTATTAGGGATAAGCAATCTGGCAAAAACA
 ATATTAGATTTTTTGAATCAGATGTTTTCGCAATCGCAATTTTATGACGCTGATCCATG
 ATGATAGTTTGAATTAAGAAGACATTTCAAAAAGCACAAGTGTCTGGACAAGGCGAT
 AGTTTACATGAAACATATGCAAATTTAGCTGGTAGCCCTGCTATTAATAAAGGATTTTTA
 CAGACTGTAAAAGTGTGATGAATGGTCAAAGTAAATGGGGCGGCATAAGCCAGAAAA
 TATCGTTATTGAAATGGCAGTGAATCAGACAACCTCAAAGGGCCAGAAAAATTCG
 GAGAGCGTATGAACGAATCGAAGAAGGTATCAAGAATTAGGAAGTCAAGATTTTAAA
 GAGCATCCTGTTGAAAATACTCAATGCAAAAATGAAAAGCTCTATCTTATATCTCCAA
 AATGGAAGAGACATGATGTTGACCAAGAATTAGATATTAATCGTTTAAAGTATGATGAT
 GTCGATCAGATTTGTTCCACAAGTTTCCTTAAAGACGATTTCAATGACAAATAAGGTTA
 ACGCGTTCTGATAAAAAATCGTGGTAAATCGGATAACGTTCCAAGTGAAGAAGTAGTCAA
 AAAGATGAAAACTATTGGAGACAACCTTCAAACGC CAAGTAACTCAACGTAAGT
 TTGATAATTTAACGAAAGCTGAACGTTGAGGTTGAGTGAACCTGATAAAGCTGGTTTTA
 TCAAACGCCAATTTGAACTCGCCAAATCACTAAGCATGTGGCAACAATTTGGATA
 GTCGATGAATACTAAATACGATGAAAATGATAAATTTATTCGAGAGGTTAAAGTGAAT

TABLE 4-continued

Sequences of Cas9 used to selectively kill *E. coli*.

ACCTTAAATCTAAATAGTTTCTGACTCCGAAAAGATTTCCAATTCATAAAGTACGT
 GAGATTAACAATTACCATCATGCCCATGATGCGTATCTAAATGCCGTCGTTGGAAGTCT
 TTGATTAAGAAATATCCAAAACCTTGAATCGGAGTTTGTCTATGGTGATTATAAGTTTAT
 GATGTTTCGTAATAATGATGCTAAGTCTGAGCAAGAAATAGGCAAGCAACCCGAAAATA
 TTTCTTTTACTCTAATATCATGAACCTTCTTCAAAACAGAAATTACACTTGCAATGGAGA
 GATTTCGCAACGCCCTCTAATCGAAACTAATGGGGAAACTGGAGAAATGTCTGGGATA
 AAGGGCGAGATTTTCCACAGTGCAGCAAGTATTGTCCATGCCCAAGTCAATATTGTCA
 AGAAAACAGAAGTACAGACAGCGGATTTCCCAAGGAGTCAATTTTACAAAAGAAAT
 TCGGCAAGCTTATTGCTCGTAAAAAGACTGGGATCCAAAATAATATGGTGGTTTGTAT
 AGTCCACCGGTAGCTTATTAGTCTTAGTGGTGTCTAAGGTGAAAAAGGGAAATCGAA
 GAAGTTAAAATCCGTTAAAGAGTTACTAGGGATCACAATTATGGAAAGAAGTTCTTTG
 AAAAAATCCGATTGACTTTTAGAAGCTAAAGGATATAAGGAAGTTAAAAAGACTTA
 ATCATTAAACTACCTAAATATAGTCTTTTGGAGTAGAAAAAGGTCGTAACCGGATGCTG
 GCTAGTGCCTGAGAAATTACAAAAGGAAATGAGCTGGCTCTGCCAAGCAAAATATGTGAA
 TTTTTTATATTTAGTCTAGTCAATATGAAAAGTTGAAAGGTAGTCCAGAAGATAACGAACA
 AAAACAATTTGTTGAGGAGCAGCATAAGCATTATTTAGATGAGATTATTGAGCAAAATCAG
 TGAAATTTCTAAGCGTGTATTTTAGCAGATGCCAATTAGATAAAGTTCTTAGTGCATAT
 AACAAACATAGAGCAAACCAATACGTGAACAAGCAGAAATATTATTCATTTATTAC
 GTTGACGAATCTTGGAGCTCCCGCTGCTTTTAAATATTTGATACAACAATTTGATCGTAA
 ACGATATACGCTTACAAAAGAGTTTTAGATGCCACTCTTATCCATCAATCCTACTGG
 TCTTTATGAAACACGCATTGATTTGAGTCTAGGAGGTGACTGA

SEQ ID NO: 11 (Cas9 amino acid sequence)
 MDKKYSIGLDIGTNSVGVAVI TDEYKVPKSKFKVLGNTDRHSIKKNIIGALLFDSGETAEAT
 RLKRTARRRYTRRNRI CYLQEI FSNEMAKVDDSPFFHRLLEESFLVEEDKKHERHPIFGNIVDE
 VAYHEKYPTI YHLRKKLVSDTKADLRLLIYLALAHMIKFRGHFLIEGLDLPNDSDVDKLFIQ
 VQTYNQLPEENP INASGVDAKAIL SARLSKSRRL ENLIAQLPGEKKNGLFGNLI ALSGLTPNF
 KSNPDLAEDAKLQSKD TYDDDLNLLAQI GDQYADLFLAAKNLSDAILLSDI LRVNTEITKA
 PLSASMIKRYDEHHQDL TLLKALVRQQLPEKYKE IFFDQSKNGYAGYIDGGASQEEFYKFKP
 ILEKMDGTEELLVKLNREDDL RKQRTPFDNGSI PHQIHLGELHAILRRQEDFYPLKDNREKIEK
 ILTFRIPYVYVGLARGNSRFAMTRKSEETITPWNFEEVVDKGSASQSFIERMTNFDKNLPNE
 KVLPHKSLLYEYFTYVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQLKE
 DYFKKIECFDSVEISGVEDRNASLGTYHDLKI IKDKDFLDNEENEDILEDIVLTLTLFEDRE
 MIEERLKYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTI LDPLKSDGFANRN
 FMQLIHDDSLTFKEDIQKAQVSGQDLSLHEHIANLAGSPAIKKGI LQTVKVVDELVKVMGRH
 KPEIVIMEMARENQTTQKQKNSRERMKRI EEGI KELGSQILKEHPVENTQLQNEKLYLYLQ
 NGRDMYVDQELINRLSDYVDHIVPQSFLKDDSDNKNVLRSDKNRGSNDNVPSEEVVKK
 MKNYWRQLLNAKLI TQRKFDNLTKAERGGSELDKAGFI KRQLVETRQITKHVAQILD SRM
 NTKYDENDKLI REVKVITLKS KLVSDFRKDPQFYKVR EINNYYHHAHDAYLNAVVG TALIKKY
 PKLESEFVYGDYKVDVRKMI AKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIET
 NGETGEIVWDKGRDFATVRKVL SMPQVNI VVKTEVQTGGFSKESILPKRNSDKLIARKKDW
 DPKKYGGPDSPTVAVSVLVVAKVEKGSKLLKSVKELLGITIMERSSFEKNPIDFLKAGYKE
 VKKDLI IKLPKYSLEENGRKRLASAGELQKGNELALPSKYVNFYLASHYEKLGKSPED
 NEQQLFVEQHKHYLDEIIEQISEFSKRVI LADANLDKVL SAYNKHDKPIREQAENI IHLFTL
 TNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD

TABLE 5

Publication	Gram-neg. bacteria as causative infectious pathogen in cancer patients	<i>E. coli/K. pneumoniae/ P. aeruginosa</i> as causative pathogens in bacteraemia in cancer patients [fraction of Gram-neg. cases]	
Samonis et al	65%	54% [85%]	45
Velasco et al	45%	33% [74%]	

TABLE 5-continued

Publication	Gram-neg. bacteria as causative infectious pathogen in cancer patients	<i>E. coli/K. pneumoniae/ P. aeruginosa</i> as causative pathogens in bacteraemia in cancer patients [fraction of Gram-neg. cases]
Marin et al	55%	51% [92%]
Anatoliotaki et al	47%	34% [73%]

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11
 <210> SEQ ID NO 1
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct
 <400> SEQUENCE: 1
 taggtgaagt ccctcgcgga tgg

-continued

<210> SEQ ID NO 2
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 2

 caccgcgatt acctttatgc agg 23

<210> SEQ ID NO 3
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 3

 gcgcgggtgt ggttgctt cgg 23

<210> SEQ ID NO 4
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 4

 acaccttaac ccagccatag agtctgataa taacttc 37

<210> SEQ ID NO 5
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 5

 gacttatttg agtctataga gaaacctagt tcagtaa 37

<210> SEQ ID NO 6
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 6

 tcgacgactt cttccaaaag gttagataat eggcttc 37

<210> SEQ ID NO 7
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

 <400> SEQUENCE: 7

 gtacaggatg gaccgcgctc tgattagcta gttggta 37

<210> SEQ ID NO 8
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 8

tccctcatgga gatactgctg tttattatgc tatggta 37

<210> SEQ ID NO 9

<211> LENGTH: 2322

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 9

tcaataaat tggctatatt cattaacttat aagcacacct ttaccaaata tctgttttgt 60

atgcttattt tcgtatatat catatttata taaaagtatt tttaaatcct caagaccttt 120

cacctgtatg tcaattacat tttgttttagc tttatatatt ggtaaattta cagttttttt 180

aattatctct cttcttgctt tttttctctt agactttatt ttattttatca attgtctatc 240

atcattacta tatgcattta taagctcttg tcccaactct tcataatcct tgattaaagt 300

ttcttcaatt ttattatata taccctctgg aattactgta tatccttcaa tattttetaag 360

tatattttgt gcactcttag aacctaaact atatggcggt atagtatccta aaatatttaa 420

agcactggta aatctctttt caaaagctgt acctctctaga ctttcttttag agtatagtat 480

cttaaccatt tctaccttat atttttcttt catcttcaca cattcttttc cattttataaa 540

agttttcagt agctcaagtc cttttttctac aatatoctta tcataaatac tacctatttc 600

tgtagctttt tctgtatata taaatatatt tgggctatatt tcttcatct cagcacttct 660

ataacatccta ccaaatcggt gaaatagact gtcaagtgtt gaattttctg tatgaagctc 720

atcaaaatca atatcaaggg atgcttccac taattgtgta gtaatccaaa taccattact 780

atcactatct gcaaattcct ttatatattt ttctaatttt gctctatcct cttgtatata 840

catagaatgt aaaagattta agttgacatc tattcctttt ggcttaacta tttctctat 900

taactcatat tttttctacag cacttttaac tgtatttact attacaagaa cttttttatt 960

cattccactt tgtattattt tacctaagtt ttcactcatt gaattttcta caatagacac 1020

acaatgtcct atttttctg tattacatgt caattcagct aagttactat tcattacacc 1080

tctttttttt aattcatccta tatatatggg tggcatagta gctgtcatta tcataaatct 1140

gccacctatc ttatgatca tttctatacc ttttccaat acagctgcta tttctgggta 1200

atatgcttgt atctcatccta ttactacttt tgaatatgct aatgttgagt acactttttc 1260

atacctccta tacaaaaaag gaaatttaaa tatttggctct attgtagaaa atgtcagttt 1320

gcaagataat aactttgcta aatctacaat ctcaactgaa ttttcttgat tactttcttc 1380

tagataatct attgctgttg aatgtaacaa acctaagaat gtatcatttg attctccaac 1440

tccaactata ttttttgctc tatcaaaata tgcatttata cttactctta atggaagtgt 1500

aaaaaatgoc ttgtctttat ctatccaaat taaggcagtt tctgtttttc ccattcctgt 1560

aggtgcaatc agtattatat tcttatttct attagattta gcaaatgatt gagcctctct 1620

caactacca aactctttca ttaaataaatt ttctgtctgt tctcctatat ttataacatt 1680

attgcattcc acaacttcat gagcagaagc actgtggctc aatctatgta gtattccttt 1740

taacataata tataaattat agtacttctg atttttatct attctttttt ctacactttg 1800

tagatatact ttactcaatt tttctgtttt tattggatat ctaactttaa attcatgctg 1860

tagttcataa actttattta ttaaactctc atcctaattt ttttgattta aatttttaaa 1920

-continued

atctttatct ataaagatat ctctttcatg atgatataca ataacttgat ttaatactgc	1980
tctaagttct ttattttttt tctgtctat ataactataa tcaataaatg caggagaaag	2040
ataattatgg cctacattat tttctaaatg agttactatt ttaggttcat ttatttttagt	2100
atccattctg ctttttatta actcttgaaa cggtgaaaa gcttttccaa tatcatgaaa	2160
ttctataaca aagtcaagta attgccaaaa tatctctct tctagaaaat ctaagctatt	2220
tatatttttt ccataacttt ctcttaatac attcatttgt tttaaaagtt catcagtatg	2280
ttctctaagt gtttccactg gattagattt agcatataac at	2322

<210> SEQ ID NO 10
 <211> LENGTH: 4107
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 10

atggataaga aatactcaat aggccttagat atcggcacia atagcgtcgg atgggcggtg	60
atcactgatg aatataaggt tccgtctaaa aagttcaagg ttctgggaaa tacagaccgc	120
cacagtatca aaaaaaatct tataggggct cttttatttg acagtggaga gacagcggaa	180
gogactcgtc tcaaacggac agctcgtaga aggtatacac gtcggaagaa tcgtatttgt	240
tatctacagg agattttttc aaatgagatg gcgaaagtag atgatagttt ctttcatcga	300
cttgaagagt cttttttggt ggaagaagac aagaagcatg aacgtcatcc tatttttggg	360
aatatagtag atgaagttgc ttatcatgag aaatatccaa ctatctatca tctgcgaaaa	420
aaattggtag attctactga taaagcggat ttgcgcttaa tctatttggc cttagcgc	480
atgattaagt ttcgtgggtc ttttttgatt gagggagatt taaatcctga taatagtgat	540
gtggacaaac tatttatcca gttgttacia acctacaatc aattatttga agaaaaccct	600
attaacgcaa gtggagtaga tgctaaagcg attctttctg cactgattgag taaatcaaga	660
cgattagaaa atctcattgc tcagctcccc ggtgagaaga aaaatggctt atttgggaat	720
ctcattgctt tgtcattggg tttgaccctc aattttaaat caaattttga tttggcagaa	780
gatgctaaat tacagctttc aaaagatact tacgatgatg atttagataa tttattggcg	840
caaattggag atcaaatatgc tgatttgtt ttggcagcta agaatttacc agatgctatt	900
ttactttcag atatcctaag agtaataact gaaataacta aggetcccc atcagcttca	960
atgattaaac gctacgatga acatcatcaa gacttgactc ttttaaaagc tttagttcga	1020
caacaacttc cagaaaagta taaagaaatc ttttttgatc aatcaaaaa cggatatgca	1080
ggttatatgg atgggggagc tagccaagaa gaattttata aatttatcaa accaatttta	1140
gaaaaaatgg atggtactga ggaattattg gtgaaactaa atcgtgaaga tttgctgcgc	1200
aagcaacgga cctttgcaa cggctctatt ccccatcaaa ttcacttggg tgagctgcat	1260
gctattttga gaagacaaga agacttttat ccatttttaa aagacaatcg tgagaagatt	1320
gaaaaaatct tgacttttcg aattccttat tatgttggtc cattggcgcg tggcaatagt	1380
cgttttgcat ggatgactcg gaagtctgaa gaaacaatta ccccatggaa ttttgaagaa	1440
gttgcgata aaggtgcttc agctcaatca tttattgaac gcatgacaaa ctttgataaa	1500
aatcttccaa atgaaaaagt actacaaaa catagtttgc tttatgagta ttttacgggt	1560
tataacgaat tgacaaaagg caaatatgtt actgaaggaa tgcgaaaacc agcatttctt	1620

-continued

tcaggtgaac	agaagaaagc	cattgttgat	ttactcttca	aaacaaatcg	aaaagtaacc	1680
gttaagcaat	taaagaaga	ttatttcaaa	aaaatagaat	gttttgatag	tggtgaaatt	1740
tcaggagttg	aagatagatt	taatgcttca	ttaggtacct	accatgattt	gctaaaaatt	1800
attaagata	aagatttttt	ggataatgaa	gaaaatgaag	atatcttaga	ggatattggt	1860
ttaacattga	ccttatttga	agatagggag	atgattgagg	aaagacttaa	aacatattgct	1920
cacctctttg	atgataaggt	gatgaaacag	cttaaacgtc	gccgttatac	tggttgggga	1980
cgtttgcttc	gaaaattgat	taatggtatt	agggataagc	aatctggcaa	aacaatatta	2040
gattttttga	aatcagatgg	ttttgccaat	cgcaatttta	tgcagctgat	cctatgatgat	2100
agtttgacat	ttaagaaga	cattcaaaaa	gcacaagtgt	ctggacaagg	cgatagttta	2160
catgaacata	ttgcaaattt	agctggtagc	cctgctatta	aaaaaggtat	ttacagact	2220
gtaaaagttg	ttgatgaatt	ggtcaaagta	atggggcggc	ataagccaga	aaatattcgtt	2280
attgaaatgg	cacgtgaaaa	tcagacaact	caaaagggcc	agaaaaattc	gcgagagcgt	2340
atgaaacgaa	tcgaagaagg	tatcaaagaa	ttaggaagtc	agattcttaa	agagcatcct	2400
gttgaataa	ctcaattgca	aaatgaaaag	ctctatctct	attatctcca	aaatggaaga	2460
gacatgtatg	tggaccaaga	attagatatt	aatcgtttaa	gtgattatga	tgtcgatcac	2520
attgttccac	aaagtttctt	taaagacgat	tcaatagaca	ataaggtctt	aacgcgttct	2580
gataaaaaatc	gtggtaaatc	ggataacggt	ccaagtgaag	aagtagtcaa	aaagatgaaa	2640
aactattgga	gacaacttct	aaacgccaaag	ttaatcactc	aacgtaagtt	tgataattta	2700
acgaaagctg	aacgtggagg	tttgagttaa	cttgataaag	ctggttttat	caaacgcca	2760
ttggttga	ctcgccaat	cactaagcat	gtggcacaaa	ttttggatag	tcgcatgaa	2820
actaaatacg	atgaaaatga	taaacttatt	cgagaggtta	aagtgattac	cttaaatct	2880
aaattagttt	ctgacttccg	aaaagatttc	caattctata	aagtacgtga	gattaacaat	2940
taccatcatg	cccattgatg	gtatctaaat	gccgtcgttg	gaactgcttt	gattaagaaa	3000
tatccaaaac	ttgaatcgga	gtttgtctat	ggtgattata	aagtttatga	tgttcgtaaa	3060
atgattgcta	agtctgagca	agaaataggc	aaagcaaccg	caaaatattt	cttttactct	3120
aatatcatga	acttcttcaa	aacagaaatt	acacttgcaa	atggagagat	tcgcaaacgc	3180
cctctaactg	aaactaatgg	ggaaactgga	gaaattgtct	gggataaagg	gcgagatttt	3240
gccacagtgc	gcaaagtatt	gtccatgcc	caagtcaata	ttgtcaagaa	aacagaagta	3300
cagacaggcg	gattctccaa	ggagtcaatt	ttaccaaaaa	gaaattcggga	caagcttatt	3360
gctcgtaaaa	aagactggga	tccaaaaaaa	tatggtgggt	ttgatagtcc	aacggtagct	3420
tattcagttc	tagtggttgc	taaggtgaa	aaagggaaat	cgaagaagtt	aaaatccgtt	3480
aaagagttac	tagggatcac	aattatggaa	agaagttcct	ttgaaaaaaa	tccgattgac	3540
tttttagaag	ctaaaggata	taagggaagt	aaaaaagact	taatcattaa	actacctaaa	3600
tatagtcttt	ttgagttaga	aaacggtcgt	aaacggatgc	tggctagtgc	cggagaatta	3660
caaaaaggaa	atgagctggc	tctgccaaagc	aaatatgtga	attttttata	tttagctagt	3720
cattatgaaa	agttgaaggg	tagtccagaa	gataacgaac	aaaaacaatt	gtttgtggag	3780
cagcataagc	attatttaga	tgagattatt	gagcaaatca	gtgaattttc	taagcgtggt	3840
attttagcag	atgccaat	agataaagtt	cttagtgcat	ataacaaca	tagagacaaa	3900
ccaatacgtg	aacaagcaga	aaatattatt	catttattta	cgttgacgaa	tcttgagct	3960
cccgtgctt	ttaaatattt	tgatacaaca	atgatcgta	aacgatatac	gtctacaaaa	4020

-continued

gaagtttttag atgccactct tatccatcaa tccatcactg gtctttatga aacacgcatt 4080

gatttgagtc agctaggagg tgactga 4107

<210> SEQ ID NO 11

<211> LENGTH: 1368

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 11

Met Asp Lys Lys Tyr Ser Ile Gly Leu Asp Ile Gly Thr Asn Ser Val
1 5 10 15

Gly Trp Ala Val Ile Thr Asp Glu Tyr Lys Val Pro Ser Lys Lys Phe
20 25 30

Lys Val Leu Gly Asn Thr Asp Arg His Ser Ile Lys Lys Asn Leu Ile
35 40 45

Gly Ala Leu Leu Phe Asp Ser Gly Glu Thr Ala Glu Ala Thr Arg Leu
50 55 60

Lys Arg Thr Ala Arg Arg Arg Tyr Thr Arg Arg Lys Asn Arg Ile Cys
65 70 75 80

Tyr Leu Gln Glu Ile Phe Ser Asn Glu Met Ala Lys Val Asp Asp Ser
85 90 95

Phe Phe His Arg Leu Glu Glu Ser Phe Leu Val Glu Glu Asp Lys Lys
100 105 110

His Glu Arg His Pro Ile Phe Gly Asn Ile Val Asp Glu Val Ala Tyr
115 120 125

His Glu Lys Tyr Pro Thr Ile Tyr His Leu Arg Lys Lys Leu Val Asp
130 135 140

Ser Thr Asp Lys Ala Asp Leu Arg Leu Ile Tyr Leu Ala Leu Ala His
145 150 155 160

Met Ile Lys Phe Arg Gly His Phe Leu Ile Glu Gly Asp Leu Asn Pro
165 170 175

Asp Asn Ser Asp Val Asp Lys Leu Phe Ile Gln Leu Val Gln Thr Tyr
180 185 190

Asn Gln Leu Phe Glu Glu Asn Pro Ile Asn Ala Ser Gly Val Asp Ala
195 200 205

Lys Ala Ile Leu Ser Ala Arg Leu Ser Lys Ser Arg Arg Leu Glu Asn
210 215 220

Leu Ile Ala Gln Leu Pro Gly Glu Lys Lys Asn Gly Leu Phe Gly Asn
225 230 235 240

Leu Ile Ala Leu Ser Leu Gly Leu Thr Pro Asn Phe Lys Ser Asn Phe
245 250 255

Asp Leu Ala Glu Asp Ala Lys Leu Gln Leu Ser Lys Asp Thr Tyr Asp
260 265 270

Asp Asp Leu Asp Asn Leu Leu Ala Gln Ile Gly Asp Gln Tyr Ala Asp
275 280 285

Leu Phe Leu Ala Ala Lys Asn Leu Ser Asp Ala Ile Leu Leu Ser Asp
290 295 300

Ile Leu Arg Val Asn Thr Glu Ile Thr Lys Ala Pro Leu Ser Ala Ser
305 310 315 320

Met Ile Lys Arg Tyr Asp Glu His His Gln Asp Leu Thr Leu Leu Lys
325 330 335

Ala Leu Val Arg Gln Gln Leu Pro Glu Lys Tyr Lys Glu Ile Phe Phe

-continued

Thr Thr Gln Lys Gly Gln Lys Asn Ser Arg Glu Arg Met Lys Arg Ile
 770 775 780

Glu Glu Gly Ile Lys Glu Leu Gly Ser Gln Ile Leu Lys Glu His Pro
 785 790 795 800

Val Glu Asn Thr Gln Leu Gln Asn Glu Lys Leu Tyr Leu Tyr Tyr Leu
 805 810 815

Gln Asn Gly Arg Asp Met Tyr Val Asp Gln Glu Leu Asp Ile Asn Arg
 820 825 830

Leu Ser Asp Tyr Asp Val Asp His Ile Val Pro Gln Ser Phe Leu Lys
 835 840 845

Asp Asp Ser Ile Asp Asn Lys Val Leu Thr Arg Ser Asp Lys Asn Arg
 850 855 860

Gly Lys Ser Asp Asn Val Pro Ser Glu Glu Val Val Lys Lys Met Lys
 865 870 875 880

Asn Tyr Trp Arg Gln Leu Leu Asn Ala Lys Leu Ile Thr Gln Arg Lys
 885 890 895

Phe Asp Asn Leu Thr Lys Ala Glu Arg Gly Gly Leu Ser Glu Leu Asp
 900 905 910

Lys Ala Gly Phe Ile Lys Arg Gln Leu Val Glu Thr Arg Gln Ile Thr
 915 920 925

Lys His Val Ala Gln Ile Leu Asp Ser Arg Met Asn Thr Lys Tyr Asp
 930 935 940

Glu Asn Asp Lys Leu Ile Arg Glu Val Lys Val Ile Thr Leu Lys Ser
 945 950 955 960

Lys Leu Val Ser Asp Phe Arg Lys Asp Phe Gln Phe Tyr Lys Val Arg
 965 970 975

Glu Ile Asn Asn Tyr His His Ala His Asp Ala Tyr Leu Asn Ala Val
 980 985 990

Val Gly Thr Ala Leu Ile Lys Lys Tyr Pro Lys Leu Glu Ser Glu Phe
 995 1000 1005

Val Tyr Gly Asp Tyr Lys Val Tyr Asp Val Arg Lys Met Ile Ala Lys
 1010 1015 1020

Ser Glu Gln Glu Ile Gly Lys Ala Thr Ala Lys Tyr Phe Phe Tyr Ser
 1025 1030 1035 1040

Asn Ile Met Asn Phe Phe Lys Thr Glu Ile Thr Leu Ala Asn Gly Glu
 1045 1050 1055

Ile Arg Lys Arg Pro Leu Ile Glu Thr Asn Gly Glu Thr Gly Glu Ile
 1060 1065 1070

Val Trp Asp Lys Gly Arg Asp Phe Ala Thr Val Arg Lys Val Leu Ser
 1075 1080 1085

Met Pro Gln Val Asn Ile Val Lys Lys Thr Glu Val Gln Thr Gly Gly
 1090 1095 1100

Phe Ser Lys Glu Ser Ile Leu Pro Lys Arg Asn Ser Asp Lys Leu Ile
 1105 1110 1115 1120

Ala Arg Lys Lys Asp Trp Asp Pro Lys Lys Tyr Gly Gly Phe Asp Ser
 1125 1130 1135

Pro Thr Val Ala Tyr Ser Val Leu Val Val Ala Lys Val Glu Lys Gly
 1140 1145 1150

Lys Ser Lys Lys Leu Lys Ser Val Lys Glu Leu Leu Gly Ile Thr Ile
 1155 1160 1165

Met Glu Arg Ser Ser Phe Glu Lys Asn Pro Ile Asp Phe Leu Glu Ala
 1170 1175 1180

-continued

Lys Gly Tyr Lys Glu Val Lys Lys Asp Leu Ile Ile Lys Leu Pro Lys
 1185 1190 1195 1200

Tyr Ser Leu Phe Glu Leu Glu Asn Gly Arg Lys Arg Met Leu Ala Ser
 1205 1210 1215

Ala Gly Glu Leu Gln Lys Gly Asn Glu Leu Ala Leu Pro Ser Lys Tyr
 1220 1225 1230

Val Asn Phe Leu Tyr Leu Ala Ser His Tyr Glu Lys Leu Lys Gly Ser
 1235 1240 1245

Pro Glu Asp Asn Glu Gln Lys Gln Leu Phe Val Glu Gln His Lys His
 1250 1255 1260

Tyr Leu Asp Glu Ile Ile Glu Gln Ile Ser Glu Phe Ser Lys Arg Val
 1265 1270 1275 1280

Ile Leu Ala Asp Ala Asn Leu Asp Lys Val Leu Ser Ala Tyr Asn Lys
 1285 1290 1295

His Arg Asp Lys Pro Ile Arg Glu Gln Ala Glu Asn Ile Ile His Leu
 1300 1305 1310

Phe Thr Leu Thr Asn Leu Gly Ala Pro Ala Ala Phe Lys Tyr Phe Asp
 1315 1320 1325

Thr Thr Ile Asp Arg Lys Arg Tyr Thr Ser Thr Lys Glu Val Leu Asp
 1330 1335 1340

Ala Thr Leu Ile His Gln Ser Ile Thr Gly Leu Tyr Glu Thr Arg Ile
 1345 1350 1355 1360

Asp Leu Ser Gln Leu Gly Gly Asp
 1365

The invention claimed is:

1. A method of treating or reducing an infection that is associated with septicemia and is caused by first bacteria in a subject, the method comprising selectively killing the first bacteria comprised by the subject by cutting a target site comprised by the genomes of the first bacteria, wherein the cutting is carried out using a Cas nuclease that is programmed by a guide RNA to cut the target site, wherein the first bacteria are Enterobacteriaceae bacteria selected from the group consisting of *Escherichia*, *Salmonella*, *Klebsiella*, *Shigella*, *Enterobacter* and *Citrobacter*, wherein the subject comprises second bacteria of one or more strains or species that are different from that of the first bacteria, wherein the genomes of the second bacteria do not comprise the target site, wherein the genomes of the second bacteria are not cut by the Cas nuclease in the subject, whereby second bacteria survive in the presence of the Cas nuclease in the subject.
2. The method of claim 1, wherein the infection is a urinary tract infection, a lung infection, or a kidney infection.
3. The method of claim 1, wherein the infection is a urinary tract infection.
4. The method of claim 3, wherein the first bacteria comprise *Escherichia coli* cells.
5. The method of claim 4, wherein the first bacteria are enterohemorrhagic *E. coli* (EHEC).
6. The method of claim 1, wherein the infection is a lung infection.
7. The method of claim 6, wherein the first bacteria comprise *E. coli* or *Klebsiella* cells.
8. The method of claim 6, wherein the first bacteria are *Klebsiella pneumoniae* cells.
9. The method of claim 1, wherein and the subject has a compromised immune system.
10. The method of claim 9, wherein the subject is a cancer patient, a transplant patient, or a patient suffering from a viral infection.
11. The method of claim 9, wherein the subject is a liver, kidney or lung transplant patient.
12. The method of claim 9, wherein the subject is on an immunosuppressant medication.
13. The method of claim 9, wherein the subject has undergone surgery.
14. The method of claim 9, wherein the subject is suffering from a chronic disease.
15. The method of claim 1, wherein the first bacteria are comprised by a gut, kidney or lung microbiota of the subject.
16. The method of claim 1, wherein the subject is a human.
17. The method of claim 16, wherein at the start of the treatment, the subject has a temperature of <math><36^{\circ}\text{C}</math>. or >math>>38^{\circ}\text{C}</math>.; a heart rate of >math>>90</math>/min, a respiratory rate of >math>>20</math> breaths/min or PaCO₂<math><4.3</math> kPa; and white blood cell count of <math><4000</math>/mm³ or >math>>12,000</math>/mm³.
18. The method of claim 1, wherein the Cas nuclease is a Cas3.
19. The method of claim 1, wherein the Cas nuclease is a Cas9.
20. The method of claim 1, wherein the method comprises reducing the infection at least 100-fold or at least 1000-fold by the first 30 minutes of the treatment.
21. The method of claim 1, wherein the method comprises maintaining reduction of the infection by at least 100-fold for at least 60 minutes after exposing the subject to the Cas nuclease and/or the guide RNA.
22. The method of claim 20, wherein the reduction of the infection persists for at least 30 minutes immediately after the first 30 minutes of the treatment.

23. The method of claim 1, wherein the infection is reduced by at least 90% for 1 hour or more.

24. The method of claim 1, wherein the infection is durably treated.

25. The method of claim 1, wherein the method comprises administering the Cas nuclease and/or the guide RNA to the subject at a first time (T1) and at a second time (T2), wherein T2 is at least 1 hour after T1.

26. The method of claim 1, wherein the method comprises administering to the subject a nucleic acid vector comprising the guide RNA or a DNA encoding the guide RNA.

27. The method of claim 26, comprising administering a second nucleic acid vector to the subject, wherein the second vector encodes the Cas nuclease.

28. The method of claim 26, wherein the Cas nuclease is an endogenous Cas nuclease of the first bacteria.

* * * * *