


United
States
of
America



To Promote the Progress

of Science and Useful Arts

The Director

of the United States Patent and Trademark Office has received an application for a patent for a new and useful invention. The title and description of the invention are enclosed. The requirements of law have been complied with, and it has been determined that a patent on the invention shall be granted under the law.

Therefore, this United States

Patent

grants to the person(s) having title to this patent the right to exclude others from making, using, offering for sale, or selling the invention throughout the United States of America or importing the invention into the United States of America, and if the invention is a process, of the right to exclude others from using, offering for sale or selling throughout the United States of America, products made by that process, for the term set forth in 35 U.S.C. 154(a)(2) or (c)(1), subject to the payment of maintenance fees as provided by 35 U.S.C. 41(b). See the Maintenance Fee Notice on the inside of the cover.

Katherine Kelly Vidal

DIRECTOR OF THE UNITED STATES PATENT AND TRADEMARK OFFICE

Maintenance Fee Notice

If the application for this patent was filed on or after December 12, 1980, maintenance fees are due three years and six months, seven years and six months, and eleven years and six months after the date of this grant, or within a grace period of six months thereafter upon payment of a surcharge as provided by law. The amount, number and timing of the maintenance fees required may be changed by law or regulation. Unless payment of the applicable maintenance fee is received in the United States Patent and Trademark Office on or before the date the fee is due or within a grace period of six months thereafter, the patent will expire as of the end of such grace period.

Patent Term Notice

If the application for this patent was filed on or after June 8, 1995, the term of this patent begins on the date on which this patent issues and ends twenty years from the filing date of the application or, if the application contains a specific reference to an earlier filed application or applications under 35 U.S.C. 120, 121, 365(c), or 386(c), twenty years from the filing date of the earliest such application ("the twenty-year term"), subject to the payment of maintenance fees as provided by 35 U.S.C. 41(b), and any extension as provided by 35 U.S.C. 154(b) or 156 or any disclaimer under 35 U.S.C. 253.

If this application was filed prior to June 8, 1995, the term of this patent begins on the date on which this patent issues and ends on the later of seventeen years from the date of the grant of this patent or the twenty-year term set forth above for patents resulting from applications filed on or after June 8, 1995, subject to the payment of maintenance fees as provided by 35 U.S.C. 41(b) and any extension as provided by 35 U.S.C. 156 or any disclaimer under 35 U.S.C. 253.



US011471531B2

(12) **United States Patent**
Clube et al.

(10) **Patent No.:** **US 11,471,531 B2**
(45) **Date of Patent:** ***Oct. 18, 2022**

- (54) **SELECTIVELY ALTERING MICROBIOTA FOR IMMUNE MODULATION**
- (71) Applicant: **SNIPR Technologies Limited**, London (GB)
- (72) Inventors: **Jasper Clube**, London (GB); **Morten Sommer**, London (GB); **Christian Grøndahl**, London (GB)
- (73) Assignee: **SNIPR Technologies Limited**, London (GB)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

8,241,498 B2	8/2012	Summer
8,252,576 B2	8/2012	Campbell
8,906,682 B2	12/2014	June
8,911,993 B2	12/2014	June
8,916,381 B1	12/2014	June
8,975,071 B1	3/2015	June
9,101,584 B2	8/2015	June
9,102,760 B2	8/2015	June
9,102,761 B2	8/2015	June
9,113,616 B2	8/2015	Stevens
9,328,156 B2	5/2016	June
9,464,140 B2	10/2016	June
9,481,728 B2	11/2016	June
9,499,629 B2	11/2016	June
9,518,123 B2	12/2016	June
9,540,445 B2	1/2017	June
9,701,964 B2	7/2017	Clube
9,758,583 B2	9/2017	Wang
9,822,372 B2	11/2017	Zhang
9,879,269 B2	1/2018	Barrangou
10,066,233 B2	9/2018	Barrangou
10,136,639 B2	11/2018	Wuest
10,136,649 B2	11/2018	Barrangou
10,195,273 B2 *	2/2019	Clube A61K 35/15

(21) Appl. No.: **17/132,624**

(22) Filed: **Dec. 23, 2020**

(65) **Prior Publication Data**
US 2021/0113689 A1 Apr. 22, 2021

Related U.S. Application Data

(63) Continuation of application No. 16/453,598, filed on Jun. 26, 2019, which is a continuation of application No. 16/192,752, filed on Nov. 15, 2018, now Pat. No. 10,363,308, which is a continuation of application No. 15/820,296, filed on Nov. 21, 2017, now Pat. No. 10,195,273, which is a continuation of application No. PCT/EP2017/063593, filed on Jun. 4, 2017.

(30) **Foreign Application Priority Data**

Jun. 5, 2016 (GB) 1609811

- (51) **Int. Cl.**
A61K 39/395 (2006.01)
A61K 39/00 (2006.01)
A61P 37/00 (2006.01)
A61P 35/00 (2006.01)
A61K 31/7105 (2006.01)
A61K 35/15 (2015.01)
A61K 35/17 (2015.01)

(52) **U.S. Cl.**
CPC *A61K 39/3955* (2013.01); *A61K 31/7105* (2013.01); *A61K 35/15* (2013.01); *A61K 35/17* (2013.01); *A61P 35/00* (2018.01); *A61P 37/00* (2018.01); *A61K 39/0011* (2013.01); *A61K 2039/505* (2013.01); *C12N 2320/30* (2013.01); *C12N 2320/50* (2013.01)

(58) **Field of Classification Search**
CPC .. *A61K 39/3955*; *A61K 39/0011*; *A61P 37/00*
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,626,504 A 12/1986 Puhler
5,633,154 A 5/1997 Schaefer

FOREIGN PATENT DOCUMENTS

CA 3010891 A1 7/2017
EP 2325332 A1 5/2011

(Continued)

OTHER PUBLICATIONS

How gut bacteria boost cancer immunotherapy; <https://microbiomepost.com/how-gut-bacteria-boost-cancer-immunotherapy/> (Year: 2020).*
Chen et al., Journal of Immunology Research 2020, vol. 2020:1-13 (Year: 2020).*
Kaiser, Science Mag, 2017; <https://www.sciencemag.org/news/2017/11/your-gut-bacteria-could-determine-how-you-respond-cutting-edge-cancer-drugs> (Year: 2017).*
Aklujkar et al. (2010) "Interference With Histidyl-tRNA Synthetase by a CRISPR Spacer Sequence as a Factor in the Evolution of Pelobacter Carbinolicus," BMC Evolutionary Biology 10:203, 15 pages.
American Lung Association (2019). "Preventing COPD," retrieved from <https://www.lung.org/lung-health-and-diseases/lung-disease-lookup/copd/symptoms-causes-risk-factors/preventing-copd.html>, last visited Aug. 5, 2019, 1 page.

(Continued)

Primary Examiner — Gary B Nickol
Assistant Examiner — Lakia J Jackson-Tongue
(74) Attorney, Agent, or Firm — Morrison & Foerster LLP

(57) **ABSTRACT**

The invention relates to methods of modulating immune cells in a patient by altering microbiota of the patient. The invention also relates to methods of modulating treatments or therapies in a subject organism by altering microbiota of the subject. The invention also relates to cell populations, systems, arrays, cells, RNA, kits and other means for effecting this. In an example, advantageously selective targeting of a particular species in a human gut microbiota using guided nucleic acid modification is carried out to effect the alteration.

16 Claims, 19 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

U.S. PATENT DOCUMENTS

10,300,138 B2	5/2019	Clube	2016/0339064 A1	11/2016	Kovarik et al.
10,300,139 B2	5/2019	Clube	2016/0345578 A1	12/2016	Barrangou
10,363,308 B2	7/2019	Clube	2016/0347836 A1	12/2016	Grosso
10,463,049 B2	11/2019	Clube	2016/0354416 A1*	12/2016	Gajewski A61P 43/00
10,506,812 B2	12/2019	Clube	2017/0022499 A1	1/2017	Lu
10,524,477 B2	1/2020	Clube	2017/0037416 A1	2/2017	Barrangou
10,596,255 B2	3/2020	Clube	2017/0106025 A1	4/2017	Kovarik
10,603,379 B2	3/2020	Clube	2017/0106026 A1	4/2017	Kovarik
10,760,065 B2	9/2020	Lu et al.	2017/0114351 A1	4/2017	Mahfouz
10,760,075 B2	9/2020	Sommer et al.	2017/0143772 A1	5/2017	Mulder
10,765,740 B2	9/2020	Clube et al.	2017/0173085 A1	6/2017	Kovarik
10,920,222 B2	2/2021	Sommer et al.	2017/0174713 A1	6/2017	Du
10,953,090 B2	3/2021	Clube et al.	2017/0175142 A1	6/2017	Zhang
11,147,830 B2	10/2021	Clube	2017/0196225 A1	7/2017	Clube
2003/0049841 A1	3/2003	Short	2017/0246221 A1	8/2017	Clube
2004/0096974 A1	5/2004	Herron	2017/0247690 A1	8/2017	Quake
2005/0118719 A1	6/2005	Schmidt	2017/0304443 A1	10/2017	Lebwohl
2009/0155768 A1	6/2009	Scholl	2017/0327582 A1	11/2017	Bissonnette
2010/0076057 A1	3/2010	Sontheimer	2017/0340733 A1	11/2017	Cao
2010/0093617 A1	4/2010	Barrangou	2018/0015131 A1*	1/2018	Gajewski A61K 45/06
2011/0002889 A1	1/2011	Barrangou	2018/0055852 A1	3/2018	Kutok
2011/0143997 A1	6/2011	Henry et al.	2018/0064114 A1	3/2018	Clube
2012/0269859 A1	10/2012	Minato	2018/0064115 A1	3/2018	Clube
2013/0011828 A1	1/2013	Barrangou	2018/0070594 A1	3/2018	Clube
2013/0109053 A1	5/2013	Macdonald	2018/0084785 A1	3/2018	Clube
2013/0121968 A1	5/2013	Quay	2018/0084786 A1	3/2018	Clube
2013/0287748 A1	10/2013	June	2018/0140698 A1	5/2018	Clube
2013/0288368 A1	10/2013	June	2018/0146681 A1	5/2018	Clube
2013/0309258 A1	11/2013	June	2018/0147221 A1*	5/2018	von Maltzahn A61P 35/04
2014/0068797 A1	3/2014	Doudna	2018/0155721 A1	6/2018	Lu
2014/0105912 A1	4/2014	Noelle	2018/0155729 A1	6/2018	Beisel
2014/0106449 A1	4/2014	June	2018/0161368 A1	6/2018	Odegard
2014/0107092 A1	4/2014	Meyerson	2018/0179547 A1	6/2018	Zhang
2014/0179726 A1	6/2014	Bajaj	2018/0200342 A1	7/2018	Bikard
2014/0199767 A1	7/2014	Barrangou	2018/0273937 A1	9/2018	Beisel et al.
2014/0234972 A1	8/2014	Zhang	2018/0273940 A1	9/2018	Sommer
2014/0341920 A1	11/2014	Noelle	2018/0303934 A1	10/2018	Clube
2014/0349400 A1	11/2014	Jakimo et al.	2018/0326057 A1	11/2018	Clube
2014/0370017 A1	12/2014	June	2018/0355378 A1	12/2018	Krom et al.
2015/0004705 A1	1/2015	Lu et al.	2018/0371405 A1	12/2018	Barrangou
2015/0031134 A1	1/2015	Zhang	2019/0015441 A1	1/2019	Shachar
2015/0032263 A1	1/2015	Keyl et al.	2019/0021343 A1	1/2019	Barrangou
2015/0050699 A1	2/2015	Siksny	2019/0070233 A1	3/2019	Yeung
2015/0050729 A1	2/2015	June	2019/0117709 A1	4/2019	Kovarik
2015/0064138 A1	3/2015	Lu	2019/0133135 A1	5/2019	Clube
2015/0093822 A1	4/2015	June	2019/0134194 A1	5/2019	Clube
2015/0099299 A1	4/2015	June	2019/0136230 A1	5/2019	Sather
2015/0118202 A1	4/2015	June	2019/0142881 A1	5/2019	Turner et al.
2015/0125463 A1	5/2015	Cogswell	2019/0160120 A1	5/2019	Haaber
2015/0132263 A1	5/2015	Liu et al.	2019/0230936 A1	8/2019	Clube
2015/0132419 A1	5/2015	Arvik	2019/0240325 A1	8/2019	Clube
2015/0139943 A1	5/2015	Campana	2019/0240326 A1	8/2019	Clube
2015/0140001 A1	5/2015	Lee	2019/0255084 A1	8/2019	Schentag
2015/0184139 A1	7/2015	Zhang	2019/0256900 A1	8/2019	Zhang
2015/0225730 A1	8/2015	Minshull et al.	2019/0298779 A1	10/2019	Falb
2015/0232881 A1	8/2015	Glucksmann	2019/0321468 A1	10/2019	Clube et al.
2015/0290244 A1	10/2015	June	2019/0321469 A1	10/2019	Clube et al.
2015/0353905 A1	12/2015	Weiss	2019/0321470 A1	10/2019	Clube
2016/0009805 A1	1/2016	Kowanetz	2019/0359933 A1	11/2019	Swee
2016/0009813 A1	1/2016	Themeli	2020/0030444 A1	1/2020	Clube
2016/0024510 A1	1/2016	Bikard	2020/0046773 A1	2/2020	Borody
2016/0040215 A1	2/2016	Henn et al.	2020/0085066 A1	3/2020	Clube
2016/0081314 A1	3/2016	Thurston	2020/0087660 A1	3/2020	Sommer
2016/0115488 A1	4/2016	Zhang	2020/0102551 A1	4/2020	Barrangou
2016/0115489 A1	4/2016	Zhang	2020/0115716 A1	4/2020	Martinez
2016/0130355 A1	5/2016	June	2020/0121787 A1	4/2020	Clube
2016/0159905 A1	6/2016	Abdiche	2020/0157237 A1	5/2020	Regev
2016/0159907 A1	6/2016	June	2020/0164070 A1	5/2020	Clube
2016/0160186 A1	6/2016	Parsley	2020/0179460 A1	6/2020	Kovarik
2016/0194404 A1	7/2016	June	2020/0205416 A1	7/2020	Clube
2016/0208012 A1	7/2016	June	2020/0282027 A1	9/2020	Bikard et al.
2016/0244784 A1	8/2016	Jacobson et al.	2020/0337313 A1	10/2020	Clube
2016/0281053 A1	9/2016	Sorek	2020/0354690 A1	11/2020	Garofolo
2016/0324938 A1	11/2016	Bikard	2020/0390886 A1*	12/2020	Clube A61K 39/0011
2016/0333348 A1	11/2016	Clube	2021/0009996 A1	1/2021	Sommer
			2021/0145006 A1	5/2021	Clube
			2021/0147827 A1	5/2021	Clube
			2021/0147857 A1	5/2021	Clube
			2021/0163960 A1	6/2021	Martinez et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2021/0189406 A1 6/2021 Martinez et al.
 2021/0198665 A1 7/2021 Sommer et al.
 2021/0230559 A1 7/2021 Clube
 2021/0283167 A1 9/2021 Clube
 2021/0290654 A1 9/2021 Clube
 2021/0386773 A1 12/2021 Clube

FOREIGN PATENT DOCUMENTS

EP 2840140 A1 2/2015
 EP 3461337 A1 4/2019
 EP 3132035 B8 4/2020
 EP 3132036 B8 4/2020
 EP 3630975 A1 4/2020
 EP 3633032 A2 4/2020
 EP 3634442 A1 4/2020
 EP 3634473 A1 4/2020
 RU 2531343 C2 10/2014
 WO 2005046579 A2 5/2005
 WO 2005046579 A3 8/2005
 WO 2007025097 A2 3/2007
 WO 2007025097 A3 7/2007
 WO 2008108989 A2 9/2008
 WO 2008108989 A3 3/2009
 WO 2010011961 A2 1/2010
 WO 2010011961 A3 6/2010
 WO 2010075424 A2 7/2010
 WO 2010075424 A3 9/2010
 WO 2012079000 A1 6/2012
 WO 2012079000 A4 8/2012
 WO 2012164565 A1 12/2012
 WO 2013063361 A1 5/2013
 WO 2013176772 A1 11/2013
 WO 2014012001 A2 1/2014
 WO 2014018423 A2 1/2014
 WO 2014018423 A3 1/2014
 WO 2014093595 A1 6/2014
 WO 2014093661 A2 6/2014
 WO 2014093661 A3 8/2014
 WO 2014124226 A1 8/2014
 WO 2014093661 A9 10/2014
 WO 2014204725 A1 12/2014
 WO 2015034872 A2 3/2015
 WO 2014012001 A3 4/2015
 WO 2015034872 A3 4/2015
 WO 2015058018 A1 4/2015
 WO 2015069682 A2 5/2015
 WO 2015070083 A1 5/2015
 WO 2015071474 A2 5/2015
 WO 2015075688 A1 5/2015
 WO 2015088643 A1 6/2015
 WO 2015089351 A1 6/2015
 WO 2015089419 A2 6/2015
 WO 2015069682 A3 7/2015
 WO 2015071474 A3 8/2015
 WO 2015089419 A3 9/2015
 WO 2015136541 A2 9/2015
 WO 2015148680 A1 10/2015
 WO 2015153940 A1 10/2015
 WO 2015155686 A2 10/2015
 WO 2015159068 A1 10/2015
 WO 2015159086 A1 10/2015
 WO 2015159087 A1 10/2015
 WO 2015136541 A3 11/2015
 WO 2016044745 A1 3/2016
 WO 2016063263 A2 4/2016
 WO 2016063263 A3 6/2016
 WO 2016196361 A1 12/2016
 WO 2016196605 A1 12/2016
 WO 2016205276 A1 12/2016
 WO 2017009399 A1 1/2017
 WO 2017042347 A1 3/2017
 WO 2017058751 A1 4/2017
 WO 2017112620 A1 6/2017
 WO 2017118598 A1 7/2017

WO 2017211753 A1 12/2017
 WO 2018064165 A2 4/2018
 WO 2018081502 A1 5/2018
 WO 2018115519 A1 6/2018
 WO 2018217351 A1 11/2018
 WO 2018217981 A1 11/2018
 WO 2018222969 A1 12/2018
 WO 2018226853 A1 12/2018
 WO 2018064165 A3 6/2019
 WO 2020072248 A1 4/2020
 WO 2020072250 A1 4/2020
 WO 2020072253 A1 4/2020
 WO 2020072254 A1 4/2020
 WO 2020152369 A1 7/2020

OTHER PUBLICATIONS

Ang, Y.L.E. et al. (2015). "Best Practice in the Treatment of Advanced Squamous Cell Lung Cancer," *Ther. Adv. Respir. Dis.* 9(5):224-235.

Anonymous (Apr. 2016). "Checkpoint Inhibition: A Promising Immunotherapeutic Approach for Colorectal Cancer," *Oncology*, 5(3):1-5, retrieved from <http://www.personalizedmedonc.com/publications/prno/april-2016-vol-5-no-3/checkpoint-inhibition-a-promising-immunotherapeutic-approach-for-colorectal-cancer-2/>, last visited Aug. 27, 2019, 5 pages.

Arnold, I.C. et al. (Apr. 8, 2015, e-pub. Mar. 4, 2015). "Helicobacter Hepaticus Infection in BALB/c Mice Abolishes Subunit-Vaccine-Induced Protection Against *M. tuberculosis*," *Vaccine* 33(15):1808-1814.

Arslan, Z. et al. (May 7, 2013). "RcsB-BglJ-Mediated Activation of Cascade Operon Does Not Induce the Maturation of CRISPR RNAs in *E. coli* K12," *RNA Biology* 10(5):708-715.

Arumugam et al. (May 12, 2011). "Enterotypes of the human gut microbiome," *Nature* 473(7346):174-180, 16 pages.

Bae, T. et al. (2006). "Prophages of *Staphylococcus aureus* Newman and Their Contribution to Virulence," *Molecular Microbiology* pp. 1-13.

Barrangou, R. et al. (Mar. 2007). "CRISPR Provides Acquired Resistance Against Viruses in Prokaryotes," *Science*, 315:1709-1712.

Beisel, C.L. et al. (2014). "A CRISPR Design for Next-Generation Antimicrobials," *Genome Biology* 15:516, 4 pages.

Belizario, J.E. et al. (Oct. 6, 2015). "Human Microbiomes and Their Roles in Dysbiosis, Common Diseases, and Novel Therapeutic Approaches," *Frontiers in Microbiology* 6(1050):1-16.

Bikard, D. et al. (2013, e-pub. Jun. 12, 2013). "Programmable Repression and Activation of Bacterial Gene Expression Using an Engineered CRISPR-Cas System," *Nucleic Acids Research* 41(15):7429-7437.

Bikard, D. et al. (2017, e-pub. Sep. 6, 2017). "Using CRISPR-Cas Systems as Antimicrobials," *Current Opinion in Microbiology* 37:155-160.

Bikard, D. et al. (Aug. 16, 2012). "CRISPR Interference Can Prevent Natural Transformation and Virulence Acquisition during In Vivo Bacterial Infection," *Cell Host & Microbe* 12(2):177-186.

Bikard, D. et al. (Nov. 2014). "Development of Sequence-Specific Antimicrobials Based on Programmable CRISPR-Cas Nucleases," *Nature Biotechnology* 32(11):1146-1151, 16 pages.

Broaders, E. et al. (Jul./Aug. 2013). "Mobile Genetic Elements of the Human Gastrointestinal Tract," *Gut Microbes* 4(4):271-280.

Brouns, S.J.J. et al. (Aug. 15, 2008). Supplemental Material for "Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes," *Science* 321:960-964.

Brouns, S.J.J. et al. (Aug. 15, 2008). "Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes," *Science* 321:960-964.

Bryksin, A. V. et al. (Oct. 8, 2010). "Rational Design of a Plasmid Origin That Replicates Efficiently in Both Gram-Positive and Gram-Negative Bacteria," *PLoS One* 5(10):e13244, 9 pages.

Bugrysheva, J.V. et al. (Jul. 2011, E-Pub. Apr. 29, 2011). "The Histone-Like Protein Hip is Essential for Growth of *Streptococcus pyogenes*: Comparison of Genetic Approaches to Study Essential Genes," *Appl. Environ. Microbiol.* 77(13):4422-4428.

(56)

References Cited

OTHER PUBLICATIONS

- Bullman, S. et al. (Nov. 23, 2017). "Analysis of Fusobacterium Persistence and Antibiotic Response in Colorectal Cancer," *Science* pp. 1443-1448, 10 pages.
- Burns, M.B. et al. (2015). "Virulence Genes are a Signature of the Microbiome in the Colorectal Tumor Microenvironment," *Genome Medicine* 7:55, 12 pages.
- Catalao, M.J. et al. (Jul. 2013, e-pub. Nov. 8, 2012). "Diversity in Bacterial Lysis Systems: Bacteriophages Show the Way," *FEMS Microbiology Reviews* 37(4):554-571.
- Chan, B.K. et al. (2013). "Phage Cocktails and the Future of Phage Therapy," *Future Microbiol.* 8(6):769-783.
- Chan, C.T.Y. et al. (Dec. 2015). "'Deadman' and 'Passcode' Microbial Kill Switches for Bacterial Containment," *Nat. Chem. Biol.* 12(2):82-86.
- Cheadle, E.J. et al. (2012). "Chimeric Antigen Receptors for T-Cell Based Therapy," *Methods Mol. Biol.* 907:645-666, 36 pages.
- Citorik, R.J. et al. (Nov. 2014, e-pub Sep. 21, 2014). "Sequence-Specific Antimicrobials Using Efficiently Delivered RNA-Guided Nucleases," *Nat. Biotechnol.* 32(11):1141-1145, 18 pages.
- Cochrane, K. et al. (2016, e-pub. Nov. 3, 2015). "Complete Genome Sequences and Analysis of the *Fusobacterium nucleatum* Subspecies *animalis* 7-1 Bacteriophage ϕ funu1 and ϕ funu2," *Anaerobe* 38:125-129. Abstract Only.
- Cong, L. et al. (Feb. 15, 2013, e-pub. Oct. 11, 2013). "Multiplex Genome Engineering Using CRISPR/Cas Systems," *Science* 339(6121):819-823, 9 pages.
- Consumer Updates (2019). "Combating Antibiotic Resistance," retrieved from <https://www.fda.gov/ForConsumers/ConsumerUpdates/ucm092810.htm>, last visited Jan. 28, 2019.
- Coyne, M.J. et al. (2014). "Evidence of Extensive DNA Transfer between *Bacteroidales* Species Within the Human Gut," *mBio* 5(3):e01305-14, 12 pages.
- Cui, L. et al. (2016, e-pub. Apr. 8, 2016). "Consequences of Cas9 Cleavage in the Chromosome of *Escherichia coli*," *Nucleic Acids Research* 44(9):4243-4251.
- Dailere, R. et al. (Oct. 18, 2016). "Enterococcus hirae and Barnesiella intestinihominis Facilitate Cyclophosphamide-Induced Therapeutic Immunomodulatory Effects," *Immunity* 95:931-943.
- Datsenko, K.A. et al. (Jul. 10, 2012). "Molecular Memory of Prior Infections Activates the CRISPR/Cas adaptive Bacterial Immunity System," *Nature Communication* 3:945, 7 pages.
- De Filippo, C. et al. (Aug. 33 2010). "Impact of Diet in Shaping Gut Microbiota Revealed by a Comparative Study in Children From Europe and Rural Africa," *Proc. Natl. Acad. Sci. USA* 107(33):14691-14696, 6 pages.
- De Paepe, M. et al. (Mar. 28, 2014). "Bacteriophages: An Underestimated Role in Human and Animal Health?" *Frontiers in Cellular and Infection Microbiology* 4(39):1-11.
- Deeks, E.D. (2014, e-pub. Jul. 15, 2014). "Nivolumab: A Review of Its Use in Patients With Malignant Melanoma," *Drugs* 74:1233-1239.
- Deghorain, M. et al. (Nov. 23, 2012). "The Staphylococci Phages Family: An Overview," *Viruses* 4:3316-3335.
- Denham, J.D. et al. (2018). "Case Report: Treatment of Enteropathogenic *Escherichia coli* Diarrhea in Cancer Patients: A Series of Three Cases," *Case Reports in Infectious Diseases* Article ID 8438701:1-3.
- Dhar, A.D. (Jul. 20, 2018). "Overview of Bacterial Skin Infections," Merck Manual retrieved from <https://www.merckmanuals.com/home/skin-disorders/bacterial-skin-infections/overview-of-bacterial-skin-infections>, last visited Jul. 20, 2018, 3 pages.
- Dickson, R.P. et al. (Jan./Feb. 2017). "Bacterial Topography of the Healthy Human Lower Respiratory Tract," *American Society for Microbiology* 8(1):e02287-6, 12 pages.
- Dierz-Villasenor, C. et al. (May 2013). "CRISPR-Spacer Integration Reporter Plasmids Reveal Distinct Genuine Acquisition Specificities Among CROSPR-Cas I-E Variants of *Escherichia coli*," *RNA Biology* 10(5):792-802.
- Dutilh, B.E. et al. (Jul. 24, 2014). "A Highly Abundant Bacteriophage Discovered in the Unknown Sequences of Human Faecal Metagenomes," *Nature Communications* 5(4498):1-10.
- Edgar et al. (Dec. 2010). "The *Escherichia coli* CRISPR System Protects From λ Lysogenization, Lysogens, and Prophage Induction," *Journal of Bacteriology* 192(23):6291-6294, Supplemental Material, 2 pages.
- Edgar et al. (Dec. 2010). "The *Escherichia coli* CRISPR System Protects From λ Lysogenization, Lysogens, and Prophage Induction," *Journal of Bacteriology* 192(23):6291-6294.
- Esvelt, K.M. et al. (Nov. 2013). "Orthogonal Cas9 Proteins for RNA-Guided Gene Regulation and Editing," *Nature Methods* 10(11):1116-1123.
- Ex Parte Re-Exam, mailed Apr. 21, 2021, for U.S. Appl. No. 90/014,705, filed Mar. 26, 2021, for U.S. Patent Reexamination 10,953,090, 15 pages. (2.34).
- Ex Parte Re-Exam, mailed Apr. 30, 2021, for U.S. Appl. No. 90/014,681, filed Feb. 16, 2021, for U.S. Patent Reexamination 10,920,222, 25 pages.
- Ex Parte Re-Exam, mailed Dec. 10, 2018, for U.S. Appl. No. 90/014,184, filed Aug. 10, 2018, for U.S. Patent Reexamination 9,701,964 102 pages.
- Ex Parte Re-Exam, mailed Feb. 22, 2021, for U.S. Appl. No. 16/700,856, filed Dec. 2, 2019, for U.S. Patent Reexamination 10,920,222, 459 pages.
- Ex Parte Re-Exam, mailed Mar. 23, 2021, for U.S. Appl. No. 16/453,604, filed Jun. 26, 2019, for U.S. Patent Reexamination 10,953,090, 235 pages.
- Ex Parte Re-Exam, mailed Mar. 24, 2021, for U.S. Appl. No. 90/014,681, filed Mar. 24, 2021, for U.S. Patent Reexamination 10,920,222, 18 pages.
- Extended European Search Report, dated Jul. 27, 2020, for European Patent Application No. 20155001.9, 9 pages.
- Extended European Search Report, dated Sep. 24, 2020, for European Patent Application No. 20154858.3, 12 pages.
- Fact Sheet (Oct. 2010). "Antimicrobial Resistance," National Institutes of Health, 1-2.
- Foca, A. et al. (2015, e-pub. Apr. 7, 2015). Gut Inflammation and Immunity: What is the Role of the Human Gut Virome? *Mediators of Inflammation* 2015(326032):1-7.
- Galperin, M.Y. (Dec. 2013). "Genome Diversity of Spore-Forming Firmicutes," *Microbiology Spectrum* 1(2):TBS-0015-2012, 27 pages.
- Garneau, J. E. et al. (Nov. 4, 2010). "The CRISPR/Cas Bacterial Immune System Cleaves Phage and Plasmid DNA," *Nature* 468(7320):67-71, 28 pages.
- Garon, E.B. et al. (Oct. 2015). "Current Perspectives in Immunotherapy for Non-Small Cell Lung Cancer," *Seminars in Oncology* 42(5 Supp. 2):S11-S18.
- Garrett W.S. et al. (Oct. 5, 2007). "Communicable Ulcerative Colitis Induced by T-Bet Deficiency in the Innate Immune System," *Cell* 131(1):33-45, 23 pages.
- Gauer, R.L. et al. (Jul. 1, 2013). "Early Recognition and Management of Sepsis in Adults: The First Six Hours," *American Family Physician* 88(1):44-53.
- Geller, L.T. et al. (Sep. 15, 2017). "Potential Role of Intratumor Bacteria in Mediating Tumor Resistance to the Chemotherapeutic Drug Gemcitabine," *Cancer*, 1156-1160, 6 pages.
- Golubovskaya, V. et al. (Mar. 15, 2016). "Different Subsets of T Cells, Memory, Effector Functions, and CAR-T Immunotherapy," *Cancers* 8(36), 12 pages.
- Gomaa et al. (Jan. 28, 2014). "Programmable Removal of Bacterial Strains by Use of Genome-Targeting CRISPR-Cas Systems," *mBio*, 5(1):e000928-13.
- Gomaa, A.A. et al. (Jan./Feb. 2014). Supplemental Material to "Programmable Removal of Bacterial Strains by Use of Genome-Targeting CRISPR-Cas Systems," *American Society for Microbiology* 5(1):1-9.
- Goodall, E.C.A. et al. (Feb. 20, 2018). "The Essential Genome of *Escherichia coli* K-12," *Am. Society for Microbiology—mBio* 9(1):e02096-17, 18 pages.
- Gopalakrishnan, V. et al. (Jan. 5, 2018). "Gut Microbiome Modulates Response to Anti-PD-1 Immunotherapy in Melanoma Patients," *Science* 359:97-103, 20 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Green, J. (Jul. 20, 2018). Colgate <https://www.colgate.com/en-us/oral-health/conditions/mouth-sores-and-infections/eight-common-oral-infections-0615>, last visited Jul. 20, 2018, 4 pages.
- Gudbergsdottir, S. et al. (2011, e-pub. Nov. 18, 2010). "Dynamic Properties of the Sulfolobus CRISPR/Cas and CRISPR/Cmr Systems When Challenged With Vector-Borne Viral and Plasmid Genes and Protospacers," *Molecular Microbiology* 79(1):35-49.
- Guedan, S. et al. (Aug. 14, 2014). "ICOS-Based Chimeric Antigen Receptors Program Bipolar TH17/TH1 Cells," *Blood* 124(7):1070-1080.
- Gupta, R. et al. (2011). "P-27/HP Endolysin as Antibacterial Agent for Antibiotic Resistant *Staphylococcus aureus* of Human Infections," *Curr. Microbiol.* 63:39-45.
- Hansen, J.J. et al. (Mar. 2015). "Therapeutic Manipulation of the Microbiome in IBD: Current Results and Future Approaches," *Curr. T. Options Gastroentrol.* 13(1):1-18.
- Hargreaves, K.R. et al. (Aug. 26, 2014). "Abundant and Diverse Clustered Regularly Interspaced Short Palindromic Repeat Spacers in *Clostridium difficile* Strains and Prophages Target Multiple Phage Types within This Pathogen," *mBio* 5(5):e01045-13.
- Harrington, L.E. (Nov. 2005, e-pub. Oct. 2, 2005). "Interleukin 17-producing CD4+ Effector T Cells Develop Via a Lineage Distinct From the T Helper Type 1 and 2 Lineages," *Nat. Immunol.* 6(11):1123-1132.
- Hartland, E.L. et al. (Apr. 30, 2013). "Enteropathogenic and Enterohemorrhagic *E. coli*: Ecology, Pathogenesis, and Evolution," *Frontiers in Cellular and Infection Microbiology* 3(15):1-3.
- Healthline (2019). "Cystic Fibrosis," retrieved from <https://www.healthline.com/health/cystic-fibrosis#prevention>, last visited Aug. 5, 2019, 14 pages.
- Hooper, L.V. et al. (Jun. 8, 2012). "Interactions Between the Microbiota and the Immune System," *Science* 336(6086):1268-1273, 16 pages.
- Horvath, P. et al. (2008, e-pub. Dec. 7, 2007). "Diversity, Activity, and Evolution of CRISPR Loci in *Streptococcus thermophilus*," *Journal of Bacteriology* 190(4):1401-1412.
- Hotta, K. et al. (2011, e-pub. Sep. 20, 2011). "Prognostic Significance of CD45RO+ Memory T Cells in Renal Cell Carcinoma," *British Journal of Cancer* 105:1191-1196.
- Huddleston, J.R. (Jun. 20, 2014). "Horizontal Gene Transfer in the Human Gastrointestinal Tract: Potential Spread of Antibiotic Resistance Genes," *Infection and Drug Resistance* 7:167-176.
- Huo, Y. et al. (Sep. 2014). "Structures of CRISPR Cas3 Offer Mechanistic Insights Into Cascade-Activated DNA Unwinding and Degradations," *Nat. Struct. Mol. Biol.* 21(9):771-777, 21 pages.
- International Search Report and The Written Opinion of the International Searching Authority for PCT/EP2018/066954, dated Oct. 23, 2018, filed Jun. 25, 2018, 14 pages.
- International Search Report and The Written Opinion of the International Searching Authority for PCT/EP2019/057453, dated Aug. 16, 2019, filed Mar. 25, 2019, 21 pages.
- International Search Report for PCT/EP2016/059803, dated Jun. 30, 2016, filed May 3, 2016, 6 pages.
- International Search Report for PCT/EP2018/082053, dated Mar. 14, 2019, filed Nov. 21, 2018, 9 pages.
- Ivanov, I.I. et al. (May 2010). "Segmented Filamentous Bacteria Take the Stage," *Mucosal Immunol.* 3(3):209-212, 7 pages.
- Jiang, W. et al. (Jan. 29, 2013). "RNA-Guided Editing of Bacterial Genomes Using CRISPR-Cas Systems," *Nat. Biotechnology* 31:233-241.
- Jiang, W. et al. (Mar. 2013, e-pub. Sep. 1, 2013). "CRISPR-Assisted Editing of Bacterial Genomes," *Nat. Biotechnol.* 31(3):233-239.
- Jiang, W. et al. (Nov. 2013). "Demonstration of CRISPR/Cas9/sgRNA-Mediated Targeted Gene Modification in *Arabidopsis*, Tobacco, Sorghum and Rice," *Nucleic Acids Research* 41(20):e188, 12 pages.
- Jin, Y. et al. (2019, e-pub. Apr. 23, 2019). "The Diversity of Gut Microbiome is Associated With Favorable Responses to Anti-Programmed Death 1 Immunotherapy in Chinese Patients With NSCLC," *Journal of Thoracic Oncology* 14(8):1378-1389.
- Jinek et al. (Aug. 17, 2012). "A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity," *Science* 337(6096):816-821.
- Khoja, L. et al. (2015). "Pembrolizumab," *Journal for ImmunoTherapy of Cancer* 3(36):1-13.
- Kochenderfer, J.N. et al. (Sep. 2009). "Construction and Pre-clinical Evaluation of an Anti-CD19 Chimeric Antigen Receptor," *J. Immunother.* 32(7):689-702, 26 pages.
- Koonin, E.V. et al. (2017, e-pub. Jun. 9, 2017). "Diversity, Classification and Evolution of CRISPR-Cas Systems," *Current Opinion in Microbiology* 37:67-78.
- Kosiewicz, M.M. et al. (2014, e-pub. Mar. 26, 2014). "Relationship Between Gut Microbiota and Development of T Cell Associated Disease," *FEBS Lett.* 588:4195-4206.
- Kostic, A.D. et al. (Aug. 14, 2013). "Fusobacterium nucleatum Potentiates Intestinal Tumorigenesis and Modulates the Tumor-Immune Microenvironment," *Cell Host Microbe.* 14(2):207-215, 18 pages.
- Krom, R.J. et al. (Jul. 5, 2015). "Engineered Phagemids for Nonlytic, Targeted Antibacterial Therapies," *Nano Letters* 15(7):4808-4813.
- La Scola, B. et al. (Sep. 4, 2008). "The Virophage as a Unique Parasite of the Giant Mimivirus," *Nature Letters* 455:100-104.
- Lopez-Sanchez, M.-J. et al. (2012, e-pub. Jul. 27, 2012). "The Highly Dynamic CRISPR1 System of *Streptococcus agalactiae* Controls the Diversity of its Mobilome," *Molecular Microbiology* 85(6):1057-1071.
- Lu, T.K. et al. (Jul. 3, 2007). "Dispersing Biofilms With Engineered Enzymatic Bacteriophage," *PNAS* 104(27):11197-11202.
- Ludwig, W. et al. (1985). "The Phylogenetic Position of *Streptococcus* and Enterococcus," *Journal of General Microbiology* 131:543-551.
- Luo, M.L. et al. (2015, e-pub. Oct. 17, 2014). "Repurposing Endogenous Type I CRISPR-Cas Systems for Programmable Gene Repression," *Nucleic Acids Research* 43(1):674-681.
- López, P. et al. (Apr. 5, 2016). "Th17 Responses and Natural IgM Antibodies are Related to Gut Microbiota Composition in Systemic Lupus Erythematosus Patients," *Sci. Rep.* 6:24072, 12 pages.
- Macon, B.L. et al. (Jan. 2, 2018). "Acute Nephritides," retrieved from healthline, <https://www.healthline.com/health/acute-nephritic-syndrome#types>, last visited Jul. 20, 2018, 13 pages.
- Magee, M.S. et al. (Nov. 2014). "Challenges to Chimeric Antigen Receptor (CAR)-T Cell Therapy for Cancer," *Discov. Med.* 18(100):265-271, 6 pages.
- Mahoney, K.M. et al. (2015). "The Next Immune-Checkpoint Inhibitors: PD-1/PD-L1 Blockade in Melanoma," *Clinical Therapeutics* 37(4):764-782.
- Makarova, K.S. (Jun. 2011). "Evolution and Classification of the CRISPR-Cas Systems," *Nat. Rev. Microbiol.* 9(6):467-477, 23 pages.
- Mali, P. et al. (Oct. 2013, e-pub. Sep. 27, 2013). "Cas9 as a Versatile Tool for Engineering Biology," *Nature Methods* 10(10):957-963. 16 pages.
- Mancha-Agresti, P. et al. (Mar. 2017). "A New Broad Range Plasmid for DNA Delivery in Eukaryotic Cells Using Lactic Acid Bacteria: In Vitro and In Vivo Assays," *Molecular Therapy: Methods & Clinical Development* 4:83-91.
- Manica, A. et al. (2011, e-pub. Mar. 8, 2011). "In vivo Activity of CRISPR-Mediated Virus Defence in a Hyperthermophilic Archaeon," *Molecular Microbiology* 80(2):481-491.
- Marraffini, L.A. et al. (Dec. 19, 2008). "CRISPR Interference Limits Horizontal Gene Transfer in Staphylococci by Targeting DNA," *Science* 322(5909):1843-1845, 12 pages.
- Martel, B. et al. (2014). "CRISPR-Cas: An Efficient Tool for Genome Engineering of Virulent Bacteriophages," *Nucleic Acids Research* 42(14):9504-9513.
- Martinez, R.M. et al. (Aug. 12, 2016). "Bloodstream Infections," *Microbial Spectrum* 4(4):DMIH2-0031-2016, 34 pages.
- Mayo Clinic (2019). "Pulmonary Embolism," retrieved from <https://www.mayoclinic.org/diseases-conditions/pulmonary-embolism/synnpntons-causes/syc-20354647>, last visited Aug. 5, 2019, 8 pages.
- Mayo Clinic (2020). "Infectious Diseases," retrieved from <https://www.mayoclinic.org/diseases-conditions/infectious-diseases/diagnosis-treatment/drc-20351179>, last visited Jan. 17, 2020, 5 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Mayo Clinic (2020). "Malaria," retrieved from <https://www.mayoclinic.org/diseases-conditions/malaria/diagnosis-treatment/drc-20351190>, last visited Jan. 17, 2020, 3 pages.
- Mayo Clinic (2020). "Sexually Transmitted Diseases (STDs)," retrieved from <https://www.mayoclinic.org/diseases-conditions/sexually-transmitted-diseases/stds/diagnosis-treatment/drc-20351246>, last visited Jan. 17, 2020, 5 pages.
- Mayo Clinic (Jul. 20, 2018). "Bacterial Vaginosis," retrieved from <https://www.mayoclinic.org/diseases-conditions/bacterial-vaginosis/symptoms-causes/syc-20352279>, last visited Jul. 20, 2018, 3 pages.
- Mayo Clinic (Jul. 20, 2018). "Cystitis," retrieved from <https://www.mayoclinic.org/diseases-conditions/cystitis/symptoms-causes/syc-20371306>, last visited Jul. 20, 2018, 10 pages.
- Mayo Clinic (Jul. 20, 2018). "Meningitis," retrieved from <https://www.mayoclinic.org/diseases-conditions/meningitis/symptoms-causes/syc-20350508>, last visited Jul. 20, 2018, 6 pages.
- Mayo Clinic (Jul. 20, 2018). "Pneumonia," retrieved from <https://www.mayoclinic.org/diseases-conditions/pneumonia/symptoms-causes/syc-20354204>, last visited Jul. 20, 2018, 5 pages.
- Mayo Clinic (Mar. 29, 2020). "Liver Disease," retrieved from <https://www.mayoclinic.org/diseases-conditions/liver-problems/diagnosis-treatment/drc-20374507>, last visited Mar. 29, 2020, 8 pages.
- Medina-Aparicio, L. et al. (May 2011, e-pub. Mar. 11, 2011). "The CRISPR/Cas Immune System is an Operon Regulated by LeuO, H—NS, and Leucine-Responsive Regulatory Protein in *Salmonella enterica* Serovar Typhi," *Journal of Bacteriology* 193(10):2396-2407.
- Mei, J.-M. et al. (1997). "Identification of *Staphylococcus aureus* Virulence Genes in a Murine Model of Bacteraemia Using Signature-Tagged Mutagenesis," *Molecular Microbiology* 26(2):399-407.
- Mercenier, A. (1990). "Molecular Genetics of *Streptococcus thermophilus*," *FEMS Microbiology Letters* 87(1-2):61-77.
- Mick, E. et al. (May 2013). "Holding a Grudge: Persisting Anti-Phage CRISPR Immunity in Multiple Human Gut Microbiomes," *RNA Biology* 10(5):900-906.
- Mills, S. et al. (Jan./Feb. 2013). "Movers and Shakers: Influence of Bacteriophages in Shaping the Mammalian Gut Microbiota," *Gut Microbes* 4(1):4-16.
- Mitsuhashi, K. et al. (Mar. 13, 2015). "Association of *fusobacterium* Species in Pancreatic Cancer Tissues With Molecular Features and Prognosis," *Oncotarget* 6(9):7209-7220.
- Nakamura, S. et al. (Nov. 2008). "Metagenomic Diagnosis of Bacterial Infections," *Emerging Infectious Diseases* 14(11):1784-1786.
- Nale, J.Y. et al. (2012). "Diverse Temperate Bacteriophage Carriage in *Clostridium Difficile* 027 Strains," *PLoS One* 7(5):e37263, 9 pages.
- Navarre, L. et al. (2007). "Silencing of Xenogenic DNA by H—NS—Facilitation of Lateral Gene Transfer in Bacteria by a Defense System That Recognizes Foreign DNA," *Genes & Development* 21:1456-1471.
- Nelson, M.H. et al. (2015). "Harnessing the Microbiome to Enhance Cancer Immunotherapy," *Journal of Immunology Research* 2015:Article 368736, 12 pages.
- News (May 22, 2018). "UK Government and Bill & Melinda Gates Foundation Join Carb-X Partnership in Fight Against Superbugs: Millions Earmarked to Boost Research Into New Life-Saving Products to Address the Global Rise of Drug-Resistant Bacteria," *Combating Antibiotic Resistant Bacteria*, 7 pages.
- Noonan, K.A. et al. (May 20, 2015). Adoptive Transfer of Activated Marrow-Infiltrating Lymphocytes Induces Measurable Antitumor Immunity in the Bone Marrow in Multiple Myeloma *Science Translational Medicine* 7(228):288ra78, 14 pages.
- Norris, J.S. et al. (2000). "Prokaryotic Gene Therapy to Combat Multidrug Resistant Bacterial Infection," *Gene Therapy* 7:723-725. Notice of Intent to Issue Ex Parte Reexamination Certificate, mailed Aug. 12, 2019, for U.S. Appl. No. 90/014,184, filed Aug. 10, 2018, 26 pages.
- Nowak, P. et al. (Nov. 28, 2015). "Gut Microbiota Diversity Predicts Immune Status in HIV-1 Infection," *AIDS* 29(18):2409-2418.
- Park, A. (Oct. 18, 2011). "A Surprising Link Between Bacteria and Colon Cancer," *Cancer* retrieved from <http://healthland.time.com/2011/10/18/a-surprising-link-between-bacteria-and-colon-cancer/>, last visited Aug. 27, 2019, 3 pages.
- Park, H. et al. (2005). "A Distinct Lineage of CD4 T Cells Regulates Tissue Inflammation by Producing Interleukin 17," *Nat. Immunol.* 6(11):1133-1141, 24 pages.
- Patterson, A.G. et al. (2017, e-pub. Mar. 27, 2017). "Regulation of CRISPR-Cas Adaptive Immune Systems," *Current Opinion in Microbiology* 37:1-7.
- Patterson, A.G. et al. (Dec. 15, 2016). "Quorum Sensing Controls Adaptive Immunity Through the Regulation of Multiple CRISPR-Cas Systems," *Mol. Cell* 64(6):1102-1108.
- Pawluk, A. et al. (Apr. 15, 2014). "A New Group of Phage Anti-CRISPR Genes Inhibits the Type I-E CRISPR-Cas System of *Pseudomonas aeruginosa*," *mBio*. 5(2):e00896.
- Perez-Chanona, E. et al. (2016, e-pub. Jan. 26, 2016). "The Role of Microbiota in Cancer Therapy," *Current Opinion in Immunology* 39:75-81.
- Pires, D.P. et al. (Sep. 2016, e-pub. Jun. 1, 2016). "Genetically Engineered Phages: A Review of Advances Over the Last Decade," *Microbiology and Molecular Biology Reviews* 80(3):523-543.
- Pul, Ü. et al. (2010, e-pub. Feb. 17, 2010). "Identification and Characterization of *E. coli* CRISPR-cas Promoters and Their Silencing by H—NS," *Molecular Microbiology* 75(6):1495-1512.
- Ramalingam, S.S. et al. (2014). "LB2-Metastatic Non-Small Cell Lung Cancer: Phase II Study of Nivolumab (Anti-PD-1, BMS-936558, ONO-4538) In Patients With Advanced, Refractory Squamous Non-Small Cell Lung Cancer," *International Journal of Radiation Oncology Biology Physics Late Breaking Abstract (LB2)*.
- Ran, F.A. et al. (Apr. 9, 2015). "In Vivo Genome Editing Using *Staphylococcus aureus* Cas9," *Nature* 570(7546):186-191, 28 pages.
- Rashid, T. et al. (2013). "The Role of *Klebsiella* in Crohn's Disease With a Potential for the Use of Antimicrobial Measures," *International Journal of Rheumatology* 2013(Article ID 610393):1-9.
- Ray, K. (Jan. 2020). "Manipulating the Gut Microbiota to Combat Alcoholic Hepatitis," *Nature Reviews Gastroenterology & Hepatology* 17:3, 1 page.
- Request for Ex Parte Reexamination mailed Aug. 10, 2018, for U.S. Appl. No. 15/160,405, now U.S. Pat. No. 9,701,964, 42 pages.
- Request for Ex Parte Reexamination mailed Nov. 1, 2018, for U.S. Appl. No. 15/160,405, now U.S. Pat. No. 9,701,964, 35 pages.
- Richter, C. et al. (2012, e-pub. Oct. 19, 2012). "Function and Regulation of Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) / CRISPR Associated (Cas) Systems," *Viruses* 4(12):2291-2311.
- Ridaura, V.K. et al. (Sep. 6, 2013). "Cultured Gut Microbiota From Twins Discordant for Obesity Modulate Adiposity and Metabolic Phenotypes in Mice," *Science* 341(6150):1241214, 22 pages.
- Roberts, A.P. et al. (Jun. 2009, e-pub. May 20, 2009). "A Modular Master on the Move: The Tn916 Family of Mobile Genetic Elements," *Trends Microbiol.* 17(6):251-258. Abstract Only.
- Rogers, L. et al. (2016). "*Escherichia coli* and Other Enterobacteriaceae: Occurrence and Detection," *Encyclopedia of Food and Health* pp. 545-551.
- Rong, Z. et al. (Mar. 14, 2014). "Homologous Recombination in Human Embryonic Stem Cells Using CRISPR/Cas9 Nickase and a Long DNA Donor Template," *Protein & Cell* 5(4):258-260.
- Routy, B. et al. (Jan. 5, 2018, e-pub. Nov. 2, 2017). "Gut Microbiome Influences Efficacy of PD-1-Based Immunotherapy Against Epithelial Tumors," *Science* 359(6371):91-97.
- Samarzija, D. et al. (2001). "Taxonomy, Physiology and Growth of *Lactococcus Lactis*: A Review," *Mljekarstvo* 51(1):35-48.
- Sapranaukas, R. et al. (Nov. 1, 2011, e-pub. Aug. 3, 2011). "The *Streptococcus thermophilus* CRISPR/Cas System Provides Immunity in *Escherichia coli*," *Nucleic Acids Research* 39(21):9275-9282.
- Schnabi, B.G. (2020). "The Role of *Enterococcus Faecalis* in Alcoholic Liver Disease," retrieved from <https://grantome.com/grant/NIH/O01-BX004594-01A2>, last visited Oct. 20, 2020, 2 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Seed, K.D. et al. (Feb. 27, 2013). "A Bacteriophage Encodes Its Own CRISPR/Cas Adaptive Response to Evade Host Innate Immunity," *Nature* 494(7438):489-491.
- Selle, K. et al. (Apr. 1, 2015). "Harnessing CRISPR-Cas Systems for Bacterial Genome Editing," *Trends in Microbiology* 23(4):225-232.
- Sepsis Alliance. (Dec. 14, 2017). "What Are Vaccines," Retrieved from <https://www.sepsis.org/sepsisand/prevention-vaccinations/>; last visited Jul. 8, 2019, 3 pages.
- Sepsis Alliance. (Jul. 8, 2019). "Prevention," Retrieved from <https://www.sepsis.org/sepsisand/prevention/>; accessed last visited Jul. 8, 2019, 5 pages.
- Shoemaker, N.B. et al. (Feb. 2001). "Evidence for Extensive Resistance Gene Transfer Among *bacteroides* spp. and Among Bacteroides and Other Genera in the Human Colon," *Appl. Environ. Microbiol.* 67(2):561-68.
- Sivan, A. et al. (Nov. 27, 2015, e-pub Nov. 5, 2015). "Commensal Bifidobacterium Promotes Antitumor Immunity and Facilitates Anti-PD-L1 Efficacy," *Science* 350(6264):1084-1089, 13 pages.
- Sivan, A. et al. (Nov. 6, 2014). "Evidence Implicating the Commensal Microbiota in Shaping Anti-Tumor Immunity in Melanoma," *Journal for ImmunoTherapy of Cancer* 2(Suppl. 3):O11, 1 page.
- Skenneron, C.T. et al. (May 2011). "Phage Encoded H-NS: A Potential Achilles Heel in the Bacterial Defence System," *PLoS One* 6(5):e20095.
- Somkuti, G. A. et al. (Apr. 1988). "Genetic Transformation of *Streptococcus thermophilus* by Electroporation," *Biochimie* 70(4):579-585. Abstract Only.
- Sorek, R. et al. (2013, e-pub. Mar. 11, 2013). "CRISPR-Mediated Adaptive Immune Systems in Bacteria and Archaea," *Annual Review of Biochemistry* 82:237-266.
- Sorg, R. A. et al. (2014). "Gene Expression Platform for Synthetic Biology in the Human Pathogen *Streptococcus pneumoniae*," *ACS Synthetic Biology* 4(3):228-239. Abstract Only.
- Soutourina, O.A. et al. (May 9, 2013). "Genome-Wide Identification of Regulatory RNAs in the Human Pathogen *Clostridium difficile*," *PLoS Genet.* 9(5):e1003493, 20 pages.
- Stern, A. et al. (2012). "CRISPR Targeting Reveals a Reservoir of Common Phages Associated With the Human Gut Microbiome," *Genome Research* 22(10):1985-1994.
- Stern, A. et al. (Aug. 2010). Self-Targeting by CRISPR: Gene Regulation or Autoimmunity? *Trends Genet.* 26(8):335-340, 10 pages.
- Stiefel, U. et al. (Aug. 2014, e-pub. May 27, 2014). "Gastrointestinal Colonization With a Cephalosporinase-Producing *Bacteroides* Species Preserves Colonization Resistance Against Vancomycin-Resistant Enterococcus and Clostridium Difficile in Cephalosporin-Treated Mice," *Antimicrob. Agents Chemother.* 58(8):4535-4542.
- Stoebel, D.M. et al. (2008). "Anti-Silencing: Overcoming H—NS-Mediated Repression of Transcription in Gramnegative Enteric Bacteria," *Microbiology* 154:2533-2545.
- Suvorov, A. (1988). "Transformation of Group A Streptococci by Electroporation," *FEMS Microbiology Letters* 56(1):95-100.
- Takaishi, H. et al. (2008). "Imbalance in Intestinal Microflora Constitution Could be Involved in the Pathogenesis of Inflammatory Bowel Disease," *Int. J. Med. Microbiol.* 298:463-472.
- Takeda, T. et al. (2011). "Distribution of Genes Encoding Nucleoid-Associated Protein Homologs in Plasmids," *International Journal of Evolutionary Biology* 2001:685015, 31 pages.
- Tan, J. (Dec. 17, 2015). "Immunotherapy Meets Microbiota," *Cell* 163:1561.
- Tarr, P.I. et al. (Mar. 19-25, 2005). "Shiga-Toxin-Producing *Escherichia coli* and Haemolytic Uraemic Syndrome," *Lancet* 365(9464):1073-1086.
- Todar, K. (2012). "The Normal Bacterial Flora of Humans," *Todar's Online Textbook of Bacteriology*, 8 pages.
- Topalian, S.L. et al. (Jun. 28, 2012). "Safety, Activity, and Immune Correlates of Anti-PD-1 Antibody in Cancer," *N. Engl. J. Med.* 366(26):2443-2454, 19 pages.
- Turnbaugh, P.J. et al. (Dec. 2006). "An Obesity-Associated Gut Microbiome With Increased Capacity for Energy Harvest," *Nature* 444:1027-1131.
- U.S. Appl. No. 62/168,355, filed May 29, 2015, Barrangou, R. et al. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. 1.98(a)(2)(iii) issued by the Office dated Sep. 21, 2004).
- U.S. Appl. No. 62/296,853, filed Feb. 18, 2016, Barrangou, R. et al. (Copy not submitted herewith pursuant to the waiver of 37 C.F.R. 1.98(a)(2)(iii) issued by the Office dated Sep. 21, 2004).
- Uchiyama, J. et al. (2013, e-pub. Mar. 8, 2013). "Characterization of *Helicobacter pylori* Bacteriophage KHP30," *Applied and Environmental Microbiology* 79(10):3176-3184.
- USPTO Interference 106,123—Declaration to Declare Interference Jun. 11, 2020, 11 pages.
- USPTO Interference 106,123—Junior Party Annotated Claims Jul. 9, 2020, 31 pages.
- USPTO Interference 106,123—Junior Party List of Motions Jul. 16, 2020, 6 pages.
- USPTO Interference 106,123—Redeclaration Jul. 21, 2020, 6 pages.
- USPTO Interference 106,123—Rockefeller Clean Claims Jun. 25, 2020, 7 pages.
- USPTO Interference 106,123—Rockefeller Motion 2 (Indefiniteness), Oct. 16, 2020, 24 pages.
- USPTO Interference 106,123—Rockefeller Notice of Lead and Backup Counsel Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Notice of Real Party in Interest Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Notice of Related Proceedings Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Power of Attorney Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Request for File Copies Jun. 25, 2020, 10 pages.
- USPTO Interference 106,123—Rockefeller Revised List of Proposed Motions Aug. 13, 2020, 4 pages.
- USPTO Interference 106,123—Senior Party List of Proposed Motions Jul. 16, 2020, 5 pages.
- USPTO Interference 106,123—SNIPR Clean Claims Jun. 25, 2020, 27 pages.
- USPTO Interference 106,123—SNIPR Motion 2 (Lack of Enablement and Written Description), Oct. 16, 2020, 32 pages.
- USPTO Interference 106,123—SNIPR Motion 4 (Deny Benefit to Count 1), Oct. 16, 2020, 16 pages.
- USPTO Interference 106,123—SNIPR Motion 5 (Substitute Count), Oct. 16, 2020, 41 pages.
- USPTO Interference 106,123—SNIPR Motion 6 (Motion to Designate Claims as Not Corresponding to Count 1 or Proposed Count 2), Oct. 16, 2020, 24 pages.
- USPTO Interference 106,123—SNIPR Notice of Lead and Backup Counsel Jun. 25, 2020, 4 pages.
- USPTO Interference 106,123—SNIPR Notice of Related Proceedings Jun. 25, 2020, 4 pages.
- USPTO Interference 106,123—SNIPR Real Party in Interest Jun. 25, 2020, 3 pages.
- USPTO Interference 106,123—SNIPR Request for File Copies Jun. 25, 2020, 10 pages.
- USPTO Interference 106,123—Standing Order Jun. 11, 2020, 81 pages.
- USPTO Interference 106,123—Joint Stipulated Extension of Time, Sep. 4, 2020, 4 pages.
- USPTO Interference 106,123—Junior Party Revised List of Motions Aug. 13, 2020, 6 pages.
- USPTO Interference 106,123—Memorandum, Jan. 19, 2021, 6 pages.
- USPTO Interference 106,123—Notice of Cross Examination—van der Oost, Dec. 1, 2020, 3 pages.
- USPTO Interference 106,123—Order Additional Applications 37 C.F.R. § 41.104(a), Sep. 3, 2020, 6 pages.
- USPTO Interference 106,123—Order Authorizing Motions and Setting Times 37 C.F.R. 11.104(c) and 121 Aug. 24, 2020, 10 pages.

(56)

References Cited

OTHER PUBLICATIONS

- USPTO Interference 106,123—Order—Additional Applications, Jan. 13, 2021, 6 pages.
- USPTO Interference 106,123—Order—Bd.R. 109(b)—Authorizing Office Records Jul. 21, 2020, 3 pages.
- USPTO Interference 106,123—Order—Video Dispositions 37 C.F.R. § 41.104(a), Sep. 25, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller List of Exhibits, Oct. 16, 2020, 4 pages.
- USPTO Interference 106,123—Rockefeller List of Exhibits, Nov. 13, 2020, 4 pages.
- USPTO Interference 106,123—Rockefeller List of Exhibits, Feb. 19, 2021, 5 pages.
- USPTO Interference 106,123—Rockefeller Motion 1 (Lack of Written Description), Oct. 16, 2020, 30 pages.
- USPTO Interference 106,123—Rockefeller Motion 3 (To Add a Claim), Nov. 13, 2020, 36 pages.
- USPTO Interference 106,123—Rockefeller Notice of Settlement Discussions, Oct. 21, 2020, 3 pages.
- USPTO Interference 106,123—Rockefeller Order—Responsive Motion 37 C.F.R. § 41.121(a)(2), Nov. 2, 2020, 2 pages.
- USPTO Interference 106,123—Rockefeller Reply 1, Feb. 19, 2021, 51 pages.
- USPTO Interference 106,123—Rockefeller Reply 2, Feb. 19, 2021, 37 pages.
- USPTO Interference 106,123—Rockefeller Reply 3, Feb. 19, 2021, 48 pages.
- USPTO Interference 106,123—Rockefeller Updated Notice of Related Proceedings, Nov. 13, 2020, 3 pages.
- USPTO Interference 106,123—SNIPR Exhibit List, Oct. 16, 2020, 7 pages.
- USPTO Interference 106,123—SNIPR Exhibit List, Feb. 19, 2021, 8 pages.
- USPTO Interference 106,123—SNIPR Motion 1 (Terminate Interference as Contrary to AIA), Oct. 16, 2020, 20 pages.
- USPTO Interference 106,123—SNIPR Request for Oral Argument, Mar. 12, 2021, 4 pages.
- USPTO Interference 106,123—Rockefeller Request for Oral Argument, Mar. 12, 2021, 3 pages.
- USPTO Interference 106,123—SNIPR Reply 1, Feb. 19, 2021, 19 pages.
- USPTO Interference 106,123—SNIPR Reply 2, Feb. 19, 2021, 42 pages.
- USPTO Interference 106,123—SNIPR Reply 4, Feb. 19, 2021, 28 pages.
- USPTO Interference 106,123—SNIPR Reply 5, Feb. 19, 2021, 44 pages.
- USPTO Interference 106,123—SNIPR Reply 6, Feb. 19, 2021, 27 pages.
- Veeranagouda, Y. et al. (Jun. 4, 2014). “Identification of Genes Required for the Survival of *B. fragilis* Using Massive Parallel Sequencing of a Saturated Transposon Mutant Library,” *BMC Genomics* 15:429, 11 pages.
- Vega, N.M. et al. (Oct. 2014). “Collective Antibiotic Resistance: Mechanisms and Implications,” *Curr. Opin. Microbiol.* 21:28-34, 14 pages.
- Vercoe, R.B. et al. (Apr. 18, 2013). “Cytotoxic Chromosomal Targeting by CRISPR/Cas Systems Can Reshape Bacterial Genomes and Expel or Remodel Pathogenicity Islands,” *PLoS Genetics* 9(4):e1003454, 13 pages.
- Villarino, N.F. et al. (Feb. 23, 2016, e-pub. Feb. 8, 2016). “Composition of the Gut Microbiota Modulates the Severity of Malaria,” *Proc. Natl. Acad. Sci. USA* 113(8):2235-2240.
- Vétizou, M. et al. (Nov. 27, 2015, e-pub Nov. 5, 2015). “Anticancer Immunotherapy by CTLA-4 Blockade Relies on the Gut Microbiota,” *Science* 350(6264):1079-1084, 13 pages.
- Walters, W.A. et al. (Nov. 17, 2014). “Meta-Analyses of Human Gut Microbes Associated With Obesity and IBD,” *FEBS Letters* 588(22):4223-4233, 34 pages.
- Wang, I.-N. et al. (2000). “HOLINS: The Protein Clocks of Bacteriophage Infections,” *Annu. Rev. Microbiol.* 54:799-825.
- Wang, J. et al. (2019). “Core Gut Microbiota Analysis of Feces in Healthy Mouse Model,” Supplementary Information, 12 pages.
- Wang, J. et al. (Apr. 2019). “Core Gut Bacteria Analysis of Healthy Mice,” *Frontiers in Microbiology* 10(887):1-14.
- Wegmann, U. et al. (Apr. 2007). “Complete Genome Sequence of the Prototype Lactic Acid Bacterium *Lactococcus lactis* Subsp. *cremoris* MG 1363,” *Journal of Bacteriology* 189(8):3256-3270.
- Wei, Y. et al. (2015, e-pub. Jan. 14, 2015). “Sequences Spanning the Leader-Repeat Junction Mediate CRISPR Adaptation to Phage in *Streptococcus thermophilus*,” *Nucleic Acids Research* 43(3):1749-1758.
- Weir, T.L. et al. (Aug. 6, 2013). “Stool Microbiome and Metabolome Differences Between Colorectal Cancer Patients and Healthy Adults,” *PLOS One* 8(8):e70803, 10 pages.
- Westra, E.R. et al. (Jun. 8, 2012). “CRISPR Immunity Relies on the Consecutive Binding and Degradation of Negatively Supercoiled Invader DNA by Cascade and Cas3,” *Molecular Cell* 46:595-605.
- Westra, E.R. et al. (Sep. 1, 2010, e-pub. Aug. 18, 2010). “H—NS-Mediated Repression of CRISPR-Based Immunity in *Escherichia coli* K12 Can be Relieved by the Transcription Activator LeuO,” *Molecular Microbiology* 77(6):1380-1393.
- Westwater, C. et al. (2002). “Development of a Pl Phagemid System for the Delivery of DNA Into Gram-Negative Bacteria,” *Microbiology* 148:943-950.
- Westwater, C. et al. (Apr. 2003). “Use of Genetically Engineered Phage to Deliver Antimicrobial Agents to Bacteria: An Alternative Therapy for Treatment of Bacterial Infections,” *Antimicrobial Agents and Chemotherapy* 47(4):1301-1307.
- Wexler, H.M. (Oct. 2007). “Bacteroides: the Good, the Bad, and the Nitty-Gritty,” *Clinical Microbiology Reviews* 20(4):593-621.
- Written Opinion for PCT Application No. PCT/EP2016/059803, dated Jun. 30, 2016, filed May 3, 2016, 6 pages.
- Written Opinion for PCT/EP2018/082053, dated Mar. 14, 2019, filed Nov. 21, 2018, 6 pages.
- Xie, Z. et al. (2013, e-pub. Aug. 9, 2013). “Development of a Tunable Wide-Range Gene Induction System Useful for the Study of Streptococcal Toxin-Antitoxin Systems,” *Applied and Environmental Microbiology* 79(20):6375-6384.
- Xu, T. et al. (Jul. 2015). “Efficient Genome Editing in *Clostridium cellulolyticum* via CRISPR-Cas9 Nickase,” *Applied and Environmental Microbiology* 81(13):4423-4431.
- Yang, Y. et al. (Jun. 5, 2014, e-pub. Apr. 13, 2014). “Focused Specificity of Intestinal Th17 Cells Towards Commensal Bacterial Antigens,” *Nature* 510(7503):152-156, 29 pages.
- Yao, J. et al. (2016, e-pub. May 9, 2016). “A Pathogen-Selective Antibiotic Minimizes Disturbance to the Microbiome,” *Antimicrob. Agents Chemother.*, 24 pages.
- Yosef, I. et al. (2011). “High-Temperature Protein G is Essential for Activity of the *Escherichia coli* Clustered Regularly Interspaced Palindromic Repeats (CRISPR)/Cas System,” *Proc. Natl. Acad. Sci. USA* 108(50):20136-20141.
- Yosef, I. et al. (Jun. 9, 2015). “Temperate and Lytic Bacteriophages Programmed to Sensitize and Kill Antibiotic-Resistant Bacteria,” *Proc. Natl. Acad. Sci. USA* 112(23):7267-7272.
- Young, R. et al. (1995). “Holins: Form and Function in Bacteriophage Lysis,” *FEMS Microbiology Reviews* 17:191-205.
- YourGenome: CRISPR/CAS9, retrieved from <https://www.yourgenome.org/facts/what-is-crispr-cas9>, last visited Jan. 6, 2020, 8 pages.
- Yu, Z. et al. (Mar. 21, 2014). “Various Applications of TALEN- and CRISPR/Cas9-Mediated Homologous Recombination to Modify the *Drosophila* Genome,” *Biology Open* 3(4):271-280.
- Zhang, R. et al. (2009, e-pub. Oct. 30, 2008). “DEG 5.0, A Database of Essential Genes in Both Prokaryotes and Eukaryotes,” *Nucleic Acids Research* 37:D455-D458.
- Zhang, T. et al. (Sep. 24, 2016). “The Efficacy and Safety of Anti-PD-1/PD-L1 Antibodies for Treatment of Advanced or Refractory Cancers: A Meta-Analysis,” *Oncotarget* 7(45):73068-73079.
- Zhang, X.Z. (2011). “Simple, Fast and High-Efficiency Transformation System for Directed Evolution of Cellulase in *Bacillus Subtilis*,” *Microbial Biotechnology* 4(1):98-105.

(56)

References Cited

OTHER PUBLICATIONS

- Zitvogel, L. et al. (Jan. 2015), "Cancer and the Gut Microbiota: An Unexpected Link," *Sci. Transl. Med.* 7(271):271ps1, 10 pages.
- Hase, K. (Nov. 2014), "Intestinal Microbiota and Immunity," *Infectious Disease (in Japanese)*. 44(6):193-200 22 pages. English Translation.
- Ichikawa, M. et al. (Dec. 2019). "The Relationship Between Gut Microbiome, Immune System, and Cancer," *Jpn. J. Cancer Chemother* 16(12):1807-1813. English Abstract.
- Abedon, S.T. et al. (Dec. 2003). "Experimental Examination of Bacteriophage Latent-Period Evolution as a Response to Bacterial Availability," *Applied and Environmental Microbiology* 69(12):7499-7506.
- Advisory Action, dated Dec. 9, 2021 for U.S. Appl. No. 90/014,705, filed Mar. 23, 2021, 9 pages.
- Bellanger, X. et al. (Jul. 1, 2014, e-pub. Jan. 27, 2014). "Conjugative and Mobilizable Genomic Islands in Bacteria: Evolution and Diversity," *FEMS Microbiology Reviews* 38(2014):720-760.
- Cronan, J.E. (Jan. 2013). "Improved Plasmid-Based System for Fully Regulated Off-To-On Gene Expression in *Escherichia coli*: Application to Production of Toxic Proteins," *Plasmid* 69(1):81-89, 17 pages.
- European Office Action, dated Jun. 29, 2021, for European Patent Application No. 16719873.8, 24 pages.
- European Search Report, dated Oct. 4, 2021, for European Patent Application No. 21170379.8, 6 pages.
- European Search Report, dated Oct. 8, 2021, for European Patent Application No. 21170380.6, 7 pages.
- Ex Parte Re-Exam Communication Transmittal Form, dated Jun. 30, 2021, for U.S. Appl. No. 90/014,705, for U.S. Patent Reexamination 10,953,090, 26 pages.
- Ex Parte Re-Exam, mailed Dec. 16, 2021, for U.S. Appl. No. 90/014,877, filed Oct. 6, 2021, for U.S. Patent Reexamination 10,953,090, 12 pages.
- Ex Parte Re-Exam, mailed Nov. 15, 2021, for U.S. Appl. No. 90/014,877, 12 pages.
- Ex Parte Re-Exam, mailed Sep. 27, 2021, for U.S. Appl. No. 90/014,705, filed Mar. 23, 2021, for U.S. Patent Reexamination 10,953,090, 16 pages.
- Gutierrez, B. et al. (Apr. 30, 2018). "Genome-Wide CRISPR-Cas9 Screen in *E. coli* Identifies Design Rules for Efficient Targeting," 22 pages.
- Hamanishi, J. et al. (2016, e-pub. Feb. 22, 2016). "PD-1/PD-L1 Blockade in Cancer Treatment: Perspectives and Issues," *International Journal of Clinical Oncology* 21:462-473.
- Johnson, C. M. et al. (Nov. 23, 2015). "Integrative and Conjugative Elements (ICEs): What They Do and How They Work," *Annual Review of Genetics* 49(1):577-601, 33 pages.
- Kaulich, M. et al. (2015, e-pub. Jan. 13, 2015). "Efficient CRISPR-AAV Engineering of Endogenous Genes to Study Protein Function by Allele-Specific RNAi," *Nucleic Acids Research* 43(7):e45, 8 pages.
- Keskin, H. et al. (Nov. 20, 2014). "Transcript-RNA-Templated DNA Recombination and Repair," *Nature* 515:436-439.
- Kugelberg, E. et al. (Aug. 2005). "Establishment of a Superficial Skin Infection Model in Mice by Using *Staphylococcus aureus* and *Streptococcus pyogenes*," *Antimicrob Agents Chemother* 49(8):3435-3441.
- Leshem, A. et al. (Sep. 29, 2020). "The Gut Microbiome and Individual-Specific Responses to Diet," *mSystems* 5(5):e00665-20, 12 pages.
- Office Action, dated Nov. 4, 2021 for U.S. Appl. No. 90/014,705, filed Mar. 23, 2021, 7 pages.
- Pastagia, N. et al. (Feb. 2011). "A Novel Chimeric Lysin Shows Superiority to Mupirocin for Skin Decolonization of Methicillin-Resistant and -Sensitive *Staphylococcus aureus* Strains," *Antimicrobial Agents and Chemotherapy* 55(2):738-744.
- Rea, K. et al. (2020, e-pub. Nov. 14, 2019). "Gut Microbiota: A Perspective for Psychiatrists," *Neuropsychobiology* 79:50-62.
- Request for Ex Parte Reexamination under 35 U.S.C. § 302 and 37 C.F.R. § 1.510, dated Feb. 16, 2021, 72 pages.
- Roberts, A.P. et al. (Dec. 1, 2003). "Development of an Integrative Vector for the Expression of Antisense RNA in *Clostridium difficile*," *Journal of Microbiological Methods* 55(3):617-624.
- Saito, H. et al. (Jun. 15, 2016, e-pub. Apr. 12, 2016). "Adoptive Transfer of CD8+ T Cells Generated From Induced Pluripotent Stem Cells Triggers Regressions of Large Tumors Along With Immunological Memory," *Cancer Research* 76(12):3473-3483.
- Svenningsen, S.L. et al. (Mar. 22, 2005). "On the Role of Cro in λ Prophage Induction," *PNAS* 102(12):4465-4469.
- Tlaskalová-Hogenová, H. et al. (2011, e-pub. Jan. 31, 2011). "The Role of Gut Microbiota (Commensal Bacteria) and the Mucosal Barrier in the Pathogenesis of Inflammatory and Autoimmune Diseases and Cancer: Contribution of Germ-Free and Gnotobiotic Animal Models of Human Diseases," *Cellular & Molecular Immunology* 8:110-120.
- USPTO Interference 106,123—Decision on Motions, Sep. 7, 2021, 18 pages.
- USPTO Interference 106,123—Judgement, Nov. 19, 2021, 3 pages.
- USPTO Interference 106,123—SNIPR Notice of Appeal, Dec. 14, 2021, 28 pages.
- USPTO Interference 106,123—Order—Show Cause, Aug. 19, 2021, 4 pages.
- USPTO Interference 106,123—Rockefeller Notice, Aug. 13, 2021, 3 pages.
- USPTO Interference 106,123—Rockefeller Response to Show Cause, Sep. 7, 2021, 7 pages.
- USPTO Interference 106,123—Rockefeller Updated Notice of Related Proceedings, Jul. 15, 2021, 3 pages.
- Waters, J. L. et al. (Nov./Dec. 2013). "Regulation of CTnDOT Conjugative Transfer is a Complex and Highly Coordinated Series of Events," *MBIO* 4(6):e00569-13, 8 pages.
- Wu, J. et al. (Jun. 2019). "Fusobacterium nucleatum Contributes to the Carcinogenesis of Colorectal Cancer by Inducing Inflammation and Suppressing Host Immunity," *Translational Oncology* 12(6):846-851.
- Ericsson, A.C. et al. (Sep. 10, 2015). "Differential Susceptibility to Colorectal Cancer Due to Naturally Occurring Gut Microbiota," *Oncotarget* 6(32):33689-33707.
- Ex Parte Re-Exam Communication Transmittal Form, dated May 25, 2022, for Control No. 901014105, for U.S. Patent Reexamination of U.S. Pat. No. 10,953,090, 19 pages.
- Gur, C. et al. (Feb. 17, 2015). "Binding of the Fap2 Protein of *Fusobacterium nucleatum* to Human Inhibitory Receptor TIGIT Protects Tumors from Immune Cell Attack," *Immunity* 42(2):344-355.
- Mima, K. et al. (Aug. 1, 2015, e-pub. Jun. 4, 2014). "Fusobacterium nucleatum and T-Cells in Colorectal Carcinoma," *JAMA Oncol.* 1(5):653-661.
- Novosiadly, R. et al. (2016, e-pub. Mar. 30, 2016). "High-Content Molecular Profiling of T-Cell Therapy in Oncology," *Molecular Therapy—Oncolytics* 3:16009, 6 pages.
- Request for Ex Parte Re-Exam Under 35 U.S.C. § 302 and 37 C.F.R. § 1.510 dated Jun. 7, 2022, for U.S. Pat. No. 11,291,723, 328 pages.
- Request for Ex Parte Re-Exam Under 35 U.S.C. § 302 and 37 C.F.R. § 1.510 dated Jun. 7, 2022, for U.S. Pat. 11,351,252, 219 pages.
- West, N.R. et al. (Dec. 14, 2015). "Immunotherapy Not Working? Check Your Microbiota," *Cancer cell* 28:687-689.

* cited by examiner

FIG. 1

pBAV1KT5-XyIR-mCherry-PldhA




FIG. 2




FIG. 3




FIG. 4

Commensal gut bacteria

E.coli- MG1655 (40 hrs)





FIG. 4 (Continued)

'Relative' of target species

S.mutans (40 hrs)



L.lactis (40 hrs)





FIG. 4 (Continued)

Target species

***S.thermophilus*-B (40 hrs)**



***S.thermophilus*-LMD9 (40hrs)**




FIG. 5




FIG. 6




FIG. 8




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 16





FIG. 17



SELECTIVELY ALTERING MICROBIOTA FOR IMMUNE MODULATION

CROSS REFERENCE TO RELATED APPLICATION

This application is a Continuation Application of U.S. patent application Ser. No. 16/453,598, which was filed on Jun. 26, 2019, which is a Continuation Application of U.S. patent application Ser. No. 16/192,752, which was filed on Nov. 15, 2018 (now U.S. Pat. No. 10,363,308), which is a Continuation Application of U.S. patent application Ser. No. 15/820,296, which was filed on Nov. 21, 2017 (now U.S. Pat. No. 10,195,273), which is a Continuation Application under 35 U.S.C. § 120 of International Patent Application No. PCT/EP2017/063593, which was filed on Jun. 4, 2017, which claims priority benefit to United Kingdom Patent Application No. GB1609811.3, which was filed on Jun. 5, 2016, the contents of each of which are incorporated herein by reference in their entirety.

SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

The content of the following submission on ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing (file name: 786212000233SEQLIST.TXT, date recorded: Dec. 22, 2020, size: 8 KB).

FIELD OF THE INVENTION

The invention relates to methods of modulating immune cells in a patient (endogenous cells of the patient and/or administered cells, such as via adoptive cell therapy) by altering microbiota of the patient. The invention also relates to methods of modulating treatments or therapies in a subject organism by altering microbiota of the subject. The invention also relates cell populations, systems, kits and other means for effecting this. In an example, advantageously selective targeting of a particular species in a human gut microbiota using guided nucleic acid modification is carried out to effect the alteration.

BACKGROUND OF THE INVENTION

One approach to immunotherapy involves engineering patients' own (or a donor's) immune cells to express cell-surface antigen receptors (CARs) that recognise and attack tumours. Although this approach, called adoptive cell transfer (ACT), has been restricted to small clinical trials so far, treatments using these engineered immune cells have generated some remarkable responses in patients with advanced cancer.

The Chimeric Antigen Receptor (CAR) consists of an antibody-derived targeting domain fused with T-cell signaling domains that, when expressed by a T-cell, endows the T-cell with antigen specificity determined by the targeting domain of the CAR. CARs can potentially redirect the effector functions of a T-cell towards any protein and non-protein target expressed on the cell surface as long as an antibody-based targeting domain is available. This strategy thereby avoids the requirement of antigen processing and presentation by the target cell and is applicable to non-classical T-cell targets like carbohydrates. This circumvention of HLA-restriction means that the CAR T-cell approach can be used as a generic tool broadening the potential of

applicability of adoptive T-cell therapy. See, eg, *Methods Mol Biol.* 2012; 907:645-66. doi: 10.1007/978-1-61779-974-7_36, "Chimeric antigen receptors for T-cell based therapy", Cheadle E J et al.

5 The first CAR-T construct was described in a 1989 paper by immunotherapy pioneer Zelig Eshhar in PNAS. The structure of the CAR now comprises a transmembrane polypeptide chain which is a chimaera of different domains from different cellular proteins. For example, the CAR has an extracellular part joined (often by a linker and/or a hinge region) to an intracellular part, with a transmembrane portion of the CAR embedding the receptor in the membrane of an immune cell, normally a T-cell. The extracellular moiety includes an antibody binding site (usually in the form of an scFv, such as derived from a mouse mAb) that recognizes a target antigen, that commonly is a tumour associated antigen (TAA) on the surface of cancer cells. Antigen recognition in this way dispenses with the need to rely on TCRs that require MHC-restricted antigen presentation, and where binding affinities may be relatively low. The intracellular moiety of the CAR typically includes a CD3-zeta (CD3) domain for intracellular signaling when antigen is bound to the extracellular binding site. Later generation CARs also include a further domain that enhances T-cell mediated responses, which often is a 4-1BB (CD137) or CD28 intracellular domain. On encountering the cognate antigen ligand for the CAR binding site, the CAR can activate intracellular signaling and thus activation of the CAR T-cell to enhance tumour cell killing.

30 Most CAR-Ts expand in vivo so dose titration in a conventional sense is difficult, and in many cases the engineered T-cells appear to be active "forever"—i.e., the observation of on-going B-cell aplasia seen in most of the CD19 CAR-T clinical studies to date. This poses a serious problem for CAR T-cell approaches. Some observed risks are discussed in *Discov Med.* 2014 November; 18(100):265-71, "Challenges to chimeric antigen receptor (CAR)-T cell therapy for cancer", Magee M S & Snook A E, which explains that the first serious adverse event following CAR-T cell treatment occurred in a patient with colorectal cancer metastatic to the lung and liver (Morgan et al., 2010). This patient was treated with T cells expressing a third-generation CAR targeting epidermal growth factor receptor 2 (ERBB2, HER2). The CAR contained an scFv derived from the 4D5 antibody (trastuzumab) that is FDA approved for the treatment of HER2-positive breast cancers (Zhao et al., 2009). The patient developed respiratory distress within 15 minutes of receiving a single dose of 1010 CAR-T cells, followed by multiple cardiac arrests over the course of 5 days, eventually leading to death. Serum analysis four hours after treatment revealed marked increases in the cytokines IFN γ , GM-CSF, TNF α , IL-6, and IL-10. CAR-T cells were found in the lung and abdominal and mediastinal lymph nodes, but not in tumour metastases. The investigators attributed toxicity to recognition of HER2 in lung epithelium resulting in inflammatory cytokine release producing pulmonary toxicity and cytokine release syndrome (CRS) causing multi-organ failure (Morgan et al., 2010). Trials utilizing second-generation HER2-targeted CARs derived from a different antibody (FRP5) following conservative dose-escalation strategies are currently underway for a variety of HER2+ malignancies by other investigators (clinicaltrials.gov identifiers NCT01109095, NCT00889954, and NCT00902044).

65 A variation on the CAR T-cell theme are antibody-coupled T-cell receptor (ACTR) therapeutics, which use CD16A (FC γ R1IIIA) to bind to Fc regions of tumour-specific

IgG (see eg, WO2015/058018, US2015139943). The aim is to enable more control of CAR T-cell activity in vivo by titrating IgG administered to patients. The CD16 binding sites of the CAR-T-cells may be free, however, to also bind to endogenous IgG of the patients and this reduces the attractiveness of the approach. The approach also needs to consider the inherently long half-life of IgG in the body (around 20 days for IgG in man), which may limit control of CAR-cell activity. Ongoing studies may assess the risk of this.

It would be desirable to provide an alternative way to modulate (downregulate or upregulate) immune cell-based therapies, like CAR-T-cell approaches and other cell-based approaches. It would also be desirable to provide a way to address diseases and conditions mediated by endogenous immune cells, such as autoimmune, inflammatory and infectious diseases and conditions.

STATEMENTS OF INVENTION

The invention provides guided nucleases, host cell modifying (HM)-CRISPR/Cas systems, gRNAs, HM-arrays, HM-crRNA, HM-Cas, HM-TALENs, HM-meganucleases, HM-zinc fingers and methods as set out in the claims herein.

Medical practice often involves the administration of antibiotics to patients. Such treatments can typically involve administration of broad-spectrum antibiotics, or antibiotics that target many gram-positive bacterial species or many gram-negative species without discrimination. Similarly, use of broad-spectrum antibiotics in farming and agriculture, for example, raise environmental concerns, including entry of such antibiotics into the human and animal food chain which may be deleterious to health and may add to development of microbial resistance. Rather, the invention involves selective targeting of a first microbiota species or strain. As shown in the worked examples herein, selective targeting of a particular bacterial species has been achieved using guided nuclease targeting of the genome of the selected species, whilst at the same time sparing phylogenetically related species and strains. Furthermore, the invention realises the role that microbiota bacteria and archaea play in shaping immune function in humans and animals, as discussed further below.

Thus, the invention relates to methods of modulating immune cells in a patient (endogenous cells of the patient and/or administered cells, such as via adoptive cell therapy) by altering microbiota of the patient. In an example, advantageously selective targeting of a species in a microbiota (eg, gut microbiota) is carried out to effect the alteration. Selective targeting may, for example, avoid targeting of related species or strains, such as species of the same phylum or such as a different strain of the same species.

For example, the invention provides for modulating immune cell-based or other therapy of diseases and conditions in patients and subjects by altering microbiota, as well as systems, kits and other means for effecting this.

For example, the invention provides for treating or reducing diseases and conditions in patients by altering microbiota, wherein the diseases and conditions are those mediated by immune cells (eg, T-cells) or addressed by altering immune cell activities or populations in patients. Embodiments are cancers, autoimmune diseases or conditions, inflammatory diseases or conditions, viral infections (eg, HIV infection of human patients), or diseases or conditions mediated or caused by viral infections.

The invention also relates to methods of modulating treatments or therapies in a subject organism (eg, a plant,

yeast, human or animal patient) by altering microbiota of the subject. Examples of therapies are adoptive cell therapy, antibody therapy (eg, immune checkpoint inhibition), radiation therapy, chemotherapy, eg, for treatment or prevention of a disease or condition in a patient.

In a first configuration the invention provides

A method of modulating a therapy of a disease or condition in a patient, the method comprising

- a. Carrying out the therapy in the patient; and
- b. Causing gut bacterial microbiota dysbiosis in the patient, whereby said dysbiosis modulates the therapy in the patient by modulating immune cells in the patient.

In another aspect, the first configuration the invention provides

A method of modulating a therapy of a disease or condition in a human or animal patient, the method comprising

- a. Carrying out the therapy in the patient; and
- b. Causing bacterial (eg, gut bacterial) microbiota dysbiosis in the patient, whereby said dysbiosis modulates the therapy in the patient by modulating immune cells in the patient; wherein the therapy comprises adoptive immune cell therapy (eg, adoptive T-cell therapy, eg, CAR-T cell administration to the patient).

In another aspect, the first configuration the invention provides

A method of modulating a therapy of a disease or condition in a human or animal patient, the method comprising

- a. Carrying out the therapy in the patient; and
- b. Causing bacterial (eg, gut bacterial) microbiota dysbiosis in the patient, whereby said dysbiosis modulates the therapy in the patient; wherein the therapy comprises administering an immune checkpoint inhibitor (eg, an anti-PD-L1, anti-PD-1, anti-CTLA4 or anti-TIM3 inhibitor, eg, an antibody) to the patient.

In another aspect, the first configuration the invention provides

A method of modulating a therapy of a disease or condition in a human or animal patient, the method comprising

- a. Carrying out the therapy in the patient; and
- b. Causing bacterial (eg, gut bacterial) microbiota dysbiosis in the patient, whereby said dysbiosis modulates the therapy in the patient; wherein the therapy comprises administering an antibody (eg, an anti-PD-L1, anti-PD-1, anti-CTLA4 or anti-TIM3 antibody; or an anti-TNF α superfamily member antibody, eg, an anti-TNF α , TNFR1 or BAFF antibody; or, an anti-IL6R or anti-IL-4Ra antibody; or an anti-PCSK9 antibody) to the patient.

In another aspect, the first configuration the invention provides

A method of modulating a treatment in a subject, the method comprising

- a. Carrying out the treatment in the subject; and
- b. Causing microbiota dysbiosis in the subject, whereby said dysbiosis modulates the treatment in the subject.

In an example, the subject or patient is a human. In an example, the subject or patient is a non-human animal. In an example, the subject is a plant, and optionally the treatment is a plant growth-promoting treatment, growth-inhibiting treatment, pesticide treatment, nitrogen fixing promotion treatment, herbicidal treatment or fertilizer treatment. In an example, the subject is a yeast, and optionally the treatment is a yeast growth-promoting treatment or growth-inhibiting treatment.

In an example, the modulating augments, upregulates, downregulates, inhibits, enhances or potentiates the treat-

5

ment or therapy of the subject or patient. In an example, the treatment or therapy is effective in the subject or patient, wherein the treatment or therapy is not effective or has reduced or increased efficacy in the subject, patient or a control subject or patient that has not been subject to the modulation. The control is of the same species as the subject or patient, and optionally the same age and/or sex. In an example, bacterial or archaeal host cells are killed or growth thereof is inhibited in the subject or patient using a method of an invention, wherein the control comprises cells of the same bacterial or archaeal species and the cells are not killed or growth inhibited by a method of the invention.

In an example, steps (a) and (b) are carried out simultaneously. In an example, step (a) is carried out before step (b). In an example, step (b) is carried out before step (a), and optionally step (b) is performed again after (a).

In an embodiment, the invention provides

A method of modulating a treatment in a plant or yeast, the method comprising

a. Carrying out the treatment in the plant or yeast; and
b. Causing bacterial microbiota dysbiosis in the plant or yeast, whereby said dysbiosis modulates the treatment in the subject;

wherein the treatment is a growth-promoting treatment, growth-inhibiting treatment, pesticide treatment, nitrogen fixing promotion treatment, herbicidal treatment or fertilizer treatment.

Causing microbial dysbiosis in the subject, patient, plant or yeast is, in an example comprises causing microbial dysbiosis on a surface of the subject, patient, plant or yeast, eg, on a leaf surface (when the subject is a plant) or on skin, lung, ocular or mucosal surface (when the subject or patient is a human or animal).

Instead of or additionally to causing bacterial dysbiosis, the invention comprises in step (b) causing archaeal microbiota dysbiosis in said subject, patient, plant or yeast.

For example, the disease or condition is an autoimmune disease or condition (eg, SLE) and the therapy is a treatment therefor, eg, administration of a tumor necrosis factor ligand superfamily member antagonist, eg, an anti-B-cell activating factor (BAFF) antibody, such as BENLYSTA® (belimumab) or a generic version thereof. For example, the disease or condition is an inflammatory disease or condition (eg, rheumatoid arthritis, IBD, Crohn's disease, colitis or psoriasis) and the therapy is a treatment therefor, eg, administration of sarilumab, dupilumab, a tumor necrosis factor ligand superfamily member antagonist, eg, an anti-TNF alpha antibody or trap, such as HUMIRA® (adalimumab), REMICADE® (infliximab), SIMPONI® (golimumab) or ENBREL® (etanercept) or a generic version thereof. For example, the disease or condition is a viral infection or mediated by a viral infection (eg, HIV infection) and the therapy is a treatment therefor, eg, administration of an anti-retroviral medicament or an anti-HIV vaccine. For example, the disease or condition is a cancer (eg, melanoma, NSCLC, breast cancer or pancreatic cancer) and the therapy is a treatment therefor, eg, administration of a chemotherapeutic agent, eg, a checkpoint inhibitor or agonist antibody such as an anti-CTLA4, PD-1, PD-L1, PD-L2, LAG3, OX40, CD28, BTLA, CD137, CD27, HVEM, KIR, TIM-3, VISTA, ICOS, GITR, TIGIT or SIRPa antibody. In an example, the antibody is a bispecific antibody that specifically binds first and second targets selected from CTLA4, PD-1, PD-L1, PD-L2, LAG3, OX40, CD28, BTLA, CD137, CD27, HVEM, KIR, TIM-3, VISTA, ICOS, GITR, TIGIT and SIRPa, eg, wherein the first target is CTLA4 and the second target is LAG3 or PD-1. Optionally, the antibody is

6

a human gamma-1 antibody and/or may be enhanced for ADCC or CDC. For example, the therapy is a vaccine therapy, eg, a cancer vaccine therapy or a vaccine therapy for treating or preventing an infection or infectious disease, such as malaria, HIV infection, *tuberculosis* infection, *cholera*, *Salmonella typhimurium* infection, *C difficile* infection, *Bordetella pertussis* infection or *Chlamydia* infection.

An embodiment of the first configuration provides

A method of modulating a cell therapy of a disease or condition in a patient, the method comprising

a. Carrying out cell therapy in the patient, comprising administering a population of cells to the patient, wherein administration of said cells is capable of treating the disease or condition in the patient; and
b. Causing gut bacterial microbiota dysbiosis in the patient, whereby said dysbiosis modulates the cell therapy in the patient.

In an example the cell therapy is an adoptive immune cell therapy, such as CAR-T or TILs therapy for the treatment of a cancer.

In a second configuration the invention provides

A method of treating or reducing the risk of a disease or condition in a patient, wherein the disease or condition is mediated by immune cells (eg, T-cells) in the patient, the method comprising causing gut bacterial microbiota dysbiosis in the patient, whereby said dysbiosis modulates immune cells (eg, T_H17 cells) in the patient, thereby treating or reducing the risk of said disease or condition in the patient.

For example, the disease or condition is an autoimmune disease or condition (eg, SLE), an inflammatory disease or condition (eg, rheumatoid arthritis, IBD, Crohn's disease, colitis or psoriasis), a viral infection or mediated by a viral infection (eg, HIV infection).

In an example, microbiota dysbiosis is effected by killing one or more target bacterial species in the microbiota or inhibiting growth of a population of said bacteria in the microbiota. In an example, microbiota dysbiosis is effected by killing one or more target archaeal species in the microbiota or inhibiting growth of a population of said archaea in the microbiota.

In a third configuration the invention provides

A method of modulating an adoptive immune cell therapy of a disease or condition in a patient, the method comprising

a. Carrying out adoptive immune cell therapy in the patient, comprising administering a population of immune cells to the patient, wherein administration of said immune cells is capable of treating the disease or condition in the patient; and
b. Altering the relative proportion of a sub-population of cells of a first bacterial species or strain, or archaeal species or strain, in a microbiota (eg, gut microbiota) of the patient, thereby producing an altered microbiota that modulates the immune cell therapy in the patient.

In another aspect, the third configuration the invention provides

A method of modulating a therapy of a disease or condition in a human or animal patient, the method comprising

a. Carrying out the therapy in the patient; and
b. Altering the relative proportion of a sub-population of cells of a first bacterial species or strain, or archaeal species or strain, in a microbiota (eg, gut microbiota) of the patient, thereby producing an altered microbiota that modulates the therapy in the patient;

wherein the therapy comprises administering an immune checkpoint inhibitor (eg, an anti-PD-L1, anti-PD-1, anti-CTLA4 or anti-TIM3 inhibitor, eg, an antibody) to the patient.

In another aspect, the third configuration the invention provides

A method of modulating a therapy of a disease or condition in a human or animal patient, the method comprising

- a. Carrying out the therapy in the patient; and
- b. Altering the relative proportion of a sub-population of cells of a first bacterial species or strain, or archaeal species or strain, in a microbiota (eg, gut microbiota) of the patient, thereby producing an altered microbiota that modulates the therapy in the patient;

wherein the therapy comprises administering an antibody (eg, an anti-PD-L1, anti-PD-1, anti-CTLA4 or anti-TIM3 antibody; or an anti-TNF α superfamily member antibody, eg, an anti-TNF α , TNFR1 or BAFF antibody; or, an anti-IL6R or anti-IL-4Ra antibody; or an anti-PCSK9 antibody) to the patient.

In another aspect, the third configuration the invention provides

A method of modulating a treatment in a subject, the method comprising

- a. Carrying out the treatment in the subject; and
- b. Altering the relative proportion of a sub-population of cells of a first bacterial species or strain, or archaeal species or strain, in a microbiota of the subject, whereby said dysbiosis modulates the treatment in the subject.

In an example, the subject or patient is a human. In an example, the subject or patient is a non-human animal. In an example, the subject is a plant, and optionally the treatment is a plant growth-promoting treatment, growth-inhibiting treatment, pesticide treatment, nitrogen fixing promotion treatment, herbicidal treatment or fertilizer treatment. In an example, the subject is a yeast, and optionally the treatment is a yeast growth-promoting treatment or growth-inhibiting treatment.

In an example, the modulating augments, upregulates, downregulates, inhibits, enhances or potentiates the treatment or therapy of the subject or patient. In an example, the treatment or therapy is effective in the subject or patient, wherein the treatment or therapy is not effective or has reduced or increased efficacy in the subject, patient or a control subject or patient that has not been subject to the modulation. The control is of the same species as the subject or patient, and optionally the same age and/or sex. In an example, bacterial or archaeal host cells are killed or growth thereof is inhibited in the subject or patient using a method of an invention, wherein the control comprises cells of the same bacterial or archaeal species and the cells are not killed or growth inhibited by a method of the invention.

In an example, steps (a) and (b) are carried out simultaneously. In an example, step (a) is carried out before step (b). In an example, step (b) is carried out before step (a), and optionally step (b) is performed again after (a).

In an embodiment, the invention provides

A method of modulating a treatment in a plant or yeast, the method comprising

- a. Carrying out the treatment in the plant or yeast; and
- b. Altering the relative proportion of a sub-population of cells of a first bacterial species or strain, or archaeal species or strain, in a microbiota of the plant or yeast, whereby said dysbiosis modulates the treatment in the plant or yeast; wherein the treatment is a growth-promoting treatment, growth-inhibiting treatment, pesticide treatment, nitrogen fixing promotion treatment, herbicidal treatment or fertilizer treatment.

Said altering of the relative proportion of sub-population of cells in the subject, patient, plant or yeast is, in an example comprises causing microbial dysbiosis on a surface

of the subject, patient, plant or yeast, eg, on a leaf surface (when the subject is a plant) or on skin, lung, ocular or mucosal surface (when the subject or patient is a human or animal).

The proportion of the first bacteria or archaea sub-population is increased or decreased. In an example, the relative ratio of first and second bacterial species or strains is altered (eg, increased or decreased); or the relative ratio of first and second archaeal species or strains is altered (eg, increased or decreased).

In an example, the adoptive immune cell therapy is CAR-T therapy for the treatment of a cancer. In an example, the adoptive immune cell therapy is a TILs therapy for the treatment of a cancer.

In an example of the first or third configuration, the cells of step (a) are of a first type selected from the group consisting of CD4⁺ T-cells, CD8⁺ T-cells, T_H1 cells or T_H17 cells and step (b) upregulates cells of that type in the patient. This is useful for enhancing the cell based therapy. In another example the cells of step (a) are of a first type selected from the group consisting of CD4⁺ T-cells, CD8⁺ T-cells, T_H1 cells or T_H17 cells and step (b) downregulates cells of that type in the patient. This is useful for dampening down the cell based therapy or a side effect thereof (eg, CRS).

In an embodiment, the dysbiosis or step (b) is carried out using selective targeting of a bacterial or archaeal microbiota sub-population using CRISPR/Cas targeting of microbiota (eg, gut microbiota) bacteria and/or archaea. In an example, the method comprises using guided nuclease (eg RNA-guided nuclease) cutting of a respective target sequence in host cells to modify the target sequences, whereby host cells are killed or the host cell population growth is reduced, thereby reducing the proportion of said sub-population in the microbiota. Suitable systems for carrying out the guided nuclease cutting are, for example, engineered CRISPR/Cas systems, TALENs, meganucleases and zinc finger systems.

To this end, the inventors believe that they have demonstrated for the first time inhibition of population growth of a specific bacterial strain in a mixed consortium of bacteria that naturally occur together in gut microbiota with one or more of the following features:—

- Population growth inhibition using an engineered CRISPR/Cas system by
 - targeting wild-type cells;
 - harnessing of wild-type endogenous Cas nuclease activity;
 - targeting essential and antibiotic resistance genes; wherein the targets are wild-type sequences.

The inventors have demonstrated this in a mixed population of human gut microbiota bacteria with the following features:—

- targeting bacterial growth inhibition in a mixed population of human gut microbiota species; wherein the population comprises three different species; comprising selective killing of one of those species and sparing cells of the other species;
- targeting cell growth inhibition in the presence of a phylogenetically-close other human gut microbiota species, which is spared such inhibition;
- targeting cell growth inhibition in a mixed population of human gut microbiota bacteria comprising target *Firmicutes* species and non-*Firmicutes* species;
- targeting cell growth inhibition of a specific *Firmicutes* species whilst sparing a different *Firmicutes* species in a mixed population of human gut microbiota bacteria;

targeting cell growth inhibition of a specific gram positive bacterial strain whilst sparing a different gram positive bacterial species in a mixed population of human gut microbiota bacteria;
 targeting a human gut microbiota bacterial species whilst sparing a commensal human gut bacterial species;
 targeting a human gut microbiota bacterial species whilst sparing a probiotic human gut bacterial species;
 targeting cell growth inhibition in a mixed population of human gut microbiota bacteria on a surface;
 achieving at least a 10-fold growth inhibition of a specific bacterial species alone or when mixed with a plurality of other bacterial species in a consortium of human gut microbiota bacteria; and
 achieving at least a 10-fold growth inhibition of two different strains of a specific human gut microbiota bacterial species.

The invention provides:

An ex vivo population of immune cells for use in a method of adoptive cell therapy of a patient for treating or preventing a disease or condition in the patient, the method comprising

- a. Carrying out adoptive immune cell therapy in the patient, comprising administering cells of said population to the patient, wherein administration of said immune cells is capable of treating the disease or condition in the patient; and
- b. Causing gut bacterial microbiota dysbiosis in the patient, whereby said dysbiosis modulates the immune cell therapy in the patient and said disease or condition is treated or prevented.

The invention provides

An ex vivo population of immune cells for use in a method of adoptive cell therapy of a patient for treating or preventing a disease or condition in the patient, the method comprising

- a. Carrying out adoptive immune cell therapy in the patient, comprising administering cells of said population to the patient, wherein administration of said immune cells is capable of treating the disease or condition in the patient; and
- b. Altering the relative proportion of a sub-population of cells of a first bacterial species or strain, or archaeal species or strain, in the gut microbiota of the patient, thereby producing an altered gut microbiota that modulates the immune cell therapy in the patient. The invention also provides CRISPR/Cas systems, arrays, cRNAs and kits for carrying out a method of the invention.

The invention also relates to systems, kits and other means for effecting the method.

Any features on one configuration herein are, in an example, combined with a different configuration of the invention for possible inclusion of such combination in one or more claims herein.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows a xylose inducible system.

FIG. 2 shows a ST1-CRISPR array.

FIG. 3 shows a spot assay on TH-agar of the strains used in this work. All strains were grown on TH-agar at 37° C. for 20 hours. Serial dilutions of overnight cultures were done in duplicate for *E. coli*, *L. lactis* and *S. mutans*, and triplicate for both strains of *S. thermophilus* in order to count individual colonies.

FIG. 4 shows selective growth of *S. thermophilus*, *S. mutans*, *L. lactis* and *E. coli* under different culture conditions. Tetracycline cannot be used to selectively grow *S. thermophilus* LMD-9. However, 3 g l⁻¹ of PEA proved to selectively grow *S. thermophilus* LMD-9 while limiting growth of *E. coli*.

FIG. 5 illustrates construction of two xylose induction cassettes.

FIG. 6 demonstrated characterization of the xylose inducible cassette in *Streptococcus thermophilus* LMD-9 with the plasmid pBAV1KT5-XylR-mCherry-Pldha. A clear response in fluorescence can be observed with increasing amount of xylose.

FIG. 7 illustrates the design of CRISPR array in pBAV1KT5-XylR-mCherry-P_{ldha+XylA}. The array contains 2 spacer sequences that target *S. thermophilus* genes under an inducible xylose promoter and a tracrRNA under a strong constitutive promoter P_{3A}.

FIG. 8 shows transformation efficiency of *Streptococcus thermophilus* LMD-9 with the plasmid pBAV1KT5-XylR-CRISPR-P_{ldha+XylA} and with pBAV1KT5-XylR-CRISPR-P_{XylA}.

FIG. 9 shows a schematic of the xylose-inducible CRISPR device. Upon induction of xylose the CRISPR array targeting both *polIII* and *tetA* on the *S. thermophilus* LMD-9 genome are expressed. Together with the constitutively expressed tracrRNA a complex is formed with Cas9. This complex will introduce a double stranded break in the *tetA* and *polIII* genes in the *S. thermophilus* LMD-9 genome resulting in limited cell viability.

FIG. 10 shows growth inhibition of *Streptococcus thermophilus* DSM 20617(T) with the plasmid pBAV1KT5-XylR-CRISPR-PXylA or pBAV1KT5-XylR-CRISPR-Pldha+XylA, not induced and induced. Picture taken after 63H of incubation. Colony counts in bottom left corner (top row: >1000, >1000, bottom row: 336, 113).

FIG. 11 shows a maximum-likelihood phylogenetic tree of 16S sequences from *S. thermophilus*, *L. lactis* and *E. coli*.

FIG. 12 shows the selective *S. thermophilus* growth inhibition in a co-culture of *E. coli*, *L. lactis* and *S. thermophilus* harboring either the pBAV1KT5-XylR-CRISPR-PxylA or the pBAV1KT5-XylR-CRISPR-PldhA+XylA plasmid. No growth difference is observed between *E. coli* harboring the pBAV1KT5-XylR-CRISPR-PxylA or the pBAV1KT5-XylR-CRISPR-PldhA+XylA plasmid. However, *S. thermophilus* (selectively grown on TH agar supplemented with 2.5 g l⁻¹ PEA) shows a decrease in transformation efficiency between the pBAV1KT5-XylR-CRISPR-PxylA (strong) or the pBAV1KT5-XylR-CRISPR-PldhA+XylA (weak) plasmid as we expected. We thus demonstrated a selective growth inhibition of the target *S. thermophilus* sub-population in the mixed population of cells. Colony counts in bottom left corner (top row: >1000, >1000, 68, bottom row: >1000, >1000, 32).

FIG. 13 shows regulators controlling the expression of spCas9 and the self-targeting sgRNA targeting the ribosomal RNA subunit 16s.

FIG. 14 shows specific targeting of *E. coli* strain by an exogenous CRISPR-Cas system. The sgRNA target the genome of K-12 derived *E. coli* strains, like *E. coli* TOP10, while the other strain tested was unaffected.

FIG. 15 shows spot assay with serial dilutions of individual bacterial species used in this study and mixed culture in TH agar without induction of CRISPR-Cas9 system.

FIG. 16 shows spot assay of the dilution 10³ on different selective media. TH with 2.5 g l⁻¹ PEA is a selective media for *B. subtilis* alone. MacConkey supplemented with malt-

ose is a selective and differential culture medium for bacteria designed to selectively isolate Gram-negative and enteric bacilli and differentiate them based on maltose fermentation. Therefore TOP10 ΔmalK mutant makes white colonies on the plates while Nissle makes pink colonies; A is *E coli* ΔmalK, B is *E coli* Nissle, C is *B subtilis*, D is *L lactis*, E is mixed culture; the images at MacConkey-/B and E appear pink; the images at MacConkey+/B and E appear pink.

FIG. 17 shows selective growth of the bacteria used in this study on different media and selective plates.

DETAILED DESCRIPTION

In the worked Example below, growth inhibition was addressed in a mixed population of human gut microbiota bacterial species. A >10-fold population growth inhibition in a selectively targeted species (a gram positive *Firmicutes* population) was achieved, sparing non-targeted commensal bacteria in the consortium. The inventors have realised the useful application of this for altering microbiota, such as gut microbiota, in situ in patients, thereby enabling immune cell modulation in the patient in response to the altered microbiota. The inventors also realised application to modulating treatments in subjects such as plants and yeast that comprise microbiota that can be altered. The inventors furthermore realised the utility for modulating immune cell-based therapies in patients or for treating or preventing immune cell-mediated diseases or conditions in patients, such as autoimmune diseases, inflammatory diseases and viral infections (eg, HIV infection of humans). The inventors realised the utility of causing dysbiosis of gut, skin, vaginal, nasal, ocular, lung, GI tract, rectal, scrotal, ear, skin or hair microbiota for effecting such modulation in a human or animal subject or patient.

As used herein “dysbiosis” refers to a change of the bacterial and/or archaeal balance of the microbiota, eg, gut microbiota. Change is relative to the balance prior to (eg, immediately prior or no more than a day before) carrying out the method. The change can be one or more of (i) an increase in the proportion of a first species (bacterial or archaeal species) or strain in the microbiota (eg, gut microbiota) (eg, *B fragalis* or *thetaitomicron*); (ii) an increase in the relative proportion of first and second species (eg, *B fragalis* versus *C difficile*; or *S thermophilus* v *E coli* or *L lactis*), first and second strains of the same species, or first and second phyla which are different from each other (eg, Bacteriodes versus *Firmicutes*); (iii) an addition of a species or strain that was not comprised by the microbiota prior to the treatment method; (iv) a decrease in the proportion of a first species (bacterial or archaeal species) or strain in the microbiota (eg, *C difficile* or *S thermophilus*); (v) a decrease in the relative proportion of first and second species (eg, *B fragalis* versus *C difficile*; or *S thermophilus* v *E coli* or *L lactis*), first and second strains of the same species, or first and second phyla which are different from each other (eg, Bacteriodes versus *Firmicutes*); and (vi) a removal of a species or strain that was not comprised by the microbiota prior to the treatment method. Dysbiosis may be effected, for example, using one or more selective antibacterial agents (eg, CRISPR-based or other guided nucleases described herein) or by administering one or more bacterial and/or archaeal transplants to the patient or subject to alter the balance of the microbiota, eg, gut microbiota.

The impact of the immune system on microbiota composition is suggested by several immune deficiencies that alter microbial communities in ways that predispose to disease. For example, Garrett et al. studied mice that lack the

transcription factor T-bet (encoded by *Tbx21*), which governs inflammatory responses in cells of both the innate and the adaptive immune system (Cell. 2007 Oct. 5; 131(1):33-45. “Communicable ulcerative colitis induced by T-bet deficiency in the innate immune system”, Garrett W S et al.). When *Tbx21*^{-/-} mice were crossed onto *Rag2*^{-/-} mice, which lack adaptive immunity, the *Tbx21*^{-/-}/*Rag2*^{-/-} progeny developed ulcerative colitis in a microbiota-dependent manner. Remarkably, this colitis phenotype was transmissible to wild-type mice by adoptive transfer of the *Tbx21*^{-/-}/*Rag2*^{-/-} microbiota. This demonstrated that altered microbiota were sufficient to induce disease. Another example of immune-driven dysbiosis is seen in mice deficient for epithelial cell expression of the inflammasome component NLRP6. These mice develop an altered microbiota with increased abundance of members of the Bacteroidetes phylum associated with increased intestinal inflammatory cell recruitment and susceptibility to chemically-induced colitis.

It has become evident that individual commensal species influence the makeup of *Lamina propria* T lymphocyte subsets that have distinct effector functions. Homeostasis in the gut mucosa is maintained by a system of checks and balances between potentially pro-inflammatory cells, which include *T_H1* cells that produce interferon- γ , *T_H17* cells that produce IL-17a, IL-17f, and IL-22, diverse innate lymphoid cells with cytokine effector features resembling *T_H2* and *T_H17* cells, and anti-inflammatory *Foxp3*⁺ regulatory T cells (*T_{regs}*).

A particular application of the invention is found in the shaping of *T_H17* cell populations in patients. Such cells have been implicated in autoimmune and inflammatory disorders. These cells were described in: Harrington L E, Hatton R D, Mangan P R, et al., “Interleukin 17-producing CD41 effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages”, *Nat Immunol.* 2005; 6(11): 1123-1132; and Park H, Li Z, Yang X O, et al., “A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17”, *Nat Immunol.* 2005; 6(11):1133-1141. In the case of autoimmune disorders, *T_H17* cell over activation can cause an inappropriate amount of inflammation, like in the case of multiple sclerosis, rheumatoid arthritis, and psoriasis. *T_H17* cells have also been shown to be necessary for maintenance of mucosal immunity. *T_H17* cells may contribute to the development of late phase asthmatic response due to increases in gene expression relative to *T_{reg}* cells.

In HIV, the loss of *T_H17* cell populations can contribute to chronic infection. The depletion of *T_H17* cell populations in the intestine disrupts the intestinal barrier, increases levels of movement of bacteria out of the gut through microbial translocation, and contributes to chronic HIV infection and progression to AIDS. Microbial translocation results in bacteria moving from out of the gut lumen, into the *Lamina propria*, to the lymph nodes, and beyond into non-lymphatic tissues. It can cause the constant immune activation seen through the body in the late stages of HIV. Increasing *T_H17* cell populations in the intestine has been shown to be both an effective treatment as well as possibly preventative. Although all CD4⁺ T cells gut are severely depleted by HIV, the loss of intestinal *T_H17* cells in particular has been linked to symptoms of chronic, pathogenic HIV and SIV infection. Microbial translocation is a major factor that contributes to chronic inflammation and immune activation in the context of HIV. In non-pathogenic cases of SIV, microbial translocation is not observed. *T_H17* cells prevent severe HIV infection by maintaining the intestinal epithelial barrier

during HIV infection in the gut. Because of their high levels of CCR5 expression, the coreceptor for HIV, they are preferentially infected and depleted. Thus, it is through T_H17 cell depletion that microbial translocation occurs. Additionally, the loss of T_H17 cells in the intestine leads to a loss of balance between inflammatory T_H17 cells and T_{reg} cells, their anti-inflammatory counterparts. Because of their immunosuppressive properties, they are thought to decrease the anti-viral response to HIV, contributing to pathogenesis. There is more T_{reg} activity compared to T_H17 activity, and the immune response to the virus is less aggressive and effective. Revitalizing T_H17 cells has been shown to decrease symptoms of chronic infection, including decreased inflammation, and results in improved responses to highly active anti-retroviral treatment (HAART). This is an important finding—microbial translocation generally results in unresponsiveness to HAART. Patients continue to exhibit symptoms and do not show as reduced a viral load as expected. In an SIV-rhesus monkey model, it was found that administering IL-21, a cytokine shown to encourage T_H17 differentiation and proliferation, decreases microbial translocation by increasing T_H17 cell populations.

In an example of the method, IL-21, IL-15 and/or IL-2 is administered to the patient sequentially or simultaneously with the cell population. This is useful for further modulating immune cell populations in the patient.

Yang et al. observed that the presence of T_H17 cells in mice requires colonisation of mice with microbiota. Segmented filamentous bacteria (SFB) were sufficient to induce T_H17 cells and promote T_H17 -dependent autoimmune disease in animal models (Nature, 2014 Jun. 5; 510(7503):152-6. doi: 10.1038/nature13279. Epub 2014 Apr. 13, “Focused specificity of intestinal Th17 cells towards commensal bacterial antigens”, Yang Y et al.). SFB appear able to penetrate the mucus layer overlying the intestinal epithelial cells in the terminal ileum, and they interact closely with the epithelial cells, inducing host cell actin polymerization at the site of interaction and, presumably, signaling events that result in a T_H17 polarizing environment within the *Lamina propria*.

In an example, the first bacteria are of a species or strain comprising a 16s rDNA sequence that is at least 80, 85, 90, 95, 96, 97, 98 or 99% identical to a 16s rDNA sequence of a segmented filamentous bacterium. In an embodiment, the method increases the proportion of the first bacteria, wherein T_H17 cells in the patient are upregulated, eg, wherein the disease is a cancer or a viral infection (eg, HIV). In an embodiment, the method decreases the proportion of the first bacteria, wherein T_H17 cells in the patient are downregulated, eg, wherein the disease or condition is an autoimmune or inflammatory disease or condition, or for reducing the risk of CRS in a cancer patient receiving ACT.

In an example, the method treats or prevents an allergic disease or condition, eg, asthma. In an example, the method treats or prevents an IgE-mediated disease or condition, eg, asthma.

In an example, the method reduces autotoxicity in the patient mediated by T_H2 cell cytokine release.

López et al. observed that intestinal dysbiosis, characterised by a reduced *Firmicutes*/Bacteroidetes ratio, has been reported in systemic lupus erythematosus (SLE) patients. In their study, in vitro cultures revealed that microbiota isolated from SLE patient stool samples (SLE-M) promoted lymphocyte activation and T_H17 differentiation from naive $CD4^+$ lymphocytes to a greater extent than healthy control microbiota. Enrichment of SLE-M with T_{reg} -inducing bacteria showed that a mixture of two Clostridia strains significantly reduced the T_H17/T_H1 balance, whereas *Bifidobac-*

terium bifidum supplementation prevented $CD4^+$ lymphocyte over-activation. Ex vivo analyses of patient samples showed enlarged T_H17 and Foxp3⁺ IL-17⁺ populations, suggesting a possible T_{reg} - T_H17 trans-differentiation. Moreover, analyses of faecal microbiota revealed a negative correlation between IL-17⁺ populations and *Firmicutes* in healthy controls, whereas in SLE this phylum correlated directly with serum levels of IFN γ , a T_H1 cytokine slightly reduced in patients. (Sci Rep. 2016 Apr. 5; 6:24072. doi: 10.1038/srep24072, “Th17 responses and natural IgM antibodies are related to gut microbiota composition in systemic lupus erythematosus patients”, López P et al.).

Other bacteria have been shown to enhance the anti-inflammatory branches of the adaptive immune system by directing the differentiation of T_{regs} or by inducing IL-10 expression. For example, colonisation of gnotobiotic mice with a complex cocktail of 46 mouse Clostridial strains, originally isolated from mouse faeces and belonging mainly to cluster IV and XIVa of the *Clostridium* genus, results in the expansion of *Lamina propria* and systemic T_{regs} .

Bacteroides fragilis polysaccharide-A (PSA) impacts the development of systemic T cell responses. Colonization of germ-free mice with PSA-producing *B. fragilis* results in higher numbers of circulating $CD4^+$ T cells as compared to mice colonized with *B. fragilis* lacking PSA. PSA-producing *B. fragilis* also elicits higher T_H1 cell frequencies in the circulation. Together, these findings show that commensal bacteria have a general impact on immunity that reaches well beyond mucosal tissues.

The decrease in *F. prausnitzii* found in IBD patients is of interest because this bacteria is butyrate-producing, and its oral administration reduces the severity of TNBS-induced colitis in mice. In an example, the first species is a butyrate-producing bacterial species (eg, *F. prausnitzii*) and the proportion of the first species in the microbiota is reduced, wherein the method downregulates T-effector and/or T-helper cells in the patient, thereby treating or preventing said disease or condition (eg, an autoimmune or inflammatory disease or condition or CRS).

Archaea have traditionally been divided into five phyla, namely Crenarchaeota, Euryarchaeota, Korarchaeota, Nanoarchaeota and Thaumarchaeota. Based on the increasing wealth of whole genome data (mainly from environmental isolates), the archaeal phylogeny has been revisited recently: the four groups Korarchaeota, Crenarchaeota, Thaumarchaeota and the newly proposed Aigarchaeota have been comprised into one superphylum (the so-called TACK-superphylum) to the exclusion of Euryarchaeota and Nanoarchaeota. The first species in the method of the invention can be any of the archaea mentioned in this paragraph.

T cells mature in the thymus, express TCR (T cell receptor), and can express either CD8 glycoprotein on their surface and are called $CD8^+$ T cells (cytotoxic) or $CD4$ glycoprotein and are then called $CD4$ cells (helper T cells). $CD4^+$ cells differentiate into different subsets: T_H (T helper) 1, T_H 2, T_H 9, T_H17 , T_H22 , T_{reg} (regulatory T cells) and T_{fh} (follicular helper T cells), which are characterized by different cytokine profiles. These different $CD4^+$ subsets play a critical role in the immune and effector response functions of T cells. All $CD4^+$ T_H subsets are differentiated from naive $CD4^+$ T cells by specific cytokines: T_H1 by IL-12 and IFN- γ (pro-inflammatory cytokine, with multiple roles such as increase of TLR (Toll-like receptor), induction of cytokine secretion or macrophage activation); T_H2 by IL-4; T_{reg} by IL-2 and TGF-beta. Each T_H subset releases specific cytokines that can have either pro- or anti-inflammatory func-

tions, survival or protective functions. For example, T_H1 releases IFN- γ and TNF; T_H2 releases IL-4 (an important survival factor for B-type lymphocytes), IL-5 and IL-13; T_H9 produces IL-9; T_{reg} secretes IL-10 (a cytokine with an immunosuppressive function, maintaining expression of FOXP3 transcription factor needed for suppressive function of T_{reg} on other cells) and TGF- β ; T_H17 produces IL-17 (a cytokine playing an important role in host defense against bacteria, and fungi).

An embodiment of the invention finds application for modulating CAR-T and other adoptive immune-cell therapies (such as adoptive TILs therapy). Several reports have demonstrated differential roles of different types of cytokines released by CD4+ subsets, an important consideration when assessing CAR-T and other immune cell-based therapies. T_H1 and T_H2 CD4+ T cell subset cytokines were shown to drive different types of cytotoxicity generated by second generation CD28-containing CAR-T. Short-term toxicity was observed with high levels of T_H1 cytokines, while high doses of T_H2 type cytokines generated chronic autotoxicity in animals that received second generation CD19-specific CAR-T. CAR-T cells engineered to deliver inducible IL-12 modulated tumor stroma to destroy cancer. IL-12 release by engineered CAR-T cells increased anti-cancer activity by recruiting macrophages. IL-12 released by CAR-T also induced reprogramming of suppressive cells, reversing their inhibitory functions suggesting its evaluation in clinical trials. The persistence of CAR-T therapy was shown to be dependent on the number of CD4+ cells and the number of central memory cells in the infused product. CD8+ clones isolated from central memory T cells but not from CD8+ effector cells persisted long-term in vivo during adoptive T cell transfer in a nonhuman primate model, indicating the importance of specific T cell subset functions for effective adoptive immunotherapy. It has also been shown that the combination of CD8+ subset with CD4+ subset significantly enhanced T cell adoptive transfer. CD4+ cells were shown to support development of CD8+ memory functions, demonstrating the importance of both subsets and combinations in immunotherapy trials. Several preclinical models demonstrated the advantage of different T cell subsets for effective CAR-T therapy: CD8+ CD45RA+ CCR7+ CAR-T cells with closest to the T-memory stem cells phenotype cells produced greater anti-tumor activity of CAR-T cells; both CD8+ and CD4+ subsets expressed synergistic anti-tumor CAR-T activities.

In an example, the administered cell population is a population of CAR-T cells comprising a combination of a CD8+ CAR-T subset with CD4+ CAR-T subset.

In an example of the invention, the cell therapy is an adoptive T-cell therapy and optionally cells selected from the group consisting of CD4+ T-cells, CD8+ T-cells, TH1 cells and TH17 cells are administered to the patient. In an example, cell therapy is enhanced by the method of the invention, eg, immune cell cytotoxicity of cancer cells is enhanced in the patient, or treatment of the disease or condition is enhanced. In an example, cell therapy is reduced by the method of the invention, eg, immune cell cytotoxicity of cancer cells is reduced in the patient, or the risk of CRS is reduced (eg, in a cancer patient). Thus, in an embodiment the method reduces or prevents the risk of cytokine release syndrome (CRS) in the patient. In an embodiment the method reduces or prevents the risk of an unwanted side-effect of the cell therapy (eg, a CAR-T therapy side effect in a human patient, such as CRS).

In an example, the immune cell population comprises CAR-T cells and/or T-cells expressing engineered T-cell

receptors (TCRs) and/or tumour infiltrating lymphocytes (TILs, eg, Engineered TILs). WO2013063361, U.S. Pat. No. 9,113,616, US20130109053, US20160081314 and WO2016044745 (whose disclosures are incorporated herein by reference) describe suitable transgenic in vivo platforms for generating CARs and TCRs for use in generating cells for use in the present invention. The immune cell population may comprise engineered autologous or allogeneic immune cells (transplant), eg, T-cells, NK cells and/or TILs, eg, wherein the cells and patient are human. A benefit of autologous cells is that the modulation of the endogenous system is likely to be tuned similarly to modulation of the cell transplanted autologous cells. In an embodiment, the administered cells and patient are of the same species or strain, for example, human or rodent (eg, mouse), for example, HLA or MHC matched donor transplant and recipient patient.

In an example, the T-cells are CD4+ T-cells or T_H17 cells. For example, the administered CAR-T cells comprise a chimaeric antigen receptor comprising an ICOS intracellular domain and optionally the cells are T_H17 cells. In an embodiment, the administered T-cells are CD8+ $CD45RA^+$ CCR7+ CAR-T cells.

Adoptive transfer experiments in mice indicate that T_H17 cells have higher in vivo survival and self-renewal capacity than T_H1 polarized cells. In an example, therefore, T_H17 cells are modulated in the patient, eg, upregulated, eg, expanded in the patient, or downregulated. These may be endogenous T-cells of the patient and/or cells that have been administered to the patient or progeny thereof. In an embodiment, ROR γ t-expressing T_H17 cells are upregulated, eg, expanded in the patient. In an embodiment expression of one or more T_H17 -related genes is increased, eg, one or more of Rorc, Il22 and Il26. In an embodiment expression of one or more T_H1 -related genes is increased, eg, one or more of IFng, Tnfa and Tbx21 (T-bet). In an embodiment, in this case the disease or condition is a cancer.

In an example, T_{reg} cells are modulated in the patient, eg, upregulated, eg, expanded in the patient, or downregulated. These may be endogenous T-cells of the patient and/or cells that have been administered to the patient or progeny thereof. In an embodiment, in this case the disease or condition is an autoimmune, inflammatory or infectious disease or condition when the T_{reg} cells are upregulated.

In an example, CD4+ cells are modulated in the patient, eg, upregulated, eg, expanded in the patient, or downregulated. These may be endogenous cells of the patient and/or cells that have been administered to the patient or progeny thereof.

In an example, CD8+ cells are modulated in the patient, eg, upregulated, eg, expanded in the patient, or downregulated. These may be endogenous cells of the patient and/or cells that have been administered to the patient or progeny thereof.

In an example, tumour infiltrating lymphocytes (TILs) are modulated in the patient, eg, upregulated, eg, expanded in the patient, or downregulated. These may be endogenous cells of the patient and/or cells that have been administered to the patient or progeny thereof.

In an example, memory cells, such as one or more of central memory T cells (T_{CM}), effector memory T cells (T_{EM}), stem cell memory cells (T_{SCM}) and effector cells (T_{eff}), are upregulated in the microbiota or patient, optionally wherein the cells are comprised by the immune cell population administered to the patient and/or are progeny thereof. In an embodiment, the memory cells are CD45RO+CD62L+ or CD25+ CD45RA- CD45RO+ CD127+.

Upregulation of a cell population may, for example, be an increase in the population size or proportion of cells of that type (eg, species or strain) in the microbiota or patient or subject and/or an increase in the activity (eg, cytotoxicity, effector function or suppressor function) of cells of that type in the microbiota or patient or subject. Downregulation of a cell population may, for example, be an decrease in the population size or proportion of cells of that type (eg, species or strain) in the microbiota or patient or subject and/or a decrease in the activity (eg, cytotoxicity, effector function or suppressor function) of cells of that type in the microbiota or patient or subject.

In an example, the cell therapy population comprises CAR-T cells (ie, respectively T-cells engineered to surface-express chimaeric antigen receptors (CARs). Alternatively, the cells are CAR-TIL or CAR-NK cells. A CAR comprises an extracellular receptor domain for binding to a target antigen (eg, a tumour cell antigen), a transmembrane moiety and an intracellular moiety comprising one or more (eg, first and second) signalling domains for signalling in the immune cell (eg, T-cell). Examples of suitable intracellular domains are well known, eg, a combination of a CD3 ζ domain and one or more of an ICOS, CD28, OX40 or 4-1BB signalling domain, eg, a combination of an ICOS and CD28; or ICOS and 41-BB; CD28 and 41-BB signalling domain.

Optionally, the cell population is comprised by a transplant that is administered to the patient to treat or prevent a disease (eg, a cancer, autoimmune disease, transplant rejection or GvHD) or the cell or transplant is for such use.

In an example, the patient is a human, eg, is a woman; or a man.

In an example, the patient or human has undergone lymphodepletion before administration of the immune cell (eg, CAR-T cell).

Techniques for producing CARs and CAR T-cells are known and routine in the art, and these can be generally applied to producing cells for use in the invention (eg, see WO2012079000A1; U.S. Pat. Nos. 8,906,682, 8,911,993, 8,916,381, 8,975,071, 9,101,584, 9,102,760, 9,102,761, 9,328,156, 9,464,140, 9,481,728, 9,499,629, 9,518,123, 9,540,445, US20130287748, US20130288368, US20130309258, US20140106449, US20140370017, US20150050729, US20150093822, US20150099299, US20150118202, US20160130355, US20160159907, US20160194404, US20160208012; J Immunother. 2009 September; 32(7): 689-702, doi: 10.1097/CJI.0b013e3181ac6138, "Construction and Pre-clinical Evaluation of an Anti-CD19 Chimeric Antigen Receptor", James N. Kochenderfer et al; also WO2014012001 and US20150290244 for general methods applicable to the present invention). For example, use of electroporation, retroviral vectors or lentiviral vectors—as will be known by the skilled addressee—can be used to introduce nucleotide sequences encoding elements of the CAR into T-cells, NK cells, TILs or other immune cells to produce the CAR-cells. Cells isolated from the patient (autologous cell sample) or from another donor of the same species (allogeneic sample) can be used to provide ancestor cells that are genetically engineered to include the CAR-encoding sequences. Expansion of cells can be used in the process, as known in the art. For example, after engineering CAR-cells, the cell population can be massively expanded using routine techniques to produce a transplant that is administered (eg, transfused) into the patient. The patient can be a human or non-human animal Nucleotide sequences for one or more of the CAR elements (eg, for one or more of three signalling domains)

can be cloned or sequenced using a cell obtained from the patient or from another donor.

For example, the CAR comprises a first intracellular signalling domain, which is a human CD3 ζ domain and the cells administered to the patient are human cells comprising an endogenous nucleotide sequence encoding said human CD3 ζ domain. In an example, the CD3 zeta signaling domain comprises SEQ ID NO: 1, i.e., the amino acid sequence of SEQ ID NO: 24 as disclosed in WO2012079000A1, which sequence is explicitly incorporated herein for use in the present invention and possible inclusion in one or more claims herein. In an example, the CD3 zeta signaling domain is encoded by SEQ ID NO: 2, i.e., the nucleic acid sequence of SEQ ID NO: 18 as disclosed in WO2012079000A1, which sequence is explicitly incorporated herein for use in the present invention and possible inclusion in one or more claims herein.

For example, the first signalling domain is a human CD28 domain and the cell population of the invention is a population of human cells each comprising an endogenous nucleotide sequence encoding said human CD28 domain.

For example, the first signalling domain is a human 4-1BB domain and the cell population of the invention is a population of human cells each comprising an endogenous nucleotide sequence encoding said human 4-1BB domain.

For example, the first signalling domain is a human OX40 domain and the cell population of the invention is a population of human cells each comprising an endogenous nucleotide sequence encoding said human OX40 domain.

In an example, the first signalling domain is a CD3' domain, and the first and second intracellular signalling domains do not naturally occur together in a single cell (eg, a human wild-type cell or a cell isolated from the patient), eg, the second domain is a CD28, CD27, OX40 or 4-1BB domain.

In an example, the first intracellular domain is a CD3' domain, CD28 domain or 4-1BB domain.

In an example, the CAR is an engineered single polypeptide comprising (in N- to C-terminal direction) an antigen binding site (eg, an antibody scFv, which may be human); an optional hinge (eg, a human CD8 α hinge); a transmembrane domain (eg, a human CD8 α or CD28 transmembrane domain); and a human CD3' domain. In an example, the CAR is a complex of two or more of said polypeptides. Optionally, the CAR comprises a further intracellular signalling domain (i) between the transmembrane and CD3 ζ domains. Optionally, the CAR comprises a further intracellular signalling domain, wherein the CD3 ζ domain is between the further signaling domain and the transmembrane domain. In an example, the further signalling domain is a human CD27 domain, CD28 domain, ICOS domain, OX40 domain, CD40 domain, 4-1BB domain, a Fc ϵ R1 γ domain, CD64 domain or CD16 domain. In an alternative, instead of a single polypeptide, the CAR comprises an engineered complex of at least 2 polypeptides comprising said domains.

The immune cells may be administered either alone, or as a pharmaceutical composition in combination with diluents and/or with other components such as IL-2 or other cytokines or cell populations.

In an embodiment, the immune cells (eg, CAR cells or cells bearing TCRs) comprise cell surface binding sites (eg, provided by the CAR or TCR) that bind a TAA. Tumour antigens (TAA) are proteins that are produced by tumour cells that elicit an immune response, particularly T-cell mediated immune responses. The selection of the antigen binding specificity will depend on the particular type of

cancer to be treated. Tumour antigens are well known in the art and include in the context of an embodiment of the invention, for example, a glioma-associated antigen, carcinoembryonic antigen (CEA), β -human chorionic gonadotropin, alphafetoprotein (AFP), lectin-reactive AFP, thyroglobulin, RAGE-1, MN-CA IX, human telomerase reverse transcriptase, RU1, RU2 (AS), intestinal carboxy esterase, mut hsp70-2, M-CSF, prostase, prostate-specific antigen (PSA), PAP, NY-ESO-1, LAGE-1a, p53, prostein, PSMA, Her2/neu, survivin and telomerase, prostate-carcinoma tumour antigen-1 (PCTA-1), MAGE, ELF2M, neutrophil elastase, ephrinB2, CD22, insulin growth factor (IGF)-I, IGF-II, IGF-I receptor and mesothelin.

In one embodiment, the tumour antigen comprises one or more antigenic cancer epitopes associated with a malignant tumour. Malignant tumours express a number of proteins that can serve as target antigens for an immune attack. These molecules include but are not limited to tissue-specific antigens such as MART-1, tyrosinase and GP 100 in melanoma and prostatic acid phosphatase (PAP) and prostate-specific antigen (PSA) in prostate cancer. Other target molecules belong to the group of transformation-related molecules such as the oncogene HER-2/Neu ErbB-2. Yet another group of target antigens are onco-foetal antigens such as carcinoembryonic antigen (CEA). In B-cell lymphoma the tumour-specific idiotype immunoglobulin constitutes a truly tumour-specific immunoglobulin antigen that is unique to the individual tumour. B-cell differentiation antigens such as CD19, CD20 and CD37 are other candidates for target antigens in B-cell lymphoma. Some of these antigens (CEA, HER-2, CD19, CD20, idiotype) have been used as targets for passive immunotherapy with monoclonal antibodies with limited success. The first antigen or fourth binding moiety can be any of these TAAs or can be an antigenic sequence of any of these TAAs.

Non-limiting examples of TAA antigens in an embodiment of the invention include the following: Differentiation antigens such as MART-1/MelanA (MART-1), g 1 OO (Pmel 17), tyrosinase, TRP-1, TRP-2 and tumour-specific multi-lineage antigens such as MAGE-1, MAGE-3, BAGE, GAGE-1, GAGE-2, p 5; overexpressed embryonic antigens such as CEA; overexpressed oncogenes and mutated tumour-suppressor genes such as p53, Ras, HER-2/neu; unique tumour antigens resulting from chromosomal translocations; such as BCR-ABL, E2A-PRL, H4-RET, 1GH-IGK, MYL-RAR; and viral antigens, such as the Epstein Barr virus antigens EBVA and the human papillomavirus (HPV) antigens E6 and E7. Other large, protein-based antigens include TSP-180, MAGE-4, MAGE-5, MAGE-6, RAGE, NY-ESO, p1 85erbB2, p 1 80erbB-3, c-met, nm-23H1, PSA, TAG-72, CA 19-9, CA 72-4, CAM 17.1, NuMa, K-ras, beta-Catenin, CDK4, Mum-1, p 15, p 16, 43-9F, 5T4(791Tgp72} alpha-fetoprotein, beta-HCG, BCA225, BTAA, CA 125, CA 15-3\CA 27.29\BCAA, CA 195, CA 242, CA-50, CAM43, CD68\ I, CO-029, FGF-5, G250, Ga733VEpCAM, HTgp-175, M344, MA-50, MG7-AG, MOV 18, NB/70K, NY-CO-1, RCAS 1, SDCCAG16, TA-90\Mac-2 binding protein\Acyclophilin C-associated protein, TAAL6, TAG72, TLP, and TPS.

In one embodiment, the CAR or TCR comprises a binding site for human CD 19, eg, for a CAR this can be provided by an anti-CD 19 scFV, optionally wherein the anti-CD19 scFV is encoded by SEQ ID NO: 3, i.e., SEQ ID: 14 disclosed in WO2012079000A1. In one embodiment, the anti-CD 19 scFV comprises SEQ ID NO: 4, i.e., the amino acid sequence of SEQ ID NO: 20 disclosed in WO2012079000A1. The sequences in this paragraph appear

in WO2012079000A1 and are explicitly incorporated herein for use in the present invention and for possible inclusion in one or more claims herein.

In one embodiment, the transmembrane domain that naturally is associated with one of the domains in the CAR is used. In some instances, the transmembrane domain can be selected or modified by amino acid substitution to avoid binding of such domains to the transmembrane domains of the same or different surface membrane proteins to minimize interactions with other members of the receptor complex.

The transmembrane domain may be derived either from a natural or from a synthetic source. Where the source is natural, the domain may be derived from any membrane-bound or transmembrane protein. Transmembrane regions of particular use in this invention may be derived from (i.e. comprise at least the transmembrane region(s) of) the alpha, beta or zeta chain of the T-cell receptor, CD28, CD3 epsilon, CD45, CD4, CD5, CDS, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD 134, CD137 or CD 154. Alternatively the transmembrane domain may be synthetic, in which case it will comprise predominantly hydrophobic residues such as leucine and valine. Optionally, a triplet of phenylalanine, tryptophan and valine will be found at each end of a synthetic transmembrane domain.

Optionally, a short oligo- or polypeptide linker, preferably between 2 and 10 amino acids in length forms a linkage between the transmembrane domain and the intracellular part of the immune cell transmembrane protein, such as the CAR. A glycine-serine doublet provides a particularly suitable linker (eg, a (G₄S)_n linker as disclosed herein).

Optionally, the transmembrane domain is the CD8 transmembrane domain encoded by SEQ ID NO: 5, i.e., the nucleic acid sequence of SEQ ID NO: 16 disclosed in WO2012079000A1. In one embodiment, the CD8 transmembrane domain comprises SEQ ID NO: 6, i.e., the amino acid sequence of SEQ ID NO: 22 disclosed in WO2012079000A1. The sequences in this paragraph appear in WO2012079000A1 and are explicitly incorporated herein for use in the present invention and for possible inclusion in one or more claims herein.

In some instances, the transmembrane domain comprises the CD8 hinge domain encoded by SEQ ID NO: 7, i.e., the nucleic acid sequence of SEQ ID NO: 15 disclosed in WO2012079000A1. In one embodiment, the CD8 hinge domain comprises SEQ ID NO: 8, i.e., the amino acid sequence of SEQ ID NO: 21 disclosed in WO2012079000A1. The sequences in this paragraph appear in WO2012079000A1 and are explicitly incorporated herein for use in the present invention and for possible inclusion in one or more claims herein.

The intracellular part or otherwise the intracellular signaling domain(s) of the transmembrane protein expressed by cells of the cell population administered to the patient is responsible for activation of at least one of the normal effector functions of the immune cell that expresses the transmembrane protein (eg, a T-cell function, such as leading to cytotoxicity (for T-effector cells for example) or suppression (for T-regulatory cells)). The term "effector function" refers to a specialized function of a cell. Effector function of a T cell, for example, may be cytolytic activity or helper activity including the secretion of cytokines. Thus the term "intracellular signaling domain" refers to the portion of a protein which transduces the effector function signal and directs the cell to perform a specialized function. While usually the entire intracellular signaling domain can be employed, in many cases it is not necessary to use the entire chain. To the extent that a truncated portion of the

intracellular signaling domain is used, such truncated portion may be used in place of the intact chain as long as it transduces the effector function signal. The term "signaling domain" is thus meant to include any truncated portion of the intracellular signaling domain sufficient to transduce the effector function signal. Examples of intracellular signaling domains for use in the transmembrane protein of the administered cells include the cytoplasmic sequences of the T cell receptor (TCR) and co-receptors that act in concert to initiate signal transduction following antigen receptor engagement, as well as any derivative or variant of these sequences and any synthetic sequence that has the same functional capability.

It is known that signals generated through the TCR alone are insufficient for full activation of the T cell and that a secondary or co-stimulatory signal is also required. Thus, T cell activation can be said to be mediated by two distinct classes of cytoplasmic signaling sequence: those that initiate antigen-dependent primary activation through the TCR (primary cytoplasmic signaling domain) and those that act in an antigen-independent manner to provide a secondary or co-stimulatory signal (secondary cytoplasmic signaling domain). Primary cytoplasmic signaling sequences regulate primary activation of the TCR complex either in a stimulatory way, or in an inhibitory way. Primary cytoplasmic signaling sequences that act in a stimulatory manner may contain signaling motifs which are known as immunoreceptor tyrosine-based activation motifs or ITAMs.

In an example, the first signalling domain is a primary cytoplasmic signaling domain (eg, CD3 ζ domain) In an example, the first signalling domain is a secondary cytoplasmic signaling domain (eg, CD28 or 4-1BB domain).

In an example, the first signalling domain comprises one or more ITAMs.

Examples of suitable ITAM containing primary cytoplasmic signaling domains that are of particular use in the invention include those derived from TCR zeta, FcR gamma, FcR beta, CD3 gamma, CD3 delta, CD3 epsilon, CDS, CD22, CD79a, CD79b, and CD66d. It is particularly preferred that cytoplasmic signaling molecule in the transmembrane protein of the invention comprises a cytoplasmic signaling sequence derived from CD3 zeta.

The intracellular part optionally comprises (eg, as the first signalling domain or a further intracellular domain) a domain of a costimulatory molecule. A costimulatory molecule is a cell surface molecule other than an antigen receptor or their ligands that is required for an efficient response of lymphocytes (eg, T- or NK cells) to an antigen. Examples of such molecules include CD27, CD28, 4-1BB (CD 137), OX40, CD30, CD40, PD-1, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, B7-H3, and a ligand that specifically binds with CD83, and the like. Thus, these and other costimulatory elements are within the scope of the invention for use in the intracellular part of the transmembrane protein.

The intracellular moiety domains may be linked together by one or more linkers, eg, a (G₄S)_n linker as disclosed herein.

In one embodiment, the intracellular moiety comprises the signaling domain of CD3-zeta and the signaling domain of CD28. In another embodiment, the intracellular moiety comprises the signaling domain of CD3-zeta and the signaling domain of 4-1BB. In yet another embodiment, the intracellular moiety comprises the signaling domain of CD3-zeta and the signaling domain of CD28 and 4-1BB.

In one embodiment, the intracellular moiety comprises the signaling domain of 4-1BB and the signaling domain of

CD3-zeta, wherein the signaling domain of 4-1BB is encoded by SEQ ID NO: 9, i.e., the nucleic acid sequence set forth in SEQ ID NO: 17 disclosed in WO2012079000A1 and the signaling domain of CD3-zeta is encoded by SEQ ID NO: 2, i.e., the nucleic acid sequence set forth in SEQ ID NO: 18 disclosed in WO2012079000A1. The sequences in this paragraph appear in WO2012079000A1 and are explicitly incorporated herein for use in the present invention and for possible inclusion in one or more claims herein.

In one embodiment, the intracellular moiety comprises the signaling domain of 4-1BB and the signaling domain of CD3-zeta, wherein the signaling domain of 4-1BB comprises SEQ ID NO: 10, i.e., the amino acid sequence of SEQ ID NO: 23 disclosed in WO2012079000A1 and the signaling domain of CD3-zeta comprises SEQ ID NO: 1, i.e., the amino acid sequence of SEQ ID NO: 24 disclosed in WO2012079000A1. The sequences in this paragraph appear in WO2012079000A1 and are explicitly incorporated herein for use in the present invention and for possible inclusion in one or more claims herein.

In one embodiment, the intracellular moiety comprises the signaling domain of 4-1BB and the signaling domain of CD3-zeta, wherein the signaling domain of 4-1BB comprises SEQ ID NO: 10, i.e., the amino acid sequence set forth in SEQ ID NO: 23 as disclosed in WO2012079000A1 and the signaling domain of CD3-zeta comprises SEQ ID NO: 1, i.e., the amino acid sequence set forth in SEQ ID NO: 24 disclosed in WO2012079000A1. The sequences in this paragraph appear in WO2012079000A1 and are explicitly incorporated herein for use in the present invention and for possible inclusion in one or more claims herein.

Sources of T-cells and other immune cells are disclosed in WO2012079000A1, U.S. Pat. Nos. 8,906,682, 8,911,993, 8,916,381, 8,975,071, 9,101,584, 9,102,760, 9,102,761, 9,328,156, 9,464,140, 9,481,728, 9,499,629, 9,518,123, 9,540,445, U.S. 20130287748, US20130288368, US20130309258, US20140106449, US20140370017, US20150050729, US20150093822, US20150099299, US20150118202, US20160130355, US20160159907, US20160194404, US20160208012, as well as methods of generating, activating and expanding these. These disclosures are referred to for possible use in working the present invention.

Cancers for Treatment or Prevention by the Method

Cancers that may be treated include tumours that are not vascularized, or not substantially vascularized, as well as vascularized tumours. The cancers may comprise non-solid tumours (such as haematological tumours, for example, leukaemias and lymphomas) or may comprise solid tumours. Types of cancers to be treated with the invention include, but are not limited to, carcinoma, blastoma, and sarcoma, and certain leukaemia or lymphoid malignancies, benign and malignant tumours, and malignancies e.g., sarcomas, carcinomas, and melanomas. Adult tumours/cancers and paediatric tumours/cancers are also included.

Haematologic cancers are cancers of the blood or bone marrow. Examples of haematological (or haematogenous) cancers include leukaemias, including acute leukaemias (such as acute lymphocytic leukaemia, acute myelocytic leukaemia, acute myelogenous leukaemia and myeloblasts, promyelocytic, myelomonocytic, monocytic and erythroleukaemia), chronic leukaemias (such as chronic myelocytic (granulocytic) leukaemia, chronic myelogenous leukaemia, and chronic lymphocytic leukaemia), polycythemia vera, lymphoma, Hodgkin's disease, non-Hodgkin's lymphoma (indolent and high grade forms), multiple myeloma,

Waldenstrom's macroglobulinemia, heavy chain disease, myelodysplastic syndrome, hairy cell leukaemia and myelodysplasia.

Solid tumours are abnormal masses of tissue that usually do not contain cysts or liquid areas. Solid tumours can be benign or malignant. Different types of solid tumours are named for the type of cells that form them (such as sarcomas, carcinomas, and lymphomas). Examples of solid tumours, such as sarcomas and carcinomas, include fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteosarcoma, and other sarcomas, synovioma, mesothelioma, Ewing's tumour, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, lymphoid malignancy, pancreatic cancer, breast cancer, lung cancers, ovarian cancer, prostate cancer, hepatocellular carcinoma, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, medullary thyroid carcinoma, papillary thyroid carcinoma, pheochromocytomas sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, Wilms' tumour, cervical cancer, testicular tumour, seminoma, bladder carcinoma, melanoma, and CNS tumours (such as a glioma (such as brainstem glioma and mixed gliomas), glioblastoma (also known as glioblastoma multiforme) astrocytoma, CNS lymphoma, germinoma, medulloblastoma, Schwannoma craniopharyngioma, ependymoma, pineaioma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, neuroblastoma, retinoblastoma and brain metastases).

In one embodiment, the administered cells express a first antigen binding site (eg, comprised by a CAR) that is designed to treat a particular cancer. For example, it specifically binds to CD19 can be used to treat cancers and disorders, eg, pre-B ALL (paediatric indication), adult ALL, mantle cell lymphoma, diffuse large B-cell lymphoma or for salvage post allogenic bone marrow transplantation. In another embodiment, the first moiety or first binding site specifically binds CD22 to treat diffuse large B-cell lymphoma.

In one embodiment, cancers and disorders include but are not limited to pre-B ALL (paediatric indication), adult ALL, mantle cell lymphoma, diffuse large B-cell lymphoma, salvage post allogenic bone marrow transplantation, and the like can be treated using a combination of bridging agents (or binding moieties or sites comprised by a single agent) that target two or three of: CD19, CD20, CD22, and ROR1 (eg, CD19 and one of the other targets).

In an example, the cells comprises first and second transmembrane proteins (eg, CARs or a CAR and an engineered TCR expressed by a T-cell) that are different, eg that differ in their target antigens (and optionally otherwise are the same). Similarly, the invention may use a mixture of immune cells (eg, a mixture of CAR-cells), eg comprised by the same transplant, wherein the mixture comprises cells comprising transmembrane proteins (eg, CARs or a CAR and an engineered TCR expressed by a T-cell) that are different, eg that differ in their target antigens (and optionally otherwise are the same). This may be useful for reducing resistance to treatment by cancers, for example, or more effectively targeting cell populations such as cancer cells that surface express a plurality of target antigens.

In one embodiment, the antigen binding site specifically binds to mesothelin to treat or prevent mesothelioma, pancreatic cancer or ovarian cancer.

In one embodiment, the antigen binding site specifically binds to CD33/IL3Ra to treat or prevent acute myelogenous leukaemia.

In one embodiment, the antigen binding site specifically binds to c-Met to treat or prevent triple negative breast cancer or non-small cell lung cancer.

In one embodiment, the antigen binding site specifically binds to PSMA to treat or prevent prostate cancer.

In one embodiment, the antigen binding site specifically binds to Glycolipid F77 to treat or prevent prostate cancer.

In one embodiment, the antigen binding site specifically binds to EGFRvIII to treat or prevent glioblastoma.

In one embodiment, the antigen binding site specifically binds to GD-2 to treat or prevent neuroblastoma or melanoma.

In one embodiment, the antigen binding site specifically binds to NY-ESO-1 TCR to treat myeloma, sarcoma or melanoma.

In one embodiment, the antigen binding site specifically binds to MAGE A3 TCR to treat myeloma, sarcoma and melanoma.

Specific antigen binding is binding with a KD of 1 mM or lower (eg, 1 mM or lower, 100 nM or lower, 10 nM or lower, 1 nM or lower, 100 pM or lower, or 10 pM or lower) as determined by Surface Plasmon Resonance (SPR) in vitro at 25 degrees celcius or rtp.

In one example, said treatment using the method reduces progression of the disease or condition or a symptom thereof. In one example, said treatment using the method reduces incidence of the disease or condition or symptom thereof, eg, for at least 1, 2, 3, 4, or 5 years.

In an example, the method is in vivo in a mammal, eg, a human, man or woman, or male child or female child, or a human infant (eg, no more than 1, 2, 3 or 4 years of age). In an example, the patient is an adult human or a paediatric human patient.

The CAR or TCR is engineered, ie, comprises a non-naturally-occurring combination of moieties and domains. In an example, the cell therapy targets a target cell, wherein the target cell is a cancer cell, eg, a leukaemic cell, lymphoma cell, adenocarcinoma cell or cancer stem cell. Optionally, the CAR or TCR of administered immune cells specifically binds to human CD19 (and optionally the target cell is a leukaemic or lymphoma cell), EpCAM (and optionally the target cell is a lung cancer cell, gastrointestinal cancer cell, an adenocarcinoma, cancer stem cell), CD20 (and optionally the target cell is a leukaemic cell), MCSP (and optionally the target cell is a melanoma cell), CEA, EGFR, EGFRvIII, sialyl Tn, CD133, CD33 (and optionally the target cell is a leukaemic cell, eg, AML cell), PMSA, WT1, CD22, L1CAM, ROR-1, MUC-16, CD30, CD47, CD52, gpA33, TAG-72, mucin, CIX, GD2, GD3, GM2, CD123, VEGFR, integrin, cMET, Her1, Her2, Her3, MAGE1, MAGE A3 TCR, NY-ESO-1, IGF1R, EPHA3, CD66e, EphA2, TRAILR1, TRAILR2, RANKL, FAP, Angiopoietin, mesothelin, Glycolipid F77 or tenascin.

Optionally, the CAR comprises the variable domains of an antibody selected from the group consisting of the CD19 binding site of blinatumomab or antibody HD37; EpCAM binding site of Catumaxomab; CD19 binding site of AFM11; CD20 binding site of Lymphomun; Her2 binding site of Ertumaxomab; CEA binding site of AMG211 (MEDI-565, MT111); PSMA binding site of Pasotuxizumab; EpCAM binding site of solitomab; VEGF or angiopoietin 2 binding site of RG7221 or RG7716; Her1 or Her3 binding site of RG7597; Her2 or Her3 binding site of MM111; IGF1R or Her3 binding site of MM141; CD123 binding site of

MGD006; gpa33 binding site of MGD007; CEA binding site of TF2; CD30 binding site of AFM13; CD19 binding site of AFM11; and Her1 or cMet binding site of LY3164530.

Optionally, the CAR comprises the variable domains of an antigen binding site of an antibody selected from the group consisting of REOPRO® (Abciximab); RITUXAN® (Rituximab); ZENAPAX® (Daclizumab); SIMULECT® (Basiliximab); SYNAGIS® (Palivizumab); REMICADE® (Infliximab); HERCEPTIN® (Trastuzumab); MYLOTARG® (Gemtuzumab ozogamicin); CAMPATH® (Alemtuzumab); ZEVALIN® (Ibritumomab); HUMIRA® (Adalimumab); XOLAIR® (Omalizumab); BEXXAR® (Tositumomab); RAPTIVA® (Efalizumab); ERBITUX® (Cetuximab); AVASTIN® (Bevacizumab); TYSABRI® (Natalizumab); ACTEMRA® (Tocilizumab); VECTIBIX® (Panitumumab); LUCENTIS® (Ranibizumab); SOLIRIS® (Eculizumab); CIMZIA® (Certolizumab); SIMPONI® (Golimumab); ILARIS® (Canakinumab); STELARA® (Ustekinumab); ARZERRA® (Ofatumumab); PROLIA® (Denosumab); NUMAX® (Motavizumab); ABTHRAX® (Raxibacumab); BENLYSTA® (Belimumab); YERVOY® (Ipilimumab); ADCETRIS® (Brentuximab vedotin); PERJETA® (Pertuzumab); KADCYLA® (Ado-trastuzumab); and GAZYVA® (Obinutuzumab).

In an example, the target cell is a blood cell, eg, a stem cell or bone marrow cell of a human or animal. Optionally, the target cell is a B- or T-cell.

In an example, the CAR or TCR comprises an antigen binding site for an autoimmune disease target and the signaling down-regulates cytotoxic activity or proliferation of the immune cells. The term "autoimmune disease" as used herein is defined as a disorder that results from an autoimmune response. An autoimmune disease is the result of an inappropriate and excessive response to a self-antigen. Examples of autoimmune diseases include but are not limited to, Addison's disease, alopecia areata, ankylosing spondylitis, autoimmune hepatitis, autoimmune parotitis, Crohn's disease, diabetes (Type I), dystrophic epidermolysis bullosa, epididymitis, glomerulonephritis, Graves' disease, Guillain-Barré syndrome, Hashimoto's disease, hemolytic anemia, systemic lupus erythematosus, multiple sclerosis, myasthenia gravis, pemphigus vulgaris, psoriasis, rheumatic fever, rheumatoid arthritis, sarcoidosis, scleroderma, Sjogren's syndrome, spondyloarthropathies, thyroiditis, vasculitis, vitiligo, myxedema, pernicious anemia, ulcerative colitis, among others.

Within the overall memory T cell population, several distinct subpopulations have been described and can be recognized by the differential expression of chemokine receptor CCR7 and L-selectin (CD62L). Stem memory T_{SCM} cells, like naive cells, are CD45RO⁻, CCR7⁺, CD45RA⁺, CD62L⁺ (L-selectin), CD27⁺, CD28⁺ and IL-7R α ⁺, but they also express large amounts of CD95, IL-2R β , CXCR3, and LFA-1, and show numerous functional attributes distinctive of memory cells. Central memory T_{CM} cells express L-selectin and the CCR7, they secrete IL-2, but not IFN γ or IL-4. Effector memory T_{EM} cells, however, do not express L-selectin or CCR7 but produce effector cytokines like IFN γ and IL-4. Memory T-cells, such as T_{SCM} may be particularly useful for establishing a sustained population of engineered immune cells in the human.

Any immune cell, target cell or stem cell herein can, in an example, be a T_{SCM} , T_{CM} or T_{EM} cell, eg, a human T_{SCM} , T_{CM} or T_{EM} cell. In an example, the immune cells of the cell therapy (eg, CAR-T cells) each is a progeny of a cell of a human suffering from an autoimmune disease, an inflam-

matory disease, a viral infection or a cancer, eg, wherein the human is suffering from lymphoblastic leukaemia, ALL (eg, T-ALL), CLL (eg, B-cell chronic lymphocytic leukaemia) or non-Hodgkin's lymphoma. The human may, for example, be the patient or a relative (eg, sibling or parent) thereof.

In an example, the administered immune cells have been engineered for enhanced signaling, wherein the signaling is selected from CD28, 4-1BB, OX40, ICOS and CD40 signaling.

Optionally, the target cells (eg, tumour cells) are killed. In an example, each target cell is a tumour cell and the method treats or reduces the risk of cancer, or treats or reduces the risk of cancer progression in the human.

Optionally, the human has cancer. In an example, the cancer is a haematological cancer. In an example, the human has a cancer of B-cell origin. In an example, the human has a cancer of T-cell origin. For example the cancer is lung cancer, melanoma, breast cancer, prostate cancer, colon cancer, renal cell carcinoma, ovarian cancer, neuroblastoma, rhabdomyosarcoma, leukaemia and lymphoma. Preferred cancer targets for use with the present invention are cancers of B cell origin, particularly including acute lymphoblastic leukaemia, B-cell chronic lymphocytic leukaemia or B-cell non-Hodgkin's lymphoma. In an example, the cancer is a cancer of T-cell or B-cell origin, eg, lymphoblastic leukaemia, ALL (eg, T-ALL), CLL (eg, B-cell chronic lymphocytic leukaemia) or non-Hodgkin's lymphoma.

Optionally, each administered immune cell (eg, CAR-cells) is a progeny of an immune cell of said human, eg, wherein the human is suffering from lymphoblastic leukaemia, Diffuse Large B-cell Lymphoma (DLBCL), ALL (eg, T-ALL or B-ALL), CLL (eg, B-cell chronic lymphocytic leukaemia) or non-Hodgkin's lymphoma. Optionally, each administered immune cell (eg, CAR-cells) is an autologous cell (eg, T-cell) of said human or is a progeny of such an autologous cell. As used herein, the term "autologous" is meant to refer to any material derived from the same individual to which it is later to be re-introduced into the individual. "Allogeneic" refers to a graft derived from a different animal of the same species.

Optionally, each administered immune cell (eg, CAR-cells) is derived from a blood or tumour sample of the human and activated and expanded in vitro before step (c). "Activation," as used herein, refers to the state of a T-cell or other immune cell that has been sufficiently stimulated to induce detectable cellular proliferation. Activation can also be associated with induced cytokine production, and detectable effector functions. The term "activated T cells" refers to, among other things, T cells that are undergoing cell division.

In an embodiment, the human has an autoimmune disease, wherein the immune cells that are administered (eg, CAR-cells) are anergic, or have reduced proliferation and/or cytotoxic activity when bound to target cells, whereby the cell transplant cells (and/or their progeny) compete with endogenous immune cells of said human that up-regulate said autoimmune disease.

The administration of immune cells in the method may be by cell infusion into the blood of the patient. The immune cells may be expanded to produce an expanded immune cell population that is administered to the patient. The immune cells may be activated produce an activated immune cell population that is administered to the patient. In methods herein, an effective amount of immune cells are administered. An "effective amount" as used herein, means an amount which provides a therapeutic or prophylactic benefit to treat or prevent the disease or condition.

In an embodiment of the method of the invention, the method treats or reduces the risk of cancer in a patient (eg, a human), wherein the patient has undergone lymphodepletion before administration of the immune cells to the patient.

In one embodiment, the human is resistant to at least one chemotherapeutic agent.

In one embodiment, the chronic lymphocytic leukaemia is refractory CD 19+ leukaemia and lymphoma.

The invention also includes a method of generating a persisting population of genetically engineered T cells in a human diagnosed with cancer, wherein the administered cells comprise T-cells and the persisting population comprises progeny thereof. In one embodiment, the method comprises administering to a human a T-cell population (eg, a CAR T-cell population), wherein the persisting population of genetically engineered T-cells persists in the human for at least one month after administration. In one embodiment, the persisting population of genetically engineered T-cells comprises a memory T-cell. In one embodiment, the persisting population of genetically engineered T-cells persists in the human for at least three months after administration. In another embodiment, the persisting population of genetically engineered T-cells persists in the human for at least four months, five months, six months, seven months, eight months, nine months, ten months, eleven months, twelve months, two years, or three years after administration.

In one embodiment, the chronic lymphocytic leukaemia is treated. The invention also provides a method of expanding a population of the engineered T-cells or NK cells in a human diagnosed with cancer, wherein the administered cells comprise T-cells and/or NK cells and the expanded population comprises progeny thereof.

Optionally, autologous lymphocyte infusion is used in the treatment. For example, autologous PBMCs are collected from a patient in need of treatment and CAR-T-cells are engineered to express the CAR transmembrane protein, activated and expanded using the methods known in the art and then infused back into the patient in step (a).

In an example, the administered cells are pluripotent or multipotent.

The stem cell cannot develop into a human. In an embodiment, the stem cell cannot develop into a human embryo or zygote.

In an example, the administered cell population comprises bone marrow stem cells, eg, human autologous or allogeneic cells.

In an example, the administered cell population comprises haematopoietic stem cells, eg, human autologous or allogeneic cells.

Modifying Microbiota

Medical practice often involves the administration of antibiotics to patients. Such treatments can typically involve administration of broad-spectrum antibiotics, or antibiotics that target many gram-positive bacterial species or many gram-negative species without discrimination. Similarly, use of broad-spectrum antibiotics in farming and agriculture, for example, raise environmental concerns, including entry of such antibiotics into the human and animal food chain which may be deleterious to health and may add to development of microbial resistance. Rather, in an example, the invention involves selective targeting of a first microbial (eg, bacterial or archaeal) species or strain of the microbiota. As shown in the worked examples herein, selective targeting of a particular bacterial species has been achieved using guided nuclease targeting of the genome of the selected species, whilst at the same time sparing related species and strains, as well as species that co-reside (in the Examples species

that co-reside in human gut microbiota). Thus, in one example, the step of causing dysbiosis or step (b) comprises killing first cells of a microbiota sub-population or inhibiting growth of said sub-population by using guided nuclease (eg, RNA guided nuclease) targeting to the genome of first cells comprised by a microbiota sub-population. Suitable systems for carrying out the guided nuclease targeting are, for example, engineered CRISPR/Cas systems, TALENs, meganucleases and zinc finger systems. By way of example, CRISPR/Cas-mediated guided targeting of a selected human gut microbiota bacterial species in a consortium is demonstrated in the Examples herein. The targeting produces nuclease cutting of target species or strain DNA, for example, which reduces the relative proportion of the species in the microbiota or inhibits growth of the sub-population of said species in the microbiota. Selective targeting of species in the method is generally advantageous to enable finer control over change in the relative proportions of bacterial and/or archaeal species in the microbiota. In this way, the invention provides the ability to alter the microbiota with the aim of influencing the upregulation or downregulation of particular immune cell populations, such as T_H1 , T_H17 and/or T_{reg} cells (be these cells endogenous to the patient and/or comprised by adoptive immune cell populations that are administered to the patient), or other outcomes of modulating the microbiota as described herein.

In an example first cell population growth is reduced by at least 5-fold compared to the growth before said dysbiosis or step (b). The method may comprise inhibiting first cell population growth on a gut surface. The method may comprise inhibiting first cell population growth on a plant (eg, leaf and/or stem) surface.

In an alternative, instead of being applied to a subject, the treatment is applied to an environment or soil (eg, the treatment is a fertiliser, plant growth promoting or inhibiting, herbicide or pesticide treatment), wherein the treatment is modulated by the invention.

It will be readily apparent to the skilled addressee how to determine changes in bacteria and archaea in a gut microbiota or other microbiota. For example, this can be done by analyzing a faecal sample of the patient before and after the treatment. One may determine the types of different species or strains in each sample and the proportion of species or strains before and after treatment. Using conventional analysis of 16s ribosomal RNA-encoding DNA (16s rDNA) it is possible to identify species, for example. Additionally or alternatively, standard biochemical test can be used to identify strains or species, eg, also involving one or more of: staining, motility testing, serological testing, phage typing and identification disc testing (eg using a Kirby Baur disc diffusion method). Biochemical testing may involve one or more of: a (a) Catalase test (b) Coagulase test (c) Oxidase test (d) Sugar fermentation test (e) Indole test (f) Citrate test and (g) Urease test. Relative proportions may be determined by growing colonies on agar plates (as in the Examples herein) from each sample and counting colony numbers.

In an example, the dysbiosis or step (b) increases the proportion of *Bacteroides* (eg, *B. fragalis* and/or *B. thetaiota-micron*) in the microbiota, eg, gut microbiota.

In an example, the dysbiosis or step (b) decreases the proportion of *Bacteroides* (eg, *B. fragalis* and/or *B. thetaiota-micron*) in the microbiota, eg, gut microbiota. In an example, the dysbiosis or step (b) increases the proportion of *Bacteroidetes* to *Firmicutes* in the microbiota, eg, gut microbiota. In an example, the dysbiosis or step (b) decreases the proportion of *Bacteroidetes* to *Firmicutes* in the microbiota, eg, gut microbiota.

Accumulating evidence supports the role of commensal strains of *Bifidobacterium* and *Clostridium* spp. belonging to clusters IV and XIVa in the induction of Treg cells. See, eg, Lopez et al. In an example, the dysbiosis or step (b) reduces the proportion of one or more *Clostridium* species or strain (eg, In an example, each species is a cluster IV or XIVa *Clostridium* species) in the gut microbiota. In an example, the dysbiosis or step (b) increases the proportion of one or more *Clostridium* species or strain (eg, In an example, each species is a cluster IV or XIVa *Clostridium* species) in the gut microbiota.

In an example, the dysbiosis or step (b) reduces the proportion of *Bifidobacterium* (eg, *B bifidum*) in the gut microbiota. In an example, the dysbiosis or step (b) increases the proportion of *Bifidobacterium* (eg, *B bifidum*) in the gut microbiota.

For example, by selectively altering the human gut microbiota the invention provides for upregulation of CAR-T or other ACT treatment (eg, wherein the altered microbiota downregulates T_{reg} cells in the patient that has received the CAR-T or ACT administration and/or upregulates T_{H1} and/or T_{H17} cells in the patient—such cells being comprised by the CAR-T or ACT transplant for example). Downregulating T_{reg} cells may reduce suppression of T-effectors and/or T-helpers in the patient, thereby enhancing the CAR-T or ACT cytotoxicity or other desirable activity against cancer or other disease-mediating cells. Upregulating T_{H1} and/or T_{H17} cells may increase T-effector activity, thereby enhancing the CAR-T or ACT cytotoxicity or other desirable activity against cancer or other disease-mediating cells.

In another example, alteration of the microbiota can be used as a switch to dampen down CAR-T or other ACT treatment (eg, wherein the altered microbiota upregulates T_{reg} cells in the patient that has received the CAR-T or ACT administration and/or downregulates T_{H1} and/or T_{H17} cells in the patient—such cells being comprised by the CAR-T or ACT transplant for example). Upregulating T_{reg} cells may increase suppression of T-effectors and/or T-helpers in the patient, thereby reducing the CAR-T or ACT ability to promote cytokine release or other undesirable activity. Downregulating T_{H1} and/or T_{H17} cells may decrease T-effector activity, thereby reducing the CAR-T or ACT ability to promote cytokine release or other undesirable activity. This may be useful for limiting the risk of cytokine release syndrome (CRS) in the patient. Subsequent further modification of the gut microbiota of the patient using the method of the invention can be performed to upregulate the CAR-T or ACT treatment when it is desired to use this once more to address the disease or condition at hand (eg, a cancer, such as a haematological cancer). In this instance, memory T-cell CAR-T or ACT populations may be present in the patient from the earlier treatment, and the upregulation using microbiota alteration according to the invention may upregulate memory T-cells to differentiate into effector and/or helper cells to address the disease or condition. Thus, in one example, the cell therapy of the invention comprises administering an immune cell population comprising immune memory cells (eg, memory T-cells, such as central memory T cells (T_{CM}) and/or stem cell memory cells (T_{SCM}); and/or the administered population comprises cells that spawn such memory cells following the initial microbiota alteration.

Whilst one aspect of the invention recognizes utility for modulating cell-based therapy in a patient, another aspect recognizes utility for modulating (ie, treating or preventing) cell-mediated diseases and conditions in patients, such as autoimmune and inflammatory diseases and conditions which are mediated, for example by T-cells or other immune

cells of the patient. In a further aspect, the invention recognizes utility as a means for modulating (eg, enhancing) another therapy of a disease or condition; for example, for enhancing or effecting therapy with an antibody or anti-viral medicine to treat or prevent the disease or condition. For example, the medicine can be an immune checkpoint antagonist or agonist (eg, for treating or preventing a cancer, such as melanoma or NSCLC). By “effecting therapy” it is contemplated that the patient does not respond or poorly responds to the medicine and the microbiota alteration according to the invention (eg, using selective guided nuclease targeting of a bacterial or archaeal species as described herein) brings about a response (or improved response) to the medicine by the patient. For example, the method of the invention upregulates T_{H17} cells in a patient suffering from HIV infection. In one aspect, this enhances anti-retroviral therapy or HIV vaccine therapy of the patient. The T_{H17} cells may be the patient’s endogenous cells or cells provided by ACT of the patient. In another example, the method of the invention upregulates T_{H17} cells in a patient suffering from a cancer (eg, melanoma or lung cancer, such as NSCLC). In one aspect, this enhances immune checkpoint antagonism or agonism therapy of the patient. The T_{H17} cells may be the patient’s endogenous cells or cells provided by ACT of the patient. For example, the therapy is antibody therapy using an antibody selected from ipilimumab (or YERVOY™), tremelimumab, nivolumab (or OPDIVO™), pembrolizumab (or KEYTRUDA™), pidilizumab, BMS-936559, durvalumab and atezolizumab.

The invention relates to guided nuclease systems (eg, engineered CRISPR/Cas systems, TALENs, meganucleases and zinc finger systems), arrays (eg, CRISPR arrays), cRNAs, gRNAs and vectors (eg, phage comprising components of a said system) for use in a method of the invention for targeting the first cells or causing said dysbiosis by inhibiting bacterial or archaeal cell population growth or altering the relative proportion of one or more sub-populations of cells in plant, yeast, environmental, soil, human or animal microbiota, such as for the alteration of the proportion of Bacteroidetes (eg, *Bacteroides*), *Firmicutes* and/or gram positive or negative bacteria in gut microbiota of a human. The invention, for example, involves modifying (eg, cutting and/or mutating) one or more target genomic or episomal nucleotide sequences of a host bacterial cell, eg, a Bacteroidetes cell or *Firmicutes* cell, or a host archaeal cell. In an example, the first bacteria are pathogenic gut bacteria.

There have been a number of studies pointing out that the respective levels of the two main intestinal phyla, the Bacteroidetes and the *Firmicutes*, are linked to obesity, both in humans and in germ-free mice. The authors of the studies deduce that carbohydrate metabolism is the important factor. They observe that the microbiota of obese individuals are more heavily enriched with bacteria of the phylum *Firmicutes* and less with Bacteroidetes, and they surmise that this bacterial mix may be more efficient at extracting energy from a given diet than the microbiota of lean individuals (which have the opposite proportions). In some studies, they found that the relative abundance of Bacteroidetes increases as obese individuals lose weight and, further, that when the microbiota of obese mice are transferred to germfree mice, these mice gain more fat than a control group that received microbiota from lean mice. See, eg, Turnbaugh, P. J., R. E. Ley, M. A. Mahowald, V. Magrini, E. R. Mardis, and J. I. Gordon. 2006, “An obesity-associated gut microbiome with increased capacity for energy harvest”, Nature 444:1027-1131. In a further aspect, the invention recognizes utility as

a means for enhancing an anti-obesity therapy of a patient, eg, by increasing the ratio of Bacteroidetes versus *Firmicutes* in the microbiota.

Optionally the first cells are in the presence of cells of a different strain or species, wherein the different cells are Enterobacteriaceae or bacteria that are probiotic, commensal or symbiotic with humans (eg, in the human gut). In an example, each first cell is a *Firmicutes*, eg, *Streptococcus*, cell.

In an example, the invention is able to selectively kill or downregulate the target microbes in the microbiota whilst not targeting a second related strain of the same species or a different species that is nevertheless phylogenetically related (as indicated by 16s rDNA). For example, the microbiota comprises cells of a second bacterial species or strain, or archaeal species or strain, wherein the second species or strain has a 16s ribosomal RNA-encoding DNA sequence that is at least 80, 85, 90, 95, 96, 97, 98 or 99% identical to an 16s ribosomal RNA-encoding DNA sequence of the first cell species or strain, wherein the growth of the second cells in the microbiota is not inhibited by said method. In an embodiment, the growth of second strain or species is not inhibited; or the growth of said first cells is inhibited by at least 2x, 3x, 4x, 5x, 6x, 7x, 8x, 9x, 10x, 50x, 100x or 1000x the growth inhibition of the second cells.

In one aspect of the method, causing the dysbiosis or step (b) comprises altering the proportion of a sub-population of first cells (host cells) in the microbiota, eg, gut microbiota, of the patient, thereby producing an altered gut microbiota that modulates the immune cell therapy in the patient, wherein the sub-population comprises host cells of said first species or strain, the method comprising using guided nuclease (eg RNA-guided nuclease) cutting of a respective target sequence in host cells to modify the target sequences, whereby host cells are killed or the host cell population growth is reduced, thereby reducing the proportion of said sub-population in the microbiota. Suitable systems for carrying out the guided nuclease cutting are, for example, engineered CRISPR/Cas systems, TALENs, meganucleases and zinc finger systems. By way of example, CRISPR/Cas-mediated guided cutting of a selected human gut microbiota bacterial species in a consortium is demonstrated in the Examples herein.

In an example, the target sequence modification is carried out by

- a. combining the microbiota with multiple copies of engineered nucleic acid sequences encoding host modifying (HM) crRNAs, and
- b. expressing HM-crRNAs in host cells,

wherein each engineered nucleic acid sequence is operable with a Cas nuclease in a respective host cell to form a HM-CRISPR/Cas system and the engineered sequence comprises

- (i) spacer and repeat sequences encoding a HM-crRNA;
- (ii) the HM-crRNA comprising a sequence that is capable of hybridizing to a host cell target sequence to guide Cas nuclease to the target sequence in the host cell; and optionally the HM-system comprises a tracrRNA sequence or a DNA sequence expressing a tracrRNA sequence;

whereby HM-crRNAs guide Cas modification of host target sequences in host cells, whereby host cells are killed or the host cell population growth is reduced, thereby reducing the proportion of said sub-population in the microbiota.

In an alternative, HM-crRNA and tracrRNA are comprised by a single guide RNA (gRNA). In an example, each

engineered nucleic acid sequence is comprised by a respective vector, wherein each vector is optionally a plasmid (eg, a conjugative plasmid capable of transfer into a host cell), phage, phagemid or prophage. The phage is capable of infecting a said host cell.

In an example, endogenous Cas nuclease of host cells is used for modification of target nucleotide sequences. In an embodiment, therefore, each vector lacks a Cas (eg, a Cas9) nuclease-encoding sequence. By harnessing endogenous Cas nuclease, embodiments of the invention use endogenous Cas nuclease activity (ie, without the need for prior genetic modification of the host cell to activate or enhance the nuclease activity). Thus, in an example, the Cas nuclease is encoded by a wild-type gene of the host cell. In an example, the nuclease is active to achieve the cell killing or growth reduction without inhibition of an endogenous Cas nuclease (or Cas nuclease gene) repressor in the host cell. Thus, the invention can address wild-type bacterial populations without the need for prior manipulation to make bring about effective Cas-mediated cell killing or growth reduction. Thus, the population can be exposed to the crRNA when the population is in its wild-type environment (such as comprised by a plant, yeast, environmental, soil, human or animal microbiome).

In an example, the crRNA or gRNA is for administration to (or administered to) a human or non-human animal patient by mucosal, gut, oral, intranasal, intrarectal or buccal administration.

Optionally said Cas nuclease is provided by an endogenous Type II CRISPR/Cas system of each first cell. Optionally, the tracrRNA sequence or DNA sequence expressing a tracrRNA sequence is endogenous to each host cell. Optionally, each target sequence is comprised by an antibiotic resistance gene, virulence gene or essential gene of the respective host cell, for example the target sequences are identical between the host cells. Optionally, the engineered nucleic acid sequences are comprised by an antibiotic composition, wherein the sequences are in combination with an antibiotic agent (first antibiotic), and in an example the target sequences are comprised by an antibiotic resistance gene wherein the antibiotic is said first antibiotic. The antibiotic composition is administered to the patient or subject to effect said dysbiosis or step (b).

Optionally, each host cell comprises a deoxyribonucleic acid strand with a free end (HM-DNA) encoding a HM-sequence of interest and/or wherein the method comprising into the host cells such a sequence encoding the HM-DNA, wherein the HM-DNA comprises a sequence or sequences that are homologous respectively to a sequence or sequences in or flanking the target sequence for inserting the HM-DNA into the host genome (eg, into a chromosomal or episomal site).

The invention also provides vectors for introducing into first cells (host cells) for carrying out the treatment or prevention method of the invention, wherein each vector is: An engineered nucleic acid vector for modifying a bacterial or archaeal host cell comprising an endogenous CRISPR/Cas system, the vector comprising nucleic acid sequences for expressing a plurality of different crRNAs (eg, gRNAs) for use in causing the dysbiosis or for use in step (b) of the method; and optionally lacking a nucleic acid sequence encoding a Cas nuclease, wherein a first of said crRNAs is capable of hybridising to a first nucleic acid sequence in said host cell; and a second of said crRNAs is capable of hybridising to a second nucleic acid sequence in said host cell, wherein said second sequence is different from said first sequence; and

- a. the first sequence is comprised by an antibiotic resistance gene (or RNA thereof) and the second sequence is comprised by an antibiotic resistance gene (or RNA thereof); optionally wherein the genes are different;
- b. the first sequence is comprised by an antibiotic resistance gene (or RNA thereof) and the second sequence is comprised by an essential or virulence gene (or RNA thereof);
- c. the first sequence is comprised by an essential gene (or RNA thereof) and the second sequence is comprised by an essential or virulence gene (or RNA thereof); or
- d. the first sequence is comprised by a virulence gene (or RNA thereof) and the second sequence is comprised by an essential or virulence gene (or RNA thereof).

Each vector may be as described above, eg, a phage capable of infecting a host cell or conjugative plasmid capable of introduction into a host cell. In an example, the vectors are in combination with an antibiotic agent (eg, a beta-lactam antibiotic).

Each first cell (host cell) may be a *Staphylococcus*, *Streptococcus*, *Pseudomonas*, *Salmonella*, *Listeria*, *E coli*, *Desulfovibrio* or *Clostridium* host cell. In an example, each first cell (host cell) is a *Firmicutes* cell, eg, a *Staphylococcus*, *Streptococcus*, *Listeria* or *Clostridium* cell.

In an example, each engineered nucleic acid sequence comprises a sequence R1-S1-R1' for expression and production of the respective crRNA (eg, comprised by a single guide RNA) in the host cell, (i) wherein R1 is a first CRISPR repeat, R1' is a second CRISPR repeat, and R1 or R1' is optional; and (ii) S1 is a first CRISPR spacer that comprises or consists of a nucleotide sequence that is 95% or more identical to said target sequence.

In an example, R1 and R1' are at least 95% identical respectively to the first and second repeat sequences of a CRISPR array of the host cell species. In an example, R1 and R1' are at least 95% (eg, 96, 97, 98, 99 or 100%) identical respectively to the first (5'-most) and second (the repeat immediately 3' of the first repeat) repeat sequences of a CRISPR array of said species, eg, of a said host cell of said species. In an example, R1 and R1' are functional with a Type II Cas9 nuclease (eg, a *S thermophilus*, *S pyogenes* or *S aureus* Cas9) or Type I Cas3 to modify the target in a said host cell.

In one aspect, the method involves the following use, as demonstrated by the worked experimental Example:

The use of wild-type endogenous Cas nuclease activity of the first cell (host cell) population to inhibit growth of the population, wherein each host cell has an endogenous CRISPR/Cas system having wild-type Cas nuclease activity, the use comprising transforming host cells of the population, wherein each transformed host cell is transformed with an engineered nucleotide sequence for providing host modifying (HM) crRNA or guide RNA (gRNA) in the host cell, the HM-crRNA or gRNA comprising a sequence that is capable of hybridising to a host cell target protospacer sequence for guiding endogenous Cas to the target, wherein the crRNA or gRNA is cognate to an endogenous Cas nuclease of the host cell that has said wild-type nuclease activity and following transformation of the host cells growth of the population is inhibited.

By "cognate to" it is intended that the endogenous Cas is operable with crRNA or gRNA sequence to be guided to the target in the host cell. The skilled addressee will understand that such Cas guiding is generally a feature of CRISPR/Cas activity in bacterial and archaeal cells, eg, wild-type CRISPR/Cas activity in bacterial or archaeal cells having endogenous active wild-type CRISPR/Cas systems.

By "wild-type" Cas activity it is intended, as will be clear to the skilled addressee, that the endogenous Cas is not an engineered Cas or the cell has not been engineered to de-repress the endogenous Cas activity. This is in contrast to certain bacteria where Cas nuclease activity is naturally repressed (ie, there is no wild-type Cas nuclease activity or none that is useful for the present invention, which on the contrary is applicable to addressing wild-type host cells in situ for example where the endogenous Cas activity can be harnessed to effect cell population growth inhibition).

In the worked Examples below, inhibition was addressed in a bacterial population (a gram positive *Firmicutes*) on a solid surface. A >10-fold inhibition of host cell population growth was achieved. Targeting was directed to an antibiotic resistance gene and an essential gene. The demonstration of the invention's ability to inhibit host cell growth on a surface is important and desirable in embodiments where the invention is for treating or preventing diseases or conditions mediated or caused by microbiota as disclosed herein in a human or animal subject. Such microbiota are typically in contact with tissue of the subject (eg, gut, tissue) and thus we believe that the demonstration of activity to inhibit growth of a microbiota bacterial species (exemplified by *Streptococcus*) on a surface supports this utility. Targeting microbiota on plant surfaces is also a desired application.

In an example, inhibition of first cell (host cell) population growth is at least 2, 3, 4, 5, 6, 7, 8, 9 or 10-fold compared to the growth of cells of the same species or strain not exposed to said engineered nucleotide sequence. For example, growth inhibition is indicated by a lower bacterial colony number of a first sample of host cells (alone or in a mixed bacterial population, eg, a microbiota or faecal sample of the patient after treatment) by at least 2, 3, 4, 5, 6, 7, 8, 9 or 10-fold compared to the colony number of a second sample of the host cells (alone or in a mixed bacterial population, eg, a microbiota or faecal sample of the patient before treatment), wherein the first cells have been transformed by said engineered nucleotide sequence but the second sample has not been exposed to said engineered nucleotide sequence. In an embodiment, the colony count is determined 12, 24, 36 or 48 hours after the first sample has been exposed to the engineered sequence. In an embodiment, the colonies are grown on solid agar in vitro (eg, in a petri dish). It will be understood, therefore, that growth inhibition can be indicated by a reduction (<100% growth compared to no treatment, ie, control sample growth) in growth of first (host) cells or populations, or can be a complete elimination of such growth. In an example, growth of the host cell population is reduced by at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 95%, ie, over a predetermined time period (eg, 24 hours or 48 hours following combination with the crRNA or gRNA, TALEN, meganuclease, zinc finger etc in the host cells), ie, growth of the host cell population is at least such percent lower than growth of a control host cell population that has not been exposed to said crRNA or gRNA etc but otherwise has been kept in the same conditions for the duration of said predetermined period. In an example, percent reduction of growth is determined by comparing colony number in a sample of each population at the end of said period (eg, at a time of mid-exponential growth phase of the control sample). For example, after exposing the test population to the crRNA or gRNA etc at time zero, a sample of the test and control populations is taken and each sample is plated on an agar plate and incubated under identical conditions for said predetermined period. At the end of the period, the colony number of each sample is counted and the percentage difference (ie, test colony number divided by

control colony number and then times by 100, and then the result is subtracted from 100 to give percentage growth reduction). The fold difference is calculated by dividing the control colony number by the test colony number.

Inhibition of population growth can be indicated, therefore, by a reduction in proliferation of first (host) cell number in the population. This may be due to cell killing by the nuclease and/or by downregulation of host cell proliferation (division and/or cell growth) by the action of the nuclease on the target protospacer sequence. In an embodiment of a treatment or prevention as disclosed herein, host cell burden of the human or animal subject is reduced, whereby the disease or condition is treated (eg, reduced or eliminated) or prevented (ie, the risk of the subject developing the disease or condition) is reduced or eliminated.

In an example, wild-type host cell endogenous Cas9 or *cpf1* activity is used. In an example, wild-type host cell endogenous Cas3 and/or CASCADE activity is used. The engineered nucleotide sequence may not be in combination with an exogenous Cas nuclease-encoding sequence. Optionally, said Cas nuclease is a nickase.

In an example, the formation of bacterial colonies of said host cells is inhibited following said dysbiosis or step (b). In an example, proliferation of host cells is inhibited following said dysbiosis or step (b). In an example, host cells are killed following said dysbiosis or step (b).

In another aspect, the method comprises producing *ex vivo* a medicament for administration to the patient for causing said dysbiosis or step (b) for treating or preventing the disease or condition, wherein the medicament comprises a modified mixed bacterial population (eg, obtained from faeces or gut microbiota of one or more human donors or said patient), wherein the modified population is administered to the patient to cause said dysbiosis or in step (b) to alter the balance of species or strains in the patient's gut microbiota, thereby altering the proportion of the first cells in the gut microbiota. The modified mixed population can be produced *ex vivo* using guided nuclease modification techniques as described herein. Thus, for example, the method can be used to reduce the proportion of a specific *Firmicutes* sub-population and spare Bacteroidetes in the mixed population, eg, for producing a medicament for treating or preventing a metabolic or GI condition or disease disclosed herein. In this way, the invention can use a modified bacterial transplant (eg, a modified faecal transplant) medicament for such use or for said treatment or prevention in a human or animal. For example, the method can be used to modify one or more microbiota *in vitro* to produce a modified collection of bacteria for administration to a human or animal for medical use (eg, treatment or prevention of a metabolic condition (such as obesity or diabetes) or a GI tract condition (eg, any such condition mentioned herein) or a cancer (eg, a GI tract cancer). In an example, the transformation of bacterial cells with phage or plasmid vectors comprising engineered nucleic acid sequences as described herein is carried out *in vitro*, or the engineered nucleotide sequence is comprised by nucleic acid that is electroporated into host cells. In an example, the nucleic acid are RNA (eg, copies of the gRNA). In another example, the nucleic acid are DNA encoding the crRNA or gRNA for expression thereof in host cells.

Thus, in an example, the invention provides an engineered nucleotide sequence for providing host cell modifying (HM) crRNA or guide RNA (gRNA) in a population of wild-type bacterial host cells comprised by a microbiota of a plant, yeast, environmental, soil, human or animal subject for use in the method of the invention, the crRNA or gRNA com-

prising a sequence that is capable of hybridising to a host cell target protospacer sequence for guiding Cas to the target, wherein the crRNA or gRNA is cognate to an endogenous host cell Cas nuclease that has wild-type nuclease activity, wherein following transformation of host cells growth of the population is inhibited and the disease or condition is treated or prevented, or the therapy or treatment is modulated.

In an example, the engineered nucleotide sequence comprises an HM-CRISPR array. In an example, the engineered nucleotide sequence encodes a single guide RNA. In an example, the engineered nucleotide sequence is a guide RNA (eg, a single guide RNA) or crRNA. In an example, the engineered sequence is comprised by a bacteriophage that is capable of infecting the host cells, wherein the transformation comprises transduction of the host cells by the bacteriophage. The bacteriophage can be a phage as described herein. In an example, the engineered nucleotide sequence is comprised by a plasmid (eg, a conjugative plasmid) that is capable of transforming host cells. The plasmid can be a plasmid as described herein. In an example, the engineered nucleotide sequence is comprised by a transposon that is capable of transfer into and/or between host cells. The transposon can be a transposon as described herein.

Any method of the invention can comprise transforming host cells with nucleic acid vectors for producing crRNA or gRNA in the cells. For example, the vectors or nucleic acid comprising the engineered nucleotide sequence are administered orally, intravenously, topically, ocularly, intranasally, by inhalation, by rectal administration, or by any other route of administration disclosed herein or otherwise to the patient, wherein the administration transforms the first (host) cells with the vectors or nucleic acid.

In an example, the first are mixed with second bacteria in the microbiota of the patient or subject. Optionally, the second bacteria species is *E coli*, *L lactis* or *S thermophilus*, as shown in the worked Example below, such are strains that co-exist symbiotically in human and animal gut microbiota. The Example also addresses targeting in a mixed gram positive and gram negative bacterial population. Additionally, the Example addresses a population of *Firmicutes* (*S thermophilus*) and a population of Enterobacteriaceae (*E coli*), both of which are found in human microbiota. Other examples of Enterobacteriaceae are *Salmonella*, *Yersinia pestis*, *Klebsiella*, *Shigella*, *Proteus*, *Enterobacter*, *Serratia*, and *Citrobacter*.

In an example, the condition or disease is a metabolic or gastrointestinal disease or condition, eg, obesity, IBD, IBS, Crohn's disease or ulcerative colitis. In an example, the condition or disease is a cancer, eg, a solid tumour or a GI cancer (eg, stomach cancer), liver cancer or pancreatic cancer. In an example, the condition is resistance or reduced responsiveness to an antibiotic (eg, any antibiotic disclosed herein).

In an example, each first (host) cell comprises an endogenous RNase III that is operable with the engineered sequence in the production of said HM-crRNA in the cell. In an alternative, one or more of the vectors comprises a nucleotide sequence encoding such a RNase III for expression of the RNase III in the host cell.

In an example, the essential gene (comprising the target) encodes a DNA polymerase of the cell. This is exemplified below.

In an example, the crRNA or gRNA comprises a sequence that is capable of hybridising to a host cell target protospacer sequence that is adjacent a NGG, NAG, NGA, NGC, NNGNG, NNGRRT or NNAGAAW protospacer adjacent

motif (PAM), eg, a AAAGAAA or TAAGAAA PAM (these sequences are written 5' to 3'). In an embodiment, the PAM is immediately adjacent the 3' end of the protospacer sequence. In an example, the Cas is a *S aureus*, *S therophilus* or *S pyogenes* Cas. In an example, the Cas is Cpf1 and/or the PAM is TTN or CTA.

In an example, each engineered nucleotide sequence or vector comprises a said CRISPR array or a sequence encoding a said crRNA or gRNA and further comprises an antibiotic resistance gene (eg, kanamycin resistance), wherein the HM-crRNA or gRNA does not target the antibiotic resistance gene. In an example, the target sequence is comprised by an antibiotic resistance gene of the host cell, wherein the antibiotic is different from the first antibiotic (eg, kanamycin). In this way, the engineered sequence or vector is able to target the host without targeting itself. By exposing the host cells to the first antibiotic (eg, by orally or intravenously administering it to the patient), one can promote retention of the engineered sequence or vector therein by positive selection pressure since cells containing the first antibiotic resistance gene will have a survival advantage in the presence of the first antibiotic (when host cells that are not transformed by the engineered sequence or vectors are not resistant to the first antibiotic). Thus, an example provides: The method of the invention comprising exposing the first (host) cell population to said antibiotic (eg, kanamycin) and said engineered sequence or vector(s), for promoting maintenance of crRNA or gRNA-encoding sequences in host cells; or the engineered sequence, array or vector of the invention is in combination with said antibiotic.

In an example the sequence encoding the crRNA or gRNA is under a constitutive promoter (eg, a strong promoter) operable in the host cell species, or an inducible promoter.

In an example, the or each first (host) cell is a gram positive cell. In another example, the or each first (host) cell is a gram positive cell.

The invention also provides: An ex vivo mixed population of microbiota bacteria obtainable by the method by isolation of a gut microbiota sample from the patient after carrying out the method, or by isolation of a faecal sample of the patient after carrying out the method. In an example, the mixed population is in a container for medical or nutritional use. For example, the container is a sterilised container, eg, an inhaler, intranasal delivery device or connected to a syringe or IV needle. In an example, the mixed population is useful for administration to a human or animal to populate a microbiome thereof for treating a disease or condition (eg, a disease or condition disclosed herein).

Herein, in an example the *Bacteroides* is a species selected from *caccae*, *capillosus*, *cellulosilyticus*, *coprocola*, *coprophilus*, *coprosuis*, *distasonis*, *dorei*, *eggerthii*, *faecis*, *fingoldii*, *fluxus*, *fragalis*, *intestinalis*, *melaninogenicus*, *nordii*, *oleiciplenus*, *oralis*, *ovatus*, *pectinophilus*, *plebeius*, *stercoris*, *thetaiotaomicron*, *uniformis*, *vulgatus* and *xylanisolvans*. For example, the *Bacteroides* is *thetaiotaomicron*. In an example, the first (host cell) population or second bacteria comprise a plurality of different *Bacteroides* species, or a plurality of *Bacteroides* species (eg, comprising *B thetaiotaomicron* and *B fragalis*), or *Bacteroides* and *Prevotella* species. Herein, in an example, the *Prevotella* is a species selected from *bergensis*, *bivia*, *buccae*, *buccalis*, *copri*, *melaninogenica*, *oris*, *ruminicola*, *tanneriae*, *timonensis* and *veroralis*. In an alternative, the first (host) cells or second bacteria are *Firmicutes* cells, for example comprising or consisting of one or more *Firmicutes* selected from *Anaerotruncus*, *Acetanaerobacterium*, *Acetotomaculum*, *Acetivibrio*, *Anaerococcus*, *Anaerofilum*,

Anaerosinus, *Anaerostipes*, *Anaerovorax*, *Butyrivibrio*, *Clostridium*, *Capracoccus*, *Dehalobacter*, *Dialister*, *Dorea*, *Enterococcus*, *Ethanoligenens*, *Faecalibacterium*, *Fusobacterium*, *Gracilibacter*, *Guggenheimella*, *Hespellia*, *Lachnobacterium*, *Lachnospira*, *Lactobacillus*, *Leuconostoc*, *Megamonas*, *Mitsuokella*, *Oribacterium*, *Oxobacter*, *Papillibacter*, *Propionispira*, *Pseudobutyrvibrio*, *Pseudoramibacter*, *Roseburia*, *Ruminococcus*, *Sarcina*, *Seinonella*, *Shuttleworthia*, *Sporobacter*, *Sporobacterium*, *Streptococcus*, *Subdoligranulum*, *Syntrophococcus*, *Thermobacillus*, *Turibacter* and *Weisella*. In an example, the first (host) cells or second bacteria comprise or consist of *Clostridium* (eg, *dificile*) cells (and optionally the other sub-population consists of *Bacteroides* (eg, *thetaiotaomicron*) cells). In an example, the first (host) cells or second bacteria comprise or consist of *Enterococcus* cells (and optionally the other cells consist of *Bacteroides* (eg, *thetaiotaomicron*) cells). In an example, the first (host) cells or second bacteria comprise or consist of *Ruminococcus* cells (and optionally the other cells consist of *Bacteroides* (eg, *thetaiotaomicron*) cells). In an example, the first (host) cells or second bacteria comprise or consist of *Streptococcus* cells (and optionally the other cells consist of *Bacteroides* (eg, *thetaiotaomicron*) cells). In an example, the first (host) cells or second bacteria comprise or consist of *Faecalibacterium* cells (and optionally the other cells consist of *Bacteroides* (eg, *thetaiotaomicron*) cells). For example, the *Faecalibacterium* is a *Faecalibacterium prausnitzii* (eg, A2-165, L2-6, M21/2 or SL3/3).

In an example, the first (host) cells or second bacteria consist of *Streptococcus* cells (optionally *S thermophilus* and/or *pyogenes* cells) and the second bacteria consists of *Bacteroides* (eg, *thetaiotaomicron*) and/or Enterobacteriaceae (eg, *E coli*) cells.

In an example, the first (host) cells are infectious disease pathogens of humans, an animal (eg, non-human animal) or a plant.

In an example, the first (host) cells are selected from a species of *Escherichia* (eg, *E coli* O157:H7 or O104: H4), *Shigella* (eg, *dysenteriae*), *Salmonella* (eg, *typhi* or *enterica*, eg, serotype *typhimurium*, eg, DT 104), *Erwinia*, *Yersinia* (eg, *pestis*), *Bacillus*, *Vibrio*, *Legionella* (eg, *pneumophila*), *Pseudomonas* (eg, *aeruginosa*), *Neisseria* (eg, *gonorrhoea* or *meningitidis*), *Bordetella* (eg, *pertussis*), *Helicobacter* (eg, *pylori*), *Listeria* (eg, *monocytogenes*), *Agrobacterium*, *Staphylococcus* (eg, *aureus*, eg, MRSA), *Streptococcus* (eg, *pyogenes* or *thermophilus*), *Enterococcus*, *Clostridium* (eg, *dificile* or *botulinum*), *Corynebacterium* (eg, *amycolatum*), *Mycobacterium* (eg, *tuberculosis*), *Treponema*, *Borrelia* (eg, *burgdorferi*), *Francisella*, *Brucella*, *Campylobacter* (eg, *jejuni*), *Klebsiella* (eg, *pneumoniae*), *Frankia*, *Bartonella*, *Rickettsia*, *Shewanella*, *Serratia*, *Enterobacter*, *Proteus*, *Providencia*, *Brochothrix*, *Bifidobacterium*, *Brevibacterium*, *Propionibacterium*, *Lactococcus*, *Lactobacillus*, *Pediococcus*, *Leuconostoc*, *Vibrio* (eg, *cholera*, eg, O139, or *vulnificus*), *Haemophilus* (eg, *influenzae*), *Brucella* (eg, *abortus*), *Franciscella*, *Xanthomonas*, *Erlchia* (eg, *chaffeensis*), *Chlamydia* (eg, *pneumoniae*), *Parachlamydia*, *Enterococcus* (eg, *faecalis* or *faceim*, eg, *linezolid-resistant*), *Oenococcus* and *Acinetobacter* (eg, *baumannii*, eg, multiple drug resistant).

In an example, the first (host) cells are *Staphylococcus aureus* cells, eg, resistant to an antibiotic selected from methicillin, vancomycin-resistant and teicoplanin.

In an example, the first (host) cells are *Pseudomonas aeruginosa* cells, eg, resistant to an antibiotic selected from cephalosporins (eg, ceftazidime), carbapenems (eg, imipenem or meropenem), fluoroquinolones, aminoglycosides (eg, gentamicin or tobramycin) and colistin.

In an example, the first (host) cells are *Klebsiella* (eg, *pneumoniae*) cells, eg, resistant to carbapenem.

In an example, the first (host) cells are *Streptococcus* (eg, *pneumoniae* or *pyogenes*) cells, eg, resistant to an antibiotic selected from erythromycin, clindamycin, beta-lactam, macrolide, amoxicillin, azithromycin and penicillin.

In an example, the first (host) cells are *Salmonella* (eg, serotype *Typhi*) cells, eg, resistant to an antibiotic selected from ceftriaxone, azithromycin and ciprofloxacin.

In an example, the first (host) cells are *Shigella* cells, eg, resistant to an antibiotic selected from ciprofloxacin and azithromycin.

In an example, the first (host) cells are *Mycobacterium tuberculosis* cells, eg, resistant to an antibiotic selected from Resistance to isoniazid (INH), rifampicin (RMP), fluoroquinolone, amikacin, kanamycin and capreomycin.

In an example, the first (host) cells are *Enterococcus* cells, eg, resistant to vancomycin.

The method of claim 13, wherein all of the host cells are Enterobacteriaceae cells, eg, resistant to an antibiotic selected from a cephalosporin and carbapenem.

In an example, the first (host) cells are *E. coli* cells, eg, resistant to an antibiotic selected from trimethoprim, itrofurantoin, cefalexin and amoxicillin.

In an example, the first (host) cells are *Clostridium* (eg, *difficile*) cells, eg, resistant to an antibiotic selected from fluoroquinolone antibiotic and carbapenem.

In an example, the first (host) cells are *Neisseria gonorrhoea* cells, eg, resistant to an antibiotic selected from cefixime (eg, an oral cephalosporin), ceftriaxone (an injectable cephalosporin), azithromycin and tetracycline.

In an example, the first (host) cells are *Acinetobacter baumannii* cells, eg, resistant to an antibiotic selected from beta-lactam, meropenem and a carbapenem.

In an example, the first (host) cells are *Campylobacter* cells, eg, resistant to an antibiotic selected from ciprofloxacin and azithromycin.

In an example, the second species or strain is a human gut commensal species or strain and/or a human gut probiotic species or strain.

In an example, the second species or strain is a Bacteroidetes (eg, *Bacteroides*) and optionally the host cells are gram positive bacterial cells.

In an example, the first cells are *Firmicutes* cells.

In an example, causing said dysbiosis or step (b) is carried out by targeting the sub-population of first cells by administering thereto an anti-bacterial or anti-archaeal agent simultaneously or sequentially with said immune cell population, whereby first cells are killed or the sub-population growth is reduced, thereby reducing the proportion of said sub-population in the gut microbiota of the patient.

In an example, the method reduces first (host) cell population growth by at least 5, 10, 20, 50 or 100-fold compared to the growth of a control population of host cells that have not received said guided nuclease (eg, Cas) modification.

In an example, the method inhibits host cell population growth on a gut surface.

In an example, for each host cell the system comprises components according to (i) to (iv):—

- (i) at least one nucleic acid sequence encoding a Cas nuclease;
- (ii) an engineered HM-CRISPR array comprising a spacer sequence and repeats encoding a HM-crRNA, the HM-crRNA comprising a sequence that hybridises to a host cell target sequence to guide said Cas to the target in the host cell to modify the target sequence;

(iii) an optional tracrRNA sequence or a DNA sequence expressing a tracrRNA sequence;

(iv) wherein said components of the system are split between the host cell and at least one nucleic acid vector that transforms the host cell, whereby the HM-crRNA guides Cas to the target to modify the host target sequence in the host cell; and

wherein the target sequence is modified by the Cas whereby the host cell is killed or host cell growth is reduced;

the method comprising introducing the vectors of (iv) into host cells and expressing said HM-crRNA in the host cells, allowing HM-crRNA to hybridise to host cell target sequences to guide Cas to the targets in the host cells to modify target sequences, whereby host cells are killed or host cell growth is reduced, thereby altering the proportion of said sub-population in the microbiota.

In an example, component (i) is endogenous to each host cell.

In an example, each vector is a virus or phage.

In an example, each target sequence is adjacent a NNA-GAAW or NGGNG protospacer adjacent motif (PAM).

In an example, alternatively HM-crRNA and tracrRNA are comprised by a single guide RNA (gRNA), the method comprising introducing said gRNA into host cells or expressing the gRNA in host cells.

In an example, the microbiota comprises a second bacterial or archaeal species, wherein each of the first and second species is a respective species of the same phylum (eg, both *Firmicutes* species) and the growth of the second bacteria is not inhibited by the HM-system; or wherein the microbiota comprises a second bacterial or archaeal strain, wherein each of the first and second bacteria or archaea is a respective strain of the same species and the growth of the second bacteria or archaea is not inhibited by the HM-system.

In an example, the microbiota comprises a second bacterial species, wherein each of the first and second species is a respective gram-positive species and the growth of the second bacteria is not inhibited by the HM-system.

In an example, each target sequence is comprised by an antibiotic resistance gene, virulence gene or essential gene of the host cell.

In an example, each first cell is a *Staphylococcus*, *Streptococcus*, *Pseudomonas*, *Salmonella*, *Listeria*, *E coli*, *Desulfovibrio*, *Vibrio* or *Clostridium* cell.

In an example, the dysbiosis or step (b) comprises stimulating Paneth cells of the patient by gut *Bacteroides* (eg, *B thetaiotamicron*), wherein the altered microbiota produced by the dysbiosis or step (b) comprises an increased proportion of *Bacteroides* first bacteria compared with the microbiota before the dysbiosis or step (b), whereby Paneth cells are stimulated and the cell therapy is modulated.

Bacteroides can affect expression of Paneth cell proteins. Small intestinal crypts house stem cells that serve to constantly replenish epithelial cells that die and are lost from the villi. Paneth cells (immune systems cells similar to neutrophils), located adjacent to these stem cells, protect them against microbes by secreting a number of antimicrobial molecules (defensins) into the lumen of the crypt, and it is possible that their protective effect even extends to the mature cells that have migrated onto the villi. In animal models, *B. thetaiotaomicron* can stimulate production of an antibiotic Paneth cell protein (Ang4) that can kill certain pathogenic organisms (e.g., *Listeria monocytogenes*). *Listeria monocytogenes* is a strong T_H1 cell inducer, and thus by stimulating Paneth cells in certain aspects of the invention, this may be useful to skew immunity from TH1 to other cell

types, such as T_H17 . This may be useful for increasing or enhancing the immune cell therapy of the invention. For example, this may be useful when the invention comprises administering T_H17 -based cell therapy (eg, CAR- T_H17 cells) to the patient.

In an example, the dysbiosis or step (b) comprises developing an immune response in the patient to gut *Bacteroides* (eg, *B thetaiotamicron*), wherein the altered microbiota produced by the dysbiosis or step (b) comprises an increased proportion of *Bacteroides* first bacteria compared with the microbiota before the dysbiosis or step (b), whereby the cell therapy is modulated.

In an example, the dysbiosis or step (b) comprises altering the relative proportion of a or said sub-population of first cells in the gut microbiota of the patient, thereby producing an altered gut microbiota that modulates the immune cell therapy in the patient, wherein the dysbiosis or step (b) comprises killing first cells of said sub-population or inhibiting growth of said sub-population by using guided nuclease targeting to the genome of first cells comprised by the sub-population.

The invention also provides: A bacterial or archaeal transplant for administration to a patient for therapy of a disease or condition in the patient using the method of the invention. Optionally the transplant comprises cells of said first species. Optionally the transplant comprises cells of said second species (and eg, does not comprise cells of the first species).

The invention also provides: A bacterial or archaeal transplant for administration to a subject (eg, a plant or yeast) or to soil or an environment for modulating a treatment thereof using the method of the invention. Optionally the transplant comprises cells of said first species. Optionally the transplant comprises cells of said second species (and eg, does not comprise cells of the first species).

In an example, the yeast comprises a population of yeast cells of the same species or strain. For example, the yeast is comprised by the microbiota, eg, the yeast comprises a sub-population of cells of the microbiota, such as where the targeted cells are bacteria or archaea cells also comprised by the microbiota. In an embodiment, the method kills the targeted cells (eg, bacteria) or reduces growth of the target cell sub-population comprised by the microbiota, wherein release of (or the concentration of) one or more chemicals or messengers by target cells in the microbiota is reduced. For example, the one or more chemicals or messengers that mediate quorum sensing in microbes (eg, bacteria) of the microbiota is reduced. This may be useful to shape the growth of target cell and/or other microbial cell populations in the microbiota. In an example, the one or more chemicals or messengers inhibit growth of the yeast or kill the yeast, and thus reduction of the release or concentration of the chemical or messenger(s) in the microbiota is advantageous to promote growth and/or maintenance of the yeast in the microbiota. In an embodiment of these examples, the treatment is nutrition of the yeast or treatment of the yeast with a growth promoter (eg, wherein the yeast are for foodstuff or beverage production, or for production of a biotechnology or medical product, such as an antibody eg, the yeast is *Saccharaomyces*). In another example, the chemical or messenger(s) promote yeast growth, wherein the yeast treatment is treatment with a yeast killing agent (eg, an fungicide). This is useful to promote efficacy of the killing treatment when the yeast are undesirable (eg, an undesirable mould). In an example when the subject is a plant, the one or more chemicals or messengers inhibit growth of the plant or kill the plant, and thus reduction of the release or

concentration of the chemical or messenger(s) in the microbiota is advantageous to promote growth of the plant and/or inhibit killing of the plant. In an embodiment of these examples, the treatment is nutrition of the plant or treatment of the yeast with a fertilizer or nitrogen fixing agent. In an embodiment, the treatment is a pesticide treatment of the plant, such as a treatment that is inhibited or reduced in the presence of the targeted microbiota cells.

In an example, the plant is a crop plant (eg, wheat, barley, cereal or livestock fodder plant), fruit plant (eg, apple, orange, citrus, lime, lemon, raspberry, strawberry, berry or banana plant), legume plant, sugar cane plant, spice plant, garden plant, vegetable plant, grass, tree or flowering plant. For example, the plant is a tuber plant, eg, a potato or a sweet potato (eg, and the first, host or targeted bacteria are *Pectobacterium atrosepticum* cells, optionally wherein the treatment is storage (eg, cold storage), washing, a pesticide or herbicide treatment, fertilising or hydrating of the plant of a crop thereof (eg, a potato crop)). For example, the plant is a tobacco plant (eg, and the first, host or targeted bacteria are *Ralstonia solanacearum* cells, optionally wherein the treatment is storage (eg, cold storage), washing, a pesticide or herbicide treatment, fertilising or hydrating of the plant of a crop thereof (eg, a tobacco leaf crop)).

In an example, the subject is a protozoa. In an example, the subject or patient is a fish. In an example, the subject or patient is a bird. In an example, the subject or patient is a reptile. In an example, the subject or patient is an arachnid. In an example, the subject is a yeast cell (eg, a *Saccharomyces* cell). In an example, the subject or patient is an animal (eg, a rodent, mouse or rat). In an example, the subject or patient is a human (eg, an embryo or not an embryo). In an example, the subject or patient is a companion pet animal (eg, a bird, horse, dog, cat or rabbit). In an example, the subject or patient is an insect (an insect at any stage of its lifecycle, eg, egg, larva or pupa, eg, a fly or crawling insect or a beetle). In an example, the subject or patient is a cephalopod or nematode. In an example, the subject or patient is a plant or animal pest. In an example, the treatment of an animal or human may be a nutritional treatment, therapy of a disease or condition, prophylaxis of a disease or condition, ocular treatment, pesticide treatment, dental treatment, topical treatment or digestion treatment. In an example, the method enhances immunity of the subject or patient against a pathogen (eg a pathogenic parasite, protozoan, virus or bacteria). In an example, the treatment or therapy is a combination treatment or therapy practised on the human or animal, wherein the human or animal is administered a first medicament in combination with a second medicament or radiation. The first and/or second medicament may be an antibody therapy or immune cell transplant (eg, CAR-T or TILs transplant) therapy (and the immune modulation aspect of the invention may be advantageous for modulating such therapies). In an example, each medicament or treatment is selected from Table 2.

In an example, the disease or condition is diabetes (eg, Type I or II diabetes, insulin-resistant diabetes (eg, insulin-resistant Type II diabetes), onset of insulin resistance in a diabetic or pre-diabetic patient or reduction in insulin responsiveness in a diabetic or pre-diabetic patient. Optionally additionally in this example, the first or host cells that are targeted are *Prevotella copri* or *Bacteroides vulgatus* cells. In an embodiment, both *Prevotella copri* and *Bacteroides vulgatus* cells are targeted (ie, killed and/or population growth reduced, or reduced in the microbiota following

administration of a transplant as described herein) in the patient, wherein said disease or condition is treated or prevented.

The invention also provides: the HM-CRISPR/Cas system, HM-array or HM-crRNA for administration to a patient for therapy of a disease or condition in the patient using the method of the invention.

The invention also provides: A kit comprising an ex vivo population of immune cells for adoptive cell therapy of a patient, wherein the kit further comprises the transplant, system, array or crRNA, optionally wherein the immune cells are selected from CAR-T cells, T-cells expressing engineered T-cell receptors (TCRs), tumour infiltrating lymphocytes (TILs) and NK cells.

Mobile Genetic Elements (MGEs)

In an example, each vector is a nucleic acid vector comprising or consisting of a mobile genetic element (MGE), wherein the MGE comprises an origin of transfer (oriT) and a CRISPR array for modifying a target sequence of the genome of a host cell or the genome of a virus (eg, prophage) in a host cell. Examples of MGEs are ICEs, transposons, plasmids and bacteriophage. An origin of transfer (oriT) is a short sequence (eg, up to 500 bp) that is necessary for transfer of the DNA that contains it from a bacterial host to recipient during conjugation. The oriT is cis-acting—it is found on the same DNA that is being transferred, and it is transferred along with the DNA. A typical origin of transfer comprises three functionally defined domains: a nicking domain, a transfer domain, and a termination domain.

Reference is made to the ICEberg database (<http://db-mml.sjtu.edu.cn/ICEberg/>), which provides examples of suitable ICEs for the invention and sources for suitable oriT. In an example, the ICE is a member of an ICE family comprising an ICE selected from the group 1 to 28, or the oriT is an oriT of a member of such a family: 1=SXT/R391; 2=Tn916; 3=Tn4371; 4=CTnDOT/ERL; 5=ICEclc; 6=ICEBs1; 7=ICEHin1056; 8=PAPI-1; 9=ICEM1Sym(R7A); 10=ICEST1; 11=SPI-7; 12=ICE6013; 13=ICEKp1; 14=TnGBS1; 15=Tn5253; 16=ICESa2603; 17=ICEYel; 18=10270-RD.2; 19=Tn1207.3; 20=Tn1806; 21=ICEA5632; 22=ICEF-I/II; 23=ICEAPG2; 24=ICEM; 25=10270-RD.1; 26=Tn5801; 27=PPI-1; 28=ICEF-III. Family descriptions are found in the ICEberg database. For example, the Tn916 family was defined by Roberts et al (2009) (Trends Microbiol. 2009 June; 17(6):251-8. doi: 10.1016/j.tim.2009.03.002. Epub 2009 May 20; “A modular master on the move: the Tn916 family of mobile genetic elements”, Roberts A, Mullany P). Elements belonging to the Tn916 family are defined by the following criteria: they must have the general organization shown in Roberts et al, and they must have a core region (conjugation and regulation module) that is similar in sequence and structure to the original Tn916 at the DNA level. Exceptions are some conjugative transposons, such as Tn1549 which have been previously classified in this family and those with a high degree of protein similarity as described in corresponding references. Optionally, the ICE is a transposon, eg, a conjugative transposon. In an example, the MGE is a mobilisable transposon that is mobilisable in the presence of a functional helper element, optionally wherein the transposon is in combination with a said helper element.

Optionally the vector is a plasmid, optionally wherein the MGE is a transposon comprised by the plasmid. For example, the transposon is a conjugative transposon. In an example the transposon is a mobilisable transposon (eg, mobilisable using one or more factors encoded by the

plasmid, eg, by genes outside the transposon sequence of the plasmid). Optionally, the transposon is a Type I transposon. Optionally, the transposon is a Type II transposon. Optionally, the vector oriT is an oriT of a Bacteroidetes (eg, Bacteroidales or *Bacteroides*) or *Prevotella* transposon. This is useful when the first (host) cells are Bacteroidetes (eg, Bacteroidales or *Bacteroides*) or *Prevotella* respectively. Optionally, the vector oriT is a CTnDot, CTnERL SXT/R391, Tn916 or Tn4371 family transposon oriT.

Optionally, the method comprises exposing the patient's microbiota to a vector or MGE (eg, one described above) that comprises a toxin-antitoxin module that comprises an anti-toxin gene that is operable in the second bacteria, but is not operable or has reduced operation in first (host) cells. Thus, first cells are killed and second bacteria are spared, thereby altering the proportion of first cells in the patient's microbiota.

Split CRISPR/Cas System

In one aspect, endogenous Cas of the first (host) cells is harnessed and operates with the engineered sequences comprised by vectors (eg, phage) that are introduced into host cells. This aspect is advantageous to free up space in vectors, for example viruses or phage that have restricted capacity for carrying exogenous sequence. By freeing up space, one is able to include more targeting spacers or arrays, which is useful for evading host resistance. It is advantageous, for example to harness the endogenous Cas endonuclease rather than encode it in the vector—especially for bulky Cas sequences such as sp or saCas9. Additionally, there is no chance of inferior compatibility as may be seen with some exogenous Cas from non-host sources. The ability to reduce virus, eg, phage genome size, may also be beneficial for promoting host cell uptake (infection and/or maintenance of the virus in host cells). In some examples, an advantage is that invasion of the host by the vector (eg, phage) may upregulate host CRISPR/Cas activity, including increased expression of host Cas nucleases—in an attempt of the host to combat invading nucleic acid. This, however, is also useful to provide endogenous Cas for use with the invention when these use crRNA or gRNA that are recognised by the host Cas. In the case where the invention involves crRNA or gRNA targeting a host CRISPR array, this then promotes inactivation of the host CRISPR array itself, akin to a “suicidal” host cell which then uses its own Cas nuclease to inactivate its own CRISPR systems.

Thus, the vectors may lack a Cas nuclease (eg, aCas9)-encoding sequence.

Optionally, the endogenous first (host) cell system is a CRISPR/Cas9 system. Optionally, the nuclease is a Type I Cas nuclease. Optionally, the nuclease is a Type II Cas nuclease (eg, a Cas9). Optionally, the nuclease is a Type III Cas nuclease.

To save even more space, optionally a tracrRNA sequence is not provided by the vectors, but is a tracrRNA sequence of an endogenous host cell CRISPR/Cas system, wherein the tracrRNA is capable of hybridising with the HM-crRNA in the cell for subsequent processing into mature crRNA for guiding Cas to the target in the host cell.

Generally Applicable Features

The following features apply to any configuration (eg, in any of its aspects, embodiments, concepts, paragraphs or examples) of the invention:—

In an example, the target sequence is a chromosomal sequence, an endogenous host cell sequence, a wild-type host cell sequence, a non-viral chromosomal host cell sequence and/or a non-phage sequence (ie, one more or all of these), eg, the sequence is a wild-type host chromosomal

cell sequence such as antibiotic resistance gene or essential gene sequence comprised by a host cell chromosome. In an example, the sequence is a host cell plasmid sequence, eg, an antibiotic resistance gene sequence.

In an example, at least two target sequences are modified by Cas, for example an antibiotic resistance gene and an essential gene. Multiple targeting in this way may be useful to reduce evolution of escape mutant host cells.

In an example, the Cas is a wild-type endogenous host cell Cas nuclease. In an example, each host cell has constitutive Cas nuclease activity, eg, constitutive wild-type Cas nuclease activity. In an example, the host cell is a bacterial cell; in another example the host cell is an archaeal cell. Use of an endogenous Cas is advantageous as this enables space to be freed in vectors cRNA or gRNA. For example, Type II Cas9 nucleotide sequence is large and the use of endogenous Cas of the host cell instead is advantageous in that instance when a Type II CRISPR/Cas system is used for host cell modification in the present invention. The most commonly employed Cas9, measuring in at 4.2 kilobases (kb), comes from *S pyogenes*. While it is an efficient nuclease, the molecule's length pushes the limit of how much genetic material a vector can accommodate, creating a barrier to using CRISPR in the tissues of living animals and other settings described herein (see F. A. Ran et al., "In vivo genome editing using *Staphylococcus aureus* Cas9," Nature, doi:10.1038/nature14299, 2015). Thus, in an embodiment, the vector of the invention is a AAV vector or has an exogenous DNA insertion capacity no more than an AAV vector, and the Cas is an endogenous Cas of the host cell, wherein the cell is a bacterial or archaeal cell. *S thermophilus* Cas9 (UniProtKB-G3EGR1 (CAS9_STRTR)) nucleotide sequence has a size of 1.4 kb.

In an example, the vector is a viral vector. Viral vectors have a particularly limited capacity for exogenous DNA insertion, thus virus packaging capacity needs to be considered. Room needs to be left for sequences encoding vital viral functions, such as for expressing coat proteins and polymerase. In an example, the vector is a phage vector or an AAV or lentiviral vector. Phage vectors are useful where the host is a bacterial cell.

By use of the term "engineered" it will be readily apparent to the skilled addressee that the array, sequence, vector, cRNA, gRNA, MGE or any other configuration, concept, aspect, embodiment, paragraph or example etc of the invention is non-naturally occurring. For example, the MGE, vector, sequence or array comprises one or more sequences or components not naturally found together with other sequences or components of the MGE, vector, sequence or array. For example, the array or sequence is recombinant, artificial, synthetic or exogenous (ie, non-endogenous or not wild-type) to the or each host cell.

In an example, the array or sequence of the invention is an engineered version of an array or sequence isolated, for example isolated from a host cell. In an example, the engineered array or sequence is not in combination with a Cas endonuclease-encoding sequence that is naturally found in a cell.

Studies suggest that *Bacteroides* have a role in preventing infection with *Clostridium difficile*. The development of the immune response that limits entry and proliferation of potential pathogens is profoundly dependent upon *B fragilis*. Also, Paneth cell proteins may produce antibacterial peptides in response to stimulation by *B thetaiotomicron*, and these molecules may prevent pathogens from colonizing the gut. In addition, *B thetaiotomicron* can induce Paneth cells to produce a bactericidal lectin, RegIII, which exerts its

antimicrobial effect by binding to the peptidoglycan of gram-positive organisms. Thus, the use of the invention in any of its configurations for increasing the proportion of *Bacteroides* (eg, *thetaiotomicron* and/or *fragilis*) in the patient's microbiota is useful for limiting pathogenic bacterial colonisation of the population or a gut of a human or non-human animal.

Hooper et al demonstrated that *B thetaiotomicron* can modify intestinal fucosylation in a complex interaction mediated by a fucose repressor gene and a signaling system. Using transcriptional analysis it was demonstrated that *B thetaiotomicron* can modulate expression of a variety of host genes, including those involved in nutrient absorption, mucosal barrier fortification, and production of angiogenic factors.

Optionally, the host (or first and/or second bacteria) is a gram negative bacterium (eg, a spirilla or vibrio). Optionally, the host (or first and/or second bacteria) is a gram positive bacterium. Optionally, the host (or first and/or second bacteria) is a mycoplasma, chlamydiae, spirochete or *Mycobacterium*. Optionally, the host (or first and/or second bacteria) is a *Streptococcus* (eg, *pyogenes* or *thermophilus*) host. Optionally, the host (or first and/or second bacteria) is a *Staphylococcus* (eg, *aureus*, eg, MRSA) host. Optionally, the host (or first and/or second bacteria) is an *E. coli* (eg, O157: H7) host, eg, wherein the Cas is encoded by the vector or an endogenous host Cas nuclease activity is de-repressed. Optionally, the host (or first and/or second bacteria) is a *Pseudomonas* (eg, *aeruginosa*) host. Optionally, the host (or first and/or second bacteria) is a *Vibro* (eg, *cholerae* (eg, O139) or *vulnificus*) host. Optionally, the host (or first and/or second bacteria) is a *Neisseria* (eg, *gonorrhoeae* or *meningitidis*) host. Optionally, the host (or first and/or second bacteria) is a *Bordetella* (eg, *pertussis*) host. Optionally, the host (or first and/or second bacteria) is a *Haemophilus* (eg, *influenzae*) host. Optionally, the host (or first and/or second bacteria) is a *Shigella* (eg, *dysenteriae*) host. Optionally, the host (or first and/or second bacteria) is a *Brucella* (eg, *abortus*) host. Optionally, the host (or first and/or second bacteria) is a *Francisella* host. Optionally, the host (or first and/or second bacteria) is a *Xanthomonas* host. Optionally, the host (or first and/or second bacteria) is a *Agrobacterium* host. Optionally, the host (or first and/or second bacteria) is a *Erwinia* host. Optionally, the host (or first and/or second bacteria) is a *Legionella* (eg, *pneumophila*) host. Optionally, the host (or first and/or second bacteria) is a *Listeria* (eg, *monocytogenes*) host. Optionally, the host (or first and/or second bacteria) is a *Campylobacter* (eg, *jejuni*) host. Optionally, the host (or first and/or second bacteria) is a *Yersinia* (eg, *pestis*) host. Optionally, the host (or first and/or second bacteria) is a *Borelia* (eg, *burgdorferi*) host. Optionally, the host (or first and/or second bacteria) is a *Helicobacter* (eg, *pylori*) host. Optionally, the host (or first and/or second bacteria) is a *Clostridium* (eg, *difficile* or *botulinum*) host. Optionally, the host (or first and/or second bacteria) is a *Erllichia* (eg, *chaffeensis*) host. Optionally, the host (or first and/or second bacteria) is a *Salmonella* (eg, *typhi* or *enterica*, eg, serotype *typhimurium*, eg, DT 104) host. Optionally, the host (or first and/or second bacteria) is a *Chlamydia* (eg, *pneumoniae*) host. Optionally, the host (or first and/or second bacteria) is a *Parachlamydia* host. Optionally, the host (or first and/or second bacteria) is a *Corynebacterium* (eg, *amycolatum*) host. Optionally, the host (or first and/or second bacteria) is a *Klebsiella* (eg, *pneumoniae*) host. Optionally, the host (or first and/or second bacteria) is a *Enterococcus* (eg, *faecalis* or *faecium*, eg,

linezolid-resistant) host. Optionally, the host (or first and/or second bacteria) is a *Acinetobacter* (eg, *baumannii*, eg, multiple drug resistant) host.

A tracrRNA sequence may be omitted from an array or vector of the invention, for example for Cas systems of a Type that does not use tracrRNA.

In an example, the Cas guided to the target is an exonuclease. Optionally a nickase as mentioned herein is a double nickase. An example of a nickase is a Cas9 nickase, ie, a Cas9 that has one of the two nuclease domains inactivated—either the RuvC and/or HNH domain.

Mention herein of using vector DNA can also in an alternative embodiment apply mutatis mutandis to vector RNA where the context allows. For example, where the vector is an RNA vector. All features of the invention are therefore in the alternative disclosed and to be read as “RNA” instead of “DNA” when referring to vector DNA herein when the context allows. In an example, the or each vector also encodes a reverse transcriptase.

In an example, the or each array or engineered nucleotide sequence is provided by a nanoparticle vector or in liposomes.

In an example, the Cas is a Cas nuclease for cutting, dead Cas (dCas) for interrupting or a dCas conjugated to a transcription activator for activating the target.

In an example, the or each array or engineered sequence comprises an exogenous promoter functional for transcription of the crRNA or gRNA in the host.

In an embodiment the array or engineered sequence is contained in a virophage vector and the host is alternatively a virus which can infect a cell. For example, the host is a large virus that may have infected an amoeba cell. For example, the host is a Sputnik virus, Pithovirus, mimivirus, mamavirus, Megavirus or Pandoravirus, eg, wherein the host virus is in water. In an example of this embodiment, the invention is for water or sewage treatment (eg, purification, eg, waterway, river, lake, pond or sea treatment).

In an embodiment the or each vector or engineered sequence is or is comprised by a Φ NM1 phage, eg, wherein the host cell(s) is a *S. aureus* (eg, MRSA) cell.

For example the method is practised on a mammalian subject, eg, a human, rodent, mouse or rat. For example the method is practised on a vertebrate, reptile, bird or fish.

The cell population can be administered to the patient in one or more doses. For example, the method comprises administering an antibacterial agent to cause said dysbiosis, or administering a bacterial transplant to the patient to cause said dysbiosis.

Wherein the method reduces the cell therapy, the therapy can be downregulated, dampened or switched off, eg, to reduce or prevent an unwanted side-effect of the cell therapy (eg, a CAR-T therapy side effect in a human patient, such as CRS). Wherein the method increases the cell therapy, the therapy can be upregulated, enhance or switched on, eg, to enhance cytotoxicity against one target cells.

The method treats or prevents (ie, reduces the risk of) the disease or condition. This may be complete or partial treatment or prevention, ie, a reduction but not complete reduction of the disease/condition or symptoms thereof; or a reducing of the risk but not total prevention of the disease/condition or a symptom thereof. Similarly, the method treats or prevents (ie, reduces the risk of) an undesirable symptom of the disease or condition or the therapy (eg, CRS).

Concepts:

1. A method of modulating an adoptive immune cell therapy of a disease or condition in a patient, the method comprising

- a. Carrying out adoptive immune cell therapy in the patient, comprising administering a population of immune cells to the patient, wherein administration of said immune cells is capable of treating the disease or condition in the patient; and
 - b. Causing bacterial microbiota (eg, gut microbiota) dysbiosis in the patient, whereby said dysbiosis modulates the immune cell therapy in the patient.
2. A method of modulating an adoptive immune cell therapy of a disease or condition in a patient, the method comprising
 - a. Carrying out adoptive immune cell therapy in the patient, comprising administering a population of immune cells to the patient, wherein administration of said immune cells is capable of treating the disease or condition in the patient; and
 - b. Altering the relative proportion of a sub-population of cells of a first bacterial species or strain, or archaeal species or strain, in a microbiota (eg, gut microbiota) of the patient, thereby producing an altered microbiota that modulates the immune cell therapy in the patient.
 3. The method of concept 2, wherein step (b) is carried out by targeting the sub-population of first cells by administering thereto an anti-bacterial or anti-archaeal agent simultaneously or sequentially with said immune cell population, whereby first cells are killed or the sub-population growth is reduced, thereby reducing the proportion of said sub-population in the microbiota of the patient.
 4. The method of any preceding concept, wherein the cell therapy is an adoptive T-cell therapy.
 5. The method of concept 4, wherein cells selected from the group consisting of CD4⁺ T-cells, CD8⁺ T-cells, T_H1 cells and T_H17 cells are administered to the patient in step (a).
 6. The method of concept 4 or 5, wherein T_H17 cells are modulated in the patient.
 7. The method of concept 4, 5 or 6, wherein T_{reg} cells are modulated in the patient.
 8. The method of any preceding concept, wherein the cell therapy is enhanced.
 9. The method of any preceding concept wherein T_H17 cells are upregulated in the patient and/or T_{reg} cells are downregulated in the patient.
 10. The method of any preceding concept, wherein CD4⁺ T-cells are upregulated in the patient.
 11. The method of any preceding concept, wherein CD8⁺ T-cells are upregulated in the patient.
 12. The method of any preceding concept, wherein one or more of central memory T cells (T_{CM}), effector memory T cells (T_{EM}), stem cell memory cells (T_{SCM}) and effector cells (T_{eff}) are upregulated in the patient, wherein the cells are comprised by the immune cell population administered in step (a) and/or are progeny thereof.
 13. The method of any one of concepts 1 to 7, wherein the cell therapy is reduced.
 14. The method of concept 13, wherein the method reduces or prevents the risk of cytokine release syndrome (CRS) in the patient.
 15. The method of any one of concepts 1 to 7, 13 and 14, wherein, T_H17 cells are downregulated in the patient and/or T_{reg} cells are upregulated in the patient.
 16. The method of any one of concepts 1 to 7 and 13 to 15, wherein CD4⁺ T-cells are downregulated in the patient.
 17. The method of any one of concepts 1 to 7 and 13 to 16, wherein CD8⁺ T-cells are downregulated in the patient.
 18. The method of any one of concepts 1 to 7 and 13 to 17, wherein one or more of central memory T cells (T_{CM}),

- effector memory T cells (T_{EM}), stem cell memory cells (T_{SCM}) and effector cells (T_{eff}) are downregulated in the patient, wherein the cells are comprised by the immune cell population administered in step (a) and/or are progeny thereof.
19. The method of any preceding concept, wherein the immune cell population comprises CAR-T cells and/or T-cells expressing engineered T-cell receptors (TCRs) and/or tumour infiltrating lymphocytes (TILs).
20. The method of any preceding concept, wherein the immune cell population comprises engineered autologous or allogeneic immune cells, eg, T-cells, NK cells and/or TILs.
21. The method of concept 19 or 20, wherein the T-cells are $CD4^+$ T-cells or T_H17 cells.
22. The method of any preceding concept, wherein step (b) increases the proportion of *Bacteroides* (eg, *B fragalis* and/or *B thetaiotamicron*) in the microbiota.
23. The method of any one of concepts 1 to 21, wherein step (b) decreases the proportion of *Bacteroides* (eg, *B fragalis* and/or *B thetaiotamicron*) in the microbiota.
24. The method of any preceding concept, wherein step (b) increases the proportion of Bacteroidetes to *Firmicutes* in the microbiota.
25. The method of any one of concepts 1 to 23, wherein step (b) decreases the proportion of Bacteroidetes to *Firmicutes* in the microbiota.
26. The method of any preceding concept, wherein step (b) reduces the proportion of one or more *Clostridium* species or strain (eg, wherein each species is a cluster IV or XIVa *Clostridium* species) in the microbiota.
27. The method of any one of concepts 1 to 25, wherein step (b) increases the proportion of one or more *Clostridium* species or strain (eg, wherein each species is a cluster IV or XIVa *Clostridium* species) in the microbiota.
28. The method of any preceding concept, wherein step (b) reduces the proportion of *Bifidobacterium* (eg, *B bifidum*) in the microbiota.
29. The method of any one of concepts 1 to 27, wherein step (b) increases the proportion of *Bifidobacterium* (eg, *B bifidum*) in the microbiota.
30. The method of any preceding concept, wherein step (b) comprises altering the relative proportion of a or said sub-population of first cells in the microbiota of the patient, thereby producing an altered microbiota that modulates the immune cell therapy in the patient, wherein the sub-population comprises host cells of said first species or strain, the method comprising
- combining the microbiota with multiple copies of engineered nucleic acid sequences encoding host modifying (HM) crRNAs, and
 - expressing HM-crRNAs in host cells,
- wherein each engineered nucleic acid sequence is operable with a Cas nuclease in a respective host cell to form a HM-CRISPR/Cas system and the engineered sequence comprises
- spacer and repeat sequences encoding a HM-crRNA;
 - the HM-crRNA comprising a sequence that is capable of hybridizing to a host cell target sequence to guide Cas nuclease to the target sequence in the host cell; and optionally the HM-system comprises a tracrRNA sequence or a DNA sequence expressing a tracrRNA sequence;
- whereby HM-crRNAs guide Cas modification of host target sequences in host cells, whereby host cells are killed or the host cell population growth is reduced, thereby reducing the proportion of said sub-population in the microbiota.

31. The method of concept 30, comprising using endogenous Cas nuclease of host cells for modification of target nucleotide sequences.
32. The method of concept 30 or 31, comprising reducing host cell population growth by at least 5-fold compared to the growth of a control population of host cells that have not received said Cas modification.
33. The method of any one of concepts 30 to 32, comprising inhibiting host cell population growth on a gut surface.
34. The method of any one of concepts 30 to 33, wherein the microbiota comprises cells of a second bacterial species or strain, or archaeal species or strain, wherein the second species or strain has a 16s ribosomal RNA-encoding DNA sequence that is at least 80% identical to an 16s ribosomal RNA-encoding DNA sequence of the host cell species or strain, wherein the growth of the second cells in the microbiota is not inhibited by said HM-system.
35. The method of concept 34, wherein the second species or strain is a human gut commensal species or strain and/or a human gut probiotic species or strain.
36. The method of concept 34 or 35, wherein the second species or strain is a Bacteroidetes (eg, *Bacteroides*) and optionally the host cells are gram positive bacterial cells.
37. The method of any one of concepts 30 to 36, wherein the first cells are *Firmicutes* cells.
38. The method of any one of concepts 30 to 37, wherein for each host cell the system comprises components according to (i) to (iv):—
- at least one nucleic acid sequence encoding a Cas nuclease;
 - an engineered HM-CRISPR array comprising a spacer sequence and repeats encoding a HM-crRNA, the HM-crRNA comprising a sequence that hybridises to a host cell target sequence to guide said Cas to the target in the host cell to modify the target sequence;
 - an optional tracrRNA sequence or a DNA sequence expressing a tracrRNA sequence;
 - wherein said components of the system are split between the host cell and at least one nucleic acid vector that transforms the host cell, whereby the HM-crRNA guides Cas to the target to modify the host target sequence in the host cell; and
- wherein the target sequence is modified by the Cas whereby the host cell is killed or host cell growth is reduced;
- the method comprising introducing the vectors of (iv) into host cells and expressing said HM-crRNA in the host cells, allowing HM-crRNA to hybridise to host cell target sequences to guide Cas to the targets in the host cells to modify target sequences, whereby host cells are killed or host cell growth is reduced, thereby altering the proportion of said sub-population in the microbiota.
39. The method of concept 38, wherein component (i) is endogenous to each host cell.
40. The method of concept 38 or 39, wherein each vector is a virus or phage.
41. The method of any one of concepts 30 to 40, wherein each target sequence is adjacent a NNAGAAW or NGGNG protospacer adjacent motif (PAM).
42. The method of any one of concepts 30 to 41, wherein alternatively HM-crRNA and tracrRNA are comprised by a single guide RNA (gRNA), the method comprising introducing said gRNA into host cells or expressing the gRNA in host cells.
43. The method of any one of concepts 30 to 35 and 37 to 42, wherein the microbiota comprises a second bacterial or archaeal species, wherein each of the first and second

- species is a respective species of the same phylum (eg, both *Firmicutes* species) and the growth of the second bacteria is not inhibited by the HM-system; or wherein the microbiota comprises a second bacterial or archaeal strain, wherein each of the first and second bacteria or archaea is a respective strain of the same species and the growth of the second bacteria or archaea is not inhibited by the HM-system.
44. The method of any one of concepts 30 to 43, wherein the microbiota comprises a second bacterial species, wherein each of the first and second species is a respective gram-positive species and the growth of the second bacteria is not inhibited by the HM-system.
45. The method of any one of concepts 30 to 44, wherein each target sequence is comprised by an antibiotic resistance gene, virulence gene or essential gene of the host cell.
46. The method of any preceding concept, wherein each first cell is a *Staphylococcus*, *Streptococcus*, *Pseudomonas*, *Salmonella*, *Listeria*, *E coli*, *Desulfovibrio*, *Vibrio* or *Clostridium* cell.
47. The method of any preceding concept, wherein step (b) comprises stimulating Paneth cells of the patient by gut *Bacteroides* (eg, *B thetaiotamicron*), wherein the altered microbiota produced by step (b) comprises an increased proportion of *Bacteroides* first bacteria compared with the microbiota before step (b), whereby Paneth cells are stimulated and the cell therapy is modulated.
48. The method of any preceding concept, wherein step (b) comprises developing an immune response in the patient to gut *Bacteroides* (eg, *B thetaiotamicron*), wherein the altered microbiota produced by step (b) comprises an increased proportion of *Bacteroides* first bacteria compared with the microbiota before step (b), whereby the cell therapy is modulated.
49. The method of any preceding concept, wherein step (b) comprises altering the relative proportion of a or said sub-population of first cells in the microbiota of the patient, thereby producing an altered microbiota that modulates the immune cell therapy in the patient, wherein step (b) comprises killing first cells of said sub-population or inhibiting growth of said sub-population by using guided nuclease targeting to the genome of first cells comprised by the sub-population.
50. A bacterial or archaeal transplant for administration to a patient for therapy of a disease or condition in the patient using the method of any preceding concept, optionally wherein the transplant comprises cells of said first species.
51. A HM-CRISPR/Cas system, HM-array or HM-crRNA as recited in any one of concepts 30 to 45 for administration to a patient for therapy of a disease or condition in the patient using the method of any one of concepts 1 to 49.
52. A kit comprising an ex vivo population of immune cells for adoptive cell therapy of a patient, wherein the kit further comprises a transplant, system, array or crRNA of concept 50 or 51, optionally wherein the immune cells are selected from CAR-T cells, T-cells expressing engineered T-cell receptors (TCRs), tumour infiltrating lymphocytes (TILs) and NK cells.
- Aspects:
1. An ex vivo population of immune cells for use in a method of adoptive cell therapy of a patient for treating or preventing a disease or condition in the patient, the method comprising
 - a. Carrying out adoptive immune cell therapy in the patient, comprising administering cells of said popula-

- tion to the patient, wherein administration of said immune cells is capable of treating the disease or condition in the patient; and
- b. Causing gut bacterial microbiota dysbiosis in the patient, whereby said dysbiosis modulates the immune cell therapy in the patient and said disease or condition is treated or prevented.
2. An ex vivo population of immune cells for use in a method of adoptive cell therapy of a patient for treating or preventing a disease or condition in the patient, the method comprising
 - a. Carrying out adoptive immune cell therapy in the patient, comprising administering cells of said population to the patient, wherein administration of said immune cells is capable of treating, the disease or condition in the patient; and
 - b. Altering the relative proportion of a sub-population of cells of a first bacterial species or strain, or archaeal species or strain, in the gut microbiota of the patient, thereby producing an altered gut microbiota that modulates the immune cell therapy in the patient.
 3. The immune cell population of Aspect 2, wherein step (b) is carried out by targeting the sub-population of first cells by administering thereto an anti-bacterial or anti-archaeal agent (eg, a guided nuclease) simultaneously or sequentially with said immune cell population, whereby first cells are killed or the sub-population growth is reduced, thereby reducing the proportion of said sub-population in the gut microbiota of the patient.
 4. The immune cell population of any preceding Aspect, wherein the cell therapy is an adoptive T-cell therapy.
 5. The immune cell population of Aspect 4, wherein cells selected from the group consisting of CD4⁺ T-cells, CD8⁺ T-cells, T_H1 cells and T_H17 cells are administered to the patient in step (a).
 6. The immune cell population of Aspect 4 or 5, wherein T_H17 cells are modulated in the patient.
 7. The immune cell population of Aspect 4, 5 or 6, wherein T_{reg} cells are modulated in the patient.
 8. The immune cell population of any preceding Aspect, wherein the cell therapy is enhanced.
 9. The immune cell population of any preceding Aspect wherein T_H17 cells are upregulated in the patient and/or T_{reg} cells are downregulated in the patient.
 10. The immune cell population of any preceding Aspect, wherein CD4⁺ T-cell are upregulated in the patient.
 11. The immune cell population of any preceding Aspect, wherein CD8⁺ T-cells are upregulated in the patient.
 12. The immune cell population of any preceding Aspect, wherein one or more of central memory T cells (T_{CM}), effector memory T cells (T_{EM}), stem cell memory cells (T_{SCM}) and effector cells (T_{eff}) are upregulated in the patient, wherein the cells are comprised by the immune cell population administered in step (a) and/or are progeny thereof.
 13. The immune cell population of any one of Aspects 1 to 7, wherein the cell therapy is reduced.
 14. The immune cell population of Aspect 13, wherein the method reduces or prevents the risk of cytokine release syndrome (CRS) in the patient.
 15. The immune cell population of any one of Aspects 1 to 7, 13 and 14, wherein, T_H17 cells are downregulated in the patient and/or T_{reg} cells are upregulated in the patient.
 16. The immune cell population of any one of Aspects 1 to 7 and 13 to 15, wherein CD4⁺ T-cells are downregulated in the patient.

17. The immune cell population of any one of Aspects 1 to 7 and 13 to 16, wherein CD8⁺ T-cells are downregulated in the patient.
18. The immune cell population of any one of Aspects 1 to 7 and 13 to 17, wherein one or more of central memory T cells (T_{CM}), effector memory T cells (T_{EM}), stem cell memory cells (T_{SCM}) and effector cells (T_{eff}) are downregulated in the patient, wherein the cells are comprised by the immune cell population administered in step (a) and/or are progeny thereof.
19. The immune cell population of any preceding Aspect, wherein the immune cell population comprises or consists of CAR-T cells and/or T-cells expressing engineered T-cell receptors (TCRs) and/or tumour infiltrating lymphocytes (TILs).
20. The immune cell population of any preceding Aspect, wherein the immune cell population comprises or consists of engineered autologous or allogeneic immune cells, eg, T-cells, NK cells and/or TILs.
21. The immune cell population of Aspect 19 or 20, wherein the T-cells are CD4⁺ T-cells or T_H17 cells.
22. The immune cell population of any preceding Aspect, wherein step (b) increases the proportion of *Bacteroides* (eg, *B fragalis* and/or *B thetaiotamicron*) in the gut microbiota.
23. The immune cell population of any one of Aspects 1 to 21, wherein step (b) decreases the proportion of *Bacteroides* (eg, *B fragalis* and/or *B thetaiotamicron*) in the gut microbiota.
24. The immune cell population of any preceding Aspect, wherein step (b) increases the proportion of Bacteroidetes to *Firmicutes* in the gut microbiota.
25. The immune cell population of any one of Aspects 1 to 23, wherein step (b) decreases the proportion of Bacteroidetes to *Firmicutes* in the gut microbiota.
26. The immune cell population of any preceding Aspect, wherein step (b) reduces the proportion of one or more *Clostridium* species or strain (eg, wherein each species is a cluster IV or XIVa *Clostridium* species) in the gut microbiota.
27. The immune cell population of any one of Aspects 1 to 25, wherein step (b) increases the proportion of one or more *Clostridium* species or strain (eg, wherein each species is a cluster IV or XIVa *Clostridium* species) in the gut microbiota.
28. The immune cell population of any preceding Aspect, wherein step (b) reduces the proportion of *Bifidobacterium* (eg, *B bifidum*) in the gut microbiota.
29. The immune cell population of any one of Aspects 1 to 27, wherein step (b) increases the proportion of *Bifidobacterium* (eg, *B bifidum*) in the gut microbiota.
30. The immune cell population of any preceding Aspect, wherein step (b) comprises altering the relative proportion of a or said sub-population of first cells in the gut microbiota of the patient, thereby producing an altered gut microbiota that modulates the immune cell therapy in the patient, wherein the sub-population comprises host cells of said first species or strain, the method comprising
- combining the microbiota with multiple copies of engineered nucleic acid sequences encoding host modifying (HM) crRNAs, and
 - expressing HM-crRNAs in host cells,
- wherein each engineered nucleic acid sequence is operable with a Cas nuclease in a respective host cell to form a HM-CRISPR/Cas system and the engineered sequence comprises

- spacer and repeat sequences encoding a HM-crRNA;
 - the HM-crRNA comprising a sequence that is capable of hybridizing to a host cell target sequence to guide Cas nuclease to the target sequence in the host cell; and optionally the HM-system comprises a tracrRNA sequence or a DNA sequence expressing a tracrRNA sequence;
- whereby HM-crRNAs guide Cas modification of host target sequences in host cells, whereby host cells are killed or the host cell population growth is reduced, thereby reducing the proportion of said sub-population in the microbiota.
31. The immune cell population of Aspect 30, comprising using endogenous Cas nuclease of host cells for modification of target nucleotide sequences.
32. The immune cell population of Aspect 30 or 31, comprising reducing host cell population growth by at least 5-fold compared to the growth of a control population of host cells that have not received said Cas modification.
33. The immune cell population of any one of Aspects 30 to 32, comprising inhibiting host cell population growth on a gut surface.
34. The immune cell population of any one of Aspects 30 to 33, wherein the microbiota comprises cells of a second bacterial species or strain, or archaeal species or strain, wherein the second species or strain has a 16s ribosomal RNA-encoding DNA sequence that is at least 80% identical to an 16s ribosomal RNA-encoding DNA sequence of the host cell species or strain, wherein the growth of the second cells in the microbiota is not inhibited by said HM-system.
35. The immune cell population of Aspect 34, wherein the second species or strain is a human gut commensal species or strain and/or a human gut probiotic species or strain.
36. The immune cell population of Aspect 34 or 35, wherein the second species or strain is a Bacteroidetes (eg, *Bacteroides*) and optionally the host cells are gram positive bacterial cells.
37. The immune cell population of any one of Aspects 30 to 36, wherein the first cells are *Firmicutes* cells.
38. The immune cell population of any one of Aspects 30 to 37, wherein for each host cell the system comprises components according to (i) to (iv):—
- at least one nucleic acid sequence encoding a Cas nuclease;
 - an engineered HM-CRISPR array comprising a spacer sequence and repeats encoding a HM-crRNA, the HM-crRNA comprising a sequence that hybridises to a host cell target sequence to guide said Cas to the target in the host cell to modify the target sequence;
 - an optional tracrRNA sequence or a DNA sequence expressing a tracrRNA sequence;
 - wherein said components of the system are split between the host cell and at least one nucleic acid vector that transforms the host cell, whereby the HM-crRNA guides Cas to the target to modify the host target sequence in the host cell; and
- wherein the target sequence is modified by the Cas whereby the host cell is killed or host cell growth is reduced;
- the method comprising introducing the vectors of (iv) into host cells and expressing said HM-crRNA in the host cells, allowing HM-crRNA to hybridise to host cell target sequences to guide Cas to the targets in the host cells to modify target sequences, whereby host cells are killed or host cell growth is reduced, thereby altering the proportion of said sub-population in the microbiota.

39. The immune cell population of Aspect 38, wherein component (i) is endogenous to each host cell.
40. The immune cell population of Aspect 38 or 39, wherein each vector is a virus or phage.
41. The immune cell population of any one of Aspects 30 to 40, wherein each target sequence is adjacent a NNA-GAAW or NGGNG protospacer adjacent motif (PAM).
42. The immune cell population of any one of Aspects 30 to 41, wherein alternatively HM-crRNA and tracrRNA are comprised by a single guide RNA (gRNA), the method comprising introducing said gRNA into host cells or expressing the gRNA in host cells.
43. The immune cell population of any one of Aspects 30 to 35 and 37 to 42, wherein the microbiota comprises a second bacterial or archaeal species, wherein each of the first and second species is a respective species of the same phylum (eg, both *Firmicutes* species) and the growth of the second bacteria is not inhibited by the HM-system; or wherein the microbiota comprises a second bacterial or archaeal strain, wherein each of the first and second bacteria or archaea is a respective strain of the same species and the growth of the second bacteria or archaea is not inhibited by the HM-system.
44. The immune cell population of any one of Aspects 30 to 43, wherein the microbiota comprises a second bacterial species, wherein each of the first and second species is a respective gram-positive species and the growth of the second bacteria is not inhibited by the HM-system.
45. The immune cell population of any one of Aspects 30 to 44, wherein each target sequence is comprised by an antibiotic resistance gene, virulence gene or essential gene of the host cell.
46. The method of any preceding Aspect, wherein each first cell is a *Staphylococcus*, *Streptococcus*, *Pseudomonas*, *Salmonella*, *Listeria*, *E coli*, *Desulfovibrio*, *Vibrio* or *Clostridium* cell.
47. The immune cell population of any preceding Aspect, wherein step (b) comprises stimulating Paneth cells of the patient by gut *Bacteroides* (eg, *B thetaiotamicron*), wherein the altered microbiota produced by step (b) comprises an increased proportion of *Bacteroides* first bacteria compared with the microbiota before step (b), whereby Paneth cells are stimulated and the cell therapy is modulated.
48. The immune cell population of any preceding Aspect, wherein step (b) comprises developing an immune response in the patient to gut *Bacteroides* (eg, *B thetaiotamicron*), wherein the altered microbiota produced by step (b) comprises an increased proportion of *Bacteroides* first bacteria compared with the microbiota before step (b), whereby the cell therapy is modulated.
49. The immune cell population of any preceding Aspect, wherein step (b) comprises altering the relative proportion of a or said sub-population of first cells in the gut microbiota of the patient, thereby producing an altered gut microbiota that modulates the immune cell therapy in the patient, wherein step (b) comprises killing first cells of said sub-population or inhibiting growth of said sub-population by using guided nuclease targeting to the genome of first cells comprised by the sub-population.
50. A bacterial or archaeal transplant for administration to a patient for therapy of a disease or condition in the patient using the method recited in any preceding Aspect, optionally wherein the transplant comprises cells of said first species.
51. A HM-CRISPR/Cas system, HM-array or HM-crRNA as recited in any one of Aspects 30 to 45 for administration

to a patient for therapy of a disease or condition in the patient using the method recited in any one of Aspects 1 to 49.

52. A kit comprising an ex vivo population of immune cells according to any one of Aspects 1 to 49 for adoptive cell therapy of a patient to treat said disease or condition, wherein the kit further comprises a transplant, system, array or crRNA of Aspect 50 or 51, optionally wherein the immune cells are selected from CAR-T cells, T-cells expressing engineered T-cell receptors (TCRs), tumour infiltrating lymphocytes (TILs) and NK cells.

Optionally, the host cell(s), first cell(s), second cell(s) or mixed bacterial population is comprised by a human or a non-human animal subject, eg, the population is comprised by a gut microbiota, skin microbiota, oral cavity microbiota, throat microbiota, hair microbiota, armpit microbiota, vaginal microbiota, rectal microbiota, anal microbiota, ocular microbiota, nasal microbiota, tongue microbiota, lung microbiota, liver microbiota, kidney microbiota, genital microbiota, penile microbiota, scrotal microbiota, mammary gland microbiota, ear microbiota, urethra microbiota, labial microbiota, organ microbiota or dental microbiota. Optionally, the mixed bacterial population is comprised by a plant (eg, a tobacco, crop plant, fruit plant, vegetable plant or tobacco, eg on the surface of a plant or contained in a plant) or by an environment (eg, soil or water or a waterway or aqueous liquid).

Optionally, the disease or condition of a human or animal subject or patient is selected from

- (a) A neurodegenerative disease or condition;
- (b) A brain disease or condition;
- (c) A CNS disease or condition;
- (d) Memory loss or impairment;
- (e) A heart or cardiovascular disease or condition, eg, heart attack, stroke or atrial fibrillation;
- (f) A liver disease or condition;
- (g) A kidney disease or condition, eg, chronic kidney disease (CKD);
- (h) A pancreas disease or condition;
- (i) A lung disease or condition, eg, cystic fibrosis or COPD;
- (j) A gastrointestinal disease or condition;
- (k) A throat or oral cavity disease or condition;
- (l) An ocular disease or condition;
- (m) A genital disease or condition, eg, a vaginal, labial, penile or scrotal disease or condition;
- (n) A sexually-transmissible disease or condition, eg, gonorrhea, HIV infection, syphilis or *Chlamydia* infection;
- (o) An ear disease or condition;
- (p) A skin disease or condition;
- (q) A heart disease or condition;
- (r) A nasal disease or condition;
- (s) A haematological disease or condition, eg, anaemia, eg, anaemia of chronic disease or cancer;
- (t) A viral infection;
- (u) A pathogenic bacterial infection;
- (v) A cancer;
- (w) An autoimmune disease or condition, eg, SLE;
- (x) An inflammatory disease or condition, eg, rheumatoid arthritis, psoriasis, eczema, asthma, ulcerative colitis, colitis, Crohn's disease or IBD;
- (y) Autism;
- (z) ADHD;
- (aa) Bipolar disorder;
- (bb) ALS [Amyotrophic Lateral Sclerosis];
- (cc) Osteoarthritis;

(dd) A congenital or development defect or condition;
 (ee) Miscarriage;
 (ff) A blood clotting condition;
 (gg) Bronchitis;
 (hh) Dry or wet AMD;
 (ii) Neovascularisation (eg, of a tumour or in the eye);
 (jj) Common cold;
 (kk) Epilepsy;
 (ll) Fibrosis, eg, liver or lung fibrosis;
 (mm) A fungal disease or condition, eg, thrush;
 (nn) A metabolic disease or condition, eg, obesity, anorexia, diabetes, Type I or Type II diabetes;
 (oo) Ulcer(s), eg, gastric ulceration or skin ulceration;
 (pp) Dry skin;
 (qq) Sjogren's syndrome;
 (rr) Cytokine storm;
 (ss) Deafness, hearing loss or impairment;
 (tt) Slow or fast metabolism (ie, slower or faster than average for the weight, sex and age of the subject);
 (uu) Conception disorder, eg, infertility or low fertility;
 (vv) Jaundice;
 (ww) Skin rash;
 (xx) Kawasaki Disease;
 (yy) Lyme Disease;
 (zz) An allergy, eg, a nut, grass, pollen, dust mite, cat or dog fur or dander allergy;
 (aaa) Malaria, typhoid fever, *tuberculosis* or *cholera*;
 (bbb) Depression;
 (ccc) Mental retardation;
 (ddd) Microcephaly;
 (eee) Malnutrition;
 (fff) Conjunctivitis;
 (ggg) Pneumonia;
 (hhh) Pulmonary embolism;
 (iii) Pulmonary hypertension;
 (jjj) A bone disorder;
 (kkk) Sepsis or septic shock;
 (lll) Sinusitis;
 (mmm) Stress (eg, occupational stress);
 (nnn) Thalassemia, anaemia, von Willebrand Disease, or haemophilia;
 (ooo) Shingles or cold sore;
 (ppp) Menstruation;
 (qqq) Low sperm count.

Neurodegenerative or CNS Diseases or Conditions for Treatment or Prevention by the Method

In an example, the neurodegenerative or CNS disease or condition is selected from the group consisting of Alzheimer disease, geriatric psychosis, Down syndrome, Parkinson's disease, Creutzfeldt-Jakob disease, diabetic neuropathy, Parkinson syndrome, Huntington's disease, Machado-Joseph disease, amyotrophic lateral sclerosis, diabetic neuropathy, and Creutzfeldt Creutzfeldt-Jakob disease. For example, the disease is Alzheimer disease. For example, the disease is Parkinson syndrome.

In an example, wherein the method of the invention is practised on a human or animal subject for treating a CNS or neurodegenerative disease or condition, the method causes downregulation of Treg cells in the subject, thereby promoting entry of systemic monocyte-derived macrophages and/or Treg cells across the choroid plexus into the brain of the subject, whereby the disease or condition (eg, Alzheimer's disease) is treated, prevented or progression thereof is reduced. In an embodiment the method causes an increase of IFN-gamma in the CNS system (eg, in the brain and/or CSF) of the subject. In an example, the method restores nerve fibre and/or reduces the progression of nerve

fibre damage. In an example, the method restores nerve myelin and/or reduces the progression of nerve myelin damage. In an example, the method of the invention treats or prevents a disease or condition disclosed in WO2015136541 and/or the method can be used with any method disclosed in WO2015136541 (the disclosure of this document is incorporated by reference herein in its entirety, eg, for providing disclosure of such methods, diseases, conditions and potential therapeutic agents that can be administered to the subject for effecting treatment and/or prevention of CNS and neurodegenerative diseases and conditions, eg, agents such as immune checkpoint inhibitors, eg, anti-PD-1, anti-PD-L1, anti-TIM3 or other antibodies disclosed therein).

Cancers for Treatment or Prevention by the Method

Cancers that may be treated include tumours that are not vascularized, or not substantially vascularized, as well as vascularized tumours. The cancers may comprise non-solid tumours (such as haematological tumours, for example, leukaemias and lymphomas) or may comprise solid tumours. Types of cancers to be treated with the invention include, but are not limited to, carcinoma, blastoma, and sarcoma, and certain leukaemia or lymphoid malignancies, benign and malignant tumours, and malignancies e.g., sarcomas, carcinomas, and melanomas. Adult tumours/cancers and paediatric tumours/cancers are also included.

Haematologic cancers are cancers of the blood or bone marrow. Examples of haematological (or haematogenous) cancers include leukaemias, including acute leukaemias (such as acute lymphocytic leukaemia, acute myelocytic leukaemia, acute myelogenous leukaemia and myeloblasts, promyelocytic, myelomonocytic, monocytic and erythroleukaemia), chronic leukaemias (such as chronic myelocytic (granulocytic) leukaemia, chronic myelogenous leukaemia, and chronic lymphocytic leukaemia), polycythemia vera, lymphoma, Hodgkin's disease, non-Hodgkin's lymphoma (indolent and high grade forms), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, myelodysplastic syndrome, hairy cell leukaemia and myelodysplasia.

Solid tumours are abnormal masses of tissue that usually do not contain cysts or liquid areas. Solid tumours can be benign or malignant. Different types of solid tumours are named for the type of cells that form them (such as sarcomas, carcinomas, and lymphomas). Examples of solid tumours, such as sarcomas and carcinomas, include fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteosarcoma, and other sarcomas, synovialoma, mesothelioma, Ewing's tumour, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, lymphoid malignancy, pancreatic cancer, breast cancer, lung cancers, ovarian cancer, prostate cancer, hepatocellular carcinoma, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, medullary thyroid carcinoma, papillary thyroid carcinoma, pheochromocytomas sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, Wilms' tumour, cervical cancer, testicular tumour, seminoma, bladder carcinoma, melanoma, and CNS tumours (such as a glioma (such as brainstem glioma and mixed gliomas), glioblastoma (also known as glioblastoma multiforme) astrocytoma, CNS lymphoma, germinoma, medulloblastoma, Schwannoma craniopharyngioma, ependymoma, pineaioma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, neuroblastoma, retinoblastoma and brain metastases).

Autoimmune Diseases for Treatment or Prevention by the Method

Acute Disseminated Encephalomyelitis (ADEM)

Acute necrotizing hemorrhagic leukoencephalitis

Addison's disease

Agammaglobulinemia

Alopecia areata

Amyloidosis

Ankylosing spondylitis

Anti-GBM/Anti-TBM nephritis

Antiphospholipid syndrome (APS)

Autoimmune angioedema

Autoimmune aplastic anemia

Autoimmune dysautonomia

Autoimmune hepatitis

Autoimmune hyperlipidemia

Autoimmune immunodeficiency

Autoimmune inner ear disease (AIED)

Autoimmune myocarditis

Autoimmune oophoritis

Autoimmune pancreatitis

Autoimmune retinopathy

Autoimmune thrombocytopenic purpura (ATP)

Autoimmune thyroid disease

Autoimmune urticaria

Axonal & neuronal neuropathies

Balo disease

Behcet's disease

Bullous pemphigoid

Cardiomyopathy

Castleman disease

Celiac disease

Chagas disease

Chronic fatigue syndrome

Chronic inflammatory demyelinating polyneuropathy (CIDP)

Chronic recurrent multifocal osteomyelitis (CRMO)

Churg-Strauss syndrome

Cicatricial pemphigoid/benign mucosal pemphigoid

Crohn's disease

Cogans syndrome

Cold agglutinin disease

Congenital heart block

Coxsackie myocarditis

CREST disease

Essential mixed cryoglobulinemia

Demyelinating neuropathies

Dermatitis herpetiformis

Dermatomyositis

Devic's disease (neuromyelitis optica)

Discoid lupus

Dressler's syndrome

Endometriosis

Eosinophilic esophagitis

Eosinophilic fasciitis

Erythema nodosum

Experimental allergic encephalomyelitis

Evans syndrome

Fibromyalgia

Fibrosing alveolitis

Giant cell arteritis (temporal arteritis)

Giant cell myocarditis

Glomerulonephritis

Goodpasture's syndrome

Granulomatosis with Polyangiitis (GPA) (formerly called Wegener's Granulomatosis)

Graves' disease

Guillain-Barre syndrome

Hashimoto's encephalitis

Hashimoto's thyroiditis

Hemolytic anemia

Henoch-Schonlein purpura

Herpes gestationis

Hypogammaglobulinemia

Idiopathic thrombocytopenic purpura (ITP)

IgA nephropathy

IgG4-related sclerosing disease

Immunoregulatory lipoproteins

Inclusion body myositis

Interstitial cystitis

Juvenile arthritis

Juvenile diabetes (Type 1 diabetes)

Juvenile myositis

Kawasaki syndrome

Lambert-Eaton syndrome

Leukocytoclastic vasculitis

Lichen planus

Lichen sclerosus

Ligneous conjunctivitis

Linear IgA disease (LAD)

Lupus (SLE)

Lyme disease, chronic

Meniere's disease

Microscopic polyangiitis

Mixed connective tissue disease (MCTD)

Mooren's ulcer

Mucha-Habermann disease

Multiple sclerosis

Myasthenia gravis

Myositis

Narcolepsy

Neuromyelitis optica (Devic's)

Neutropenia

Ocular cicatricial pemphigoid

Optic neuritis

Palindromic rheumatism

PANDAS (Pediatric Autoimmune Neuropsychiatric Disorders Associated with *Streptococcus*)

Paraneoplastic cerebellar degeneration

Paroxysmal nocturnal hemoglobinuria (PNH)

Parry Romberg syndrome

Parsonnage-Turner syndrome

Pars planitis (peripheral uveitis)

Pemphigus

Peripheral neuropathy

Perivenous encephalomyelitis

Pernicious anemia

POEMS syndrome

Polyarteritis nodosa

Type I, II, & III autoimmune polyglandular syndromes

Polymyalgia rheumatica

Polymyositis

Postmyocardial infarction syndrome

Postpericardiotomy syndrome

Progesterone dermatitis

Primary biliary cirrhosis

Primary sclerosing cholangitis

Psoriasis

Psoriatic arthritis

Idiopathic pulmonary fibrosis

Pyoderma gangrenosum

Pure red cell aplasia

Raynauds phenomenon

Reactive Arthritis

Reflex sympathetic dystrophy
 Reiter's syndrome
 Relapsing polychondritis
 Restless legs syndrome
 Retroperitoneal fibrosis
 Rheumatic fever
 Rheumatoid arthritis
 Sarcoidosis
 Schmidt syndrome
 Scleritis
 Scleroderma
 Sjogren's syndrome
 Sperm & testicular autoimmunity
 Stiff person syndrome
 Subacute bacterial endocarditis (SBE)
 Susac's syndrome
 Sympathetic ophthalmia
 Takayasu's arteritis
 Temporal arteritis/Giant cell arteritis
 Thrombocytopenic purpura (TTP)
 Tolosa-Hunt syndrome
 Transverse myelitis
 Type 1 diabetes
 Ulcerative colitis
 Undifferentiated connective tissue disease (UCTD)
 Uveitis
 Vasculitis
 Vesiculobullous dermatosis
 Vitiligo
 Wegener's granulomatosis (now termed Granulomatosis
 with Polyangiitis (GPA)).
 Inflammatory Diseases for Treatment or Prevention by the
 Method
 Alzheimer's
 ankylosing spondylitis
 arthritis (osteoarthritis, rheumatoid arthritis (RA), psori-
 atic arthritis)
 asthma
 atherosclerosis
 Crohn's disease
 colitis
 dermatitis
 diverticulitis
 fibromyalgia
 hepatitis
 irritable bowel syndrome (IBS)
 systemic lupus erythematosus (SLE)
 nephritis
 Parkinson's disease
 ulcerative colitis.

In an example (eg, in the method of the invention involv-
 ing a mixed bacterial population), the host cell (or first cell
 or second cell) genus or species is selected from a genus or
 species listed in Table 1. In examples of the present inven-
 tion, the Cas (eg, Cas nuclease such as a Type I, II or III Cas,
 eg, a Cas3 or 9) is a Cas comprised by bacteria of a genus
 or species that is selected from a genus or species listed in
 Table 1, and optionally the host cell (or first cell or second
 cell) is of the same genus or species. In an example of this,
 the Cas is endogenous to said host cell (or first or second
 cell), which is useful for embodiments herein wherein
 endogenous Cas is used to modify a target sequence. In this
 case, the HM-array may comprise one or more repeat
 nucleotide (eg, DNA or RNA) sequences that is at least 90,
 95, 96, 97, 98 or 99% identical (or is 100% identical) to a
 repeat sequence of said cell, genus or species, whereby the
 Cas is operable with cRNA encoded by the HM-array for

modifying one or more target sequences in the cell. In an
 example, the Cas is a Type I Cas3 and is used with a Type
 I CASCADE, wherein one or both of the Cas3 and CAS-
 CADE are endogenous to the host or first cells, or are
 5 vector-borne (ie, exogenous to the host or first cells).

In an example, the method of the invention selectively
 kills first cells in the microbiota whilst not targeting second
 cells, eg, wherein the second cells are (a) of a related strain
 to the strain of the first species or (b) of a species that is
 10 different to the first species and is phylogenetically related to
 the first species, wherein the second species or strain has a
 16s ribosomal RNA-encoding DNA sequence that is at least
 80% identical to an 16s ribosomal RNA-encoding DNA
 sequence of the first cell species or strain. In an embodiment,
 15 the first cells are of a first species selected from Table 1 and
 the second cells are of a different species selected from Table
 1. In an example, the species are of the same genus or are of
 different genera.

It will be understood that particular embodiments
 20 described herein are shown by way of illustration and not as
 limitations of the invention. The principal features of this
 invention can be employed in various embodiments without
 departing from the scope of the invention. Those skilled in
 the art will recognize, or be able to ascertain using no more
 than routine study, numerous equivalents to the specific
 procedures described herein. Such equivalents are consid-
 25 ered to be within the scope of this invention and are covered
 by the claims. All publications and patent applications
 mentioned in the specification are indicative of the level of
 skill of those skilled in the art to which this invention
 pertains. All publications and patent applications and all US
 equivalent patent applications and patents are herein incor-
 porated by reference to the same extent as if each individual
 publication or patent application was specifically and indi-
 35 vidualy indicated to be incorporated by reference. The use
 of the word "a" or "an" when used in conjunction with the
 term "comprising" in the claims and/or the specification may
 mean "one," but it is also consistent with the meaning of
 "one or more," "at least one," and "one or more than one."
 40 The use of the term "or" in the claims is used to mean
 "and/or" unless explicitly indicated to refer to alternatives
 only or the alternatives are mutually exclusive, although the
 disclosure supports a definition that refers to only alterna-
 tives and "and/or." Throughout this application, the term
 45 "about" is used to indicate that a value includes the inherent
 variation of error for the device, the method being employed
 to determine the value, or the variation that exists among the
 study subjects.

As used in this specification and claim(s), the words
 50 "comprising" (and any form of comprising, such as "com-
 prise" and "comprises"), "having" (and any form of having,
 such as "have" and "has"), "including" (and any form of
 including, such as "includes" and "include") or "containing"
 (and any form of containing, such as "contains" and "con-
 55 tain") are inclusive or open-ended and do not exclude
 additional, unrecited elements or method steps

The term "or combinations thereof" or similar as used
 herein refers to all permutations and combinations of the
 listed items preceding the term. For example, "A, B, C, or
 combinations thereof" is intended to include at least one of:
 A, B, C, AB, AC, BC, or ABC, and if order is important in
 a particular context, also BA, CA, CB, CBA, BCA, ACB,
 BAC, or CAB. Continuing with this example, expressly
 included are combinations that contain repeats of one or
 60 more item or term, such as BB, AAA, MB, BBC,
 AAABCCCC, CBBAAA, CABABB, and so forth. The
 skilled artisan will understand that typically there is no limit

on the number of items or terms in any combination, unless otherwise apparent from the context.

Any part of this disclosure may be read in combination with any other part of the disclosure, unless otherwise apparent from the context.

All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

The present invention is described in more detail in the following non limiting Examples.

EXAMPLES

Example 1: Specific Microbiota Bacterial Population Growth Inhibition by Harnessing Wild-Type Endogenous Cas

1. Material and Methods

1.1. Strains

The following strains were used in the course of this Example and Examples 2 and 3: *E. coli* MG1655, *E. coli* TOP10, *Streptococcus thermophilus* LMD-9 (ATCC BAA-491, Manassas, Va.), *Streptococcus thermophilus* DSM 20617(T) (DSMZ, Braunschweig, Germany), *Lactococcus lactis* MG1363 and *Streptococcus mutans* Clarke 1924 DSM 20523 (DSMZ, Braunschweig, Germany).

During the course of media selection and testing of the genetic constructs different Streptococci strains were used. *Streptococcus thermophilus* LMD-9 (ATCC BAA-491) and *Escherichia coli* TOP10 were considered because of their compatible growth requirements. All strains were cultivated in Todd-Hewitt broth (TH) (T1438 Sigma-Aldrich), in aerobic conditions and at 37° C., unless elsewhere indicated. The strains were stored in 25% glycerol at -80° C.

1.2. Differential Growth Media

All strains were grown on TH media at 37° C. for 20 hours. Selective media for *S. thermophilus* was TH media supplemented with 3 g l⁻¹ of 2-phenylethanol (PEA). PEA was added to the media and autoclaved at 121° C. for 15 minutes at 15 psi. Agar plates were prepared by adding 1.5% (wt/vol) agar to the corresponding media. When necessary for selection or plasmid maintenance 30 µg ml⁻¹ kanamycin was used for both *S. thermophilus* strains and *E. coli*, and 500 µg ml⁻¹ for *S. mutans*.

In some cases, depending on the strain and plasmid, a longer incubation, up to 48 hours, may be needed to see growth on media supplemented with PEA. In order to control for the viability of the organisms used, a control TH agar must be done in parallel.

1.3. Cloning

E. coli (One Shot® ThermoFischer TOP10 Chemically Competent cells) was used in all subcloning procedures. PCR was carried out using Phusion polymerase. All PCR products were purified with Nucleospin Gel and PCR Clean-up by Macherey-Nagel following the manufacturer's protocol. The purified fragments were digested with restriction enzyme DpnI in 1λ FD buffer with 1 µl enzyme in a total

volume of 34 µl. The digested reaction was again purified with Nucleospin Gel and PCR Clean-up by Macherey-Nagel following the manufacturer's protocol. Gibson assembly was performed in 10 µl reactions following the manufacturer's protocol (NewEngland Biolab).

Plasmid DNA was prepared using Qiagen kits according to the manufacturer's instructions. Modifications for Gram-positive strains included growing bacteria in a medium supplemented with 0.5% glycine and lysozyme to facilitate cell lysis.

1.4. Transformation

1.4.1 Electro-Competent *E. coli* Cells and Transformation

Commercially electrocompetent cells were used for cloning and the experiments (One Shot® ThermoFischer TOP10 Chemically Competent *E. coli*). Electroporation was done using standard settings: 1800 V, 25 µF and 200Ω using an Electro Cell Manipulator (BTX Harvard Apparatus ECM630). Following the pulse, 1 ml LB-SOC media was added and the cells were incubated at 37° C. for 1 hour. The transformed cells were plated in LB-agar containing 50 µg ml⁻¹ of kanamycin.

1.4.2 Preparation of Electro-Competent *S. thermophilus* Cells

The electroporation protocol was modified from Somkuti and Steinberg, 1988. An overnight culture of *Streptococcus thermophilus* in TH Broth supplemented with 40 mM DL-threonine (T8375 Sigma-Aldrich) was diluted 100-fold in 5 ml of the same media and grown to an OD₆₀₀ between 0.3-0.5 (approximately 2.5 hours after inoculation). The cells were collected by centrifugation at 10,000×g for 10 min at 4° C. and washed three times with 5 ml of ice cold wash buffer (0.5 M sucrose+10% glycerol). After the cells were washed, they were suspended to an OD₆₀₀ of 15-30 in electroporation buffer (0.5 M sucrose, 10% glycerol and 1 mM MgCl₂). The cells in the electroporation buffer may be kept at 4° C. until use (within one hour) or aliquot 50 µl in eppendorf tubes, freezing them in liquid nitrogen and stored at -80° C. for later use.

1.4.3 Electroporation *S. thermophilus* Cells

1 µl of purified plasmid DNA was added to 50 µl of the cell suspension and electroporation was carried out in 2 mm-gap electroporation cuvettes pre-cooled. The electroporation setting were 2500 V, 25 µF and 200Ω using an Electro Cell Manipulator (BTX Harvard Apparatus ECM630) Immediately after the electric pulse, 1 ml of TH broth was added to the cells and the suspension was kept on ice for 10 minutes, subsequently the cells were incubated for 3 h at 37° C. After allowing time for expression of the resistance gene the cells were plated onto TH-agar plates containing 30 µg ml⁻¹ of kanamycin. Depending on the construct, colonies were visible between 12 and 48 h of incubation at 37° C.

1.5. Construction of XylS Plasmid

All the plasmids used in this work were based on pBAV1K-T5, which is a broad-host range expression vector derived from the a cryptic plasmid pWV01 from *Streptococcus cremoris* (Bryksin & Matsumura, 2010), the backbone was amplified using that contain overhangs for assembly with the other fragments using Gibson's method.

The xylose inducible system was constructed by cloning the promoter *gyrA* in front of the XylR repressor (FIG. 1). The XylR repressor was amplified from *Bacillus Subtilis* strain SCK6 (Zhang et al. 2011) with the a reverse primer that includes an overhang for Gibson assembly and a forward primer, that is an ultramer used to introduce the *gyrA* promoter (Xie et al. 2013) and the corresponding overhang for assembly into pBAV1KT5 backbone. The resulting fragment was flanked by an mCherry amplified from pCL002

(unpublished work) with an ultramer that include PldhA+PxyIA hybrid promoter (Xie et al. 2013). The three resulting PCR products were assembled in a Gibson Master Mix® (NewEngland Biolab) according to manufacturer's instructions. The product was finally transformed in *E. coli* TOP10 electrocompetent cells. See FIG. 1.

1.6. Design and Construction of CRISPR Array Plasmid

Streptococcus thermophilus has 4 distinct CRISPR systems (Sapranauskas, et al. 2011), for this work the type II CRISPR1 (ST1-CRISPR) system was chosen. The design of the target sequence was based on the available genome sequence of LMD-9 (GenBank: CP000419.1). The ST1-CRISPR array was designed to contain only the CRISPR array repeats and spacers under a xylose inducible promoter (Xie et al. 2013), followed by the corresponding tracrRNA under a strong constitutive promoter for Streptococci species (Sorg et al. 2014) (FIG. 2).

The tracrRNA plays a role in the maturation of crRNA and it is processed by *S. thermophilus* endogenous RNase III, forming a complex with crRNA. This complex acts as a guide for the endonuclease ST1-Cas9 (Horvath & Barrangou, 2010). After transcription of the synthetic array from the xylose inducible promoter, the endogenous Cas9 and RNases will process it into a functional gRNA. The gRNA/Cas9 complex will cause a double stranded break at the target location.

The design of the array used 2 specific target sequences high on GC content and a reduced portion of the tracrRNA (ie, a less than complete tracrRNA sequence), which has been suggested not to be necessary for proper maturation of crRNA (Horvath & Barrangou, 2010).

The 2 targets were an essential gene (DNA polymerase III subunit alpha) and an antibiotic resistance gene (tetA-like gene).

Primers were used to amplify pBAV1KT5-XylR-PldhA backbone. The CRISPR array gBlock and the backbone with overhangs were assembled in a Gibson Master Mix® according to manufacturer's instructions (NewEngland Biolabs). The product was finally transformed in *E. coli* TOP10 electrocompetent cells.

1.7. Characterization of Xylose Inducible System in *Streptococcus thermophilus* LMD-9

Overnight stationary-phase cultures were diluted 1:100 into TH broth with corresponding antibiotic. Mid-log cells were induced with different concentration of D-(+)-xylose (0, 0.001, 0.01, 0.1, 0.5 and 1% wt/vol) and the cell cultures were measured either directly in medium to assess the extent of autofluorescence of the media, on the cell suspension or the suspension buffer (PBS buffer). 20 µl samples of the cell cultures were diluted 1/10 on PBS buffer, on 96-well plates with flat bottoms. Fluorescence of cell suspensions or media was read on a plate reader. mCherry fluorescence was measured using an excitation wavelength of 558 nm and emission at 612 nm. Absorbance of the resuspended cells was measured at OD 600 nm. A minimum of three independent biological replicates was done for each experiment.

1.8. Activation of CRISPR Array in *S. thermophilus*

S. thermophilus LMD-9 and *E. coli* TOP10 both with the plasmid containing the CRISPR array targeting the DNA polymerase III and tetA of *S. thermophilus* were grown overnight in 3 ml cultures supplemented with 30 µg ml⁻¹ of kanamycin for plasmid maintenance. The next day 96 well deep well plates were inoculated with 500 µl of 1/100 of overnight culture in fresh TH media, supplemented with 30 ml kanamycin. Mid-log cell cultures were induced with 1% xylose. The killing effect was tested on *S. thermophilus* and *E. coli* alone. For each strain and condition tested a negative

control was kept without xylose. The cells were grown till ~OD 0.5 and next 10-fold serially diluted in TH media and using a 96-well replicator (Mettler Toledo Liquidator™ 96) 5 µL volume drops were spotted on TH agar and TH agar supplemented with g l⁻¹ PEA plates. The plates were incubated for 24H at 37° C. and the colony forming units (CFU) were calculated from triplicate measurements.

2. Results

2.1 Growth Condition and Selective Media

We first set out to establish the bacterial strains and cultivation protocol that would support growth for all strains we planned to use for the co-cultivation experiments. We used *S. thermophilus* strain LMD-9 which was able to support a similar growth as *E. coli* in TH broth at 37° C. (FIG. 3).

Distinguishing the different bacteria from a mixed culture is important in order to determine cell number of the different species. With MacConkey agar is possible to selectively grow *E. coli*, however there is no specific media for selective growth of *S. thermophilus*. PEA agar is a selective medium that is used for the isolation of gram-positive (*S. thermophilus*) from gram-negative (*E. coli*). Additionally, we found that different concentrations of PEA partially inhibit the growth of other gram positives, which allow for selection between the other gram-positive bacteria used in this work (FIG. 4). 3 g l⁻¹ of PEA proved to selectively grow *S. thermophilus* LMD-9 while limiting growth of *E. coli*.

2.2 Design and Validation of Inducible System

An induction system for *Streptococcus* species was previously developed based on the *Bacillus megaterium* xylose operon (FIG. 5) by creating a heterologous xylose induction cassette (Xyl-S). The xylR and xylA promoters were replaced with *S. mutans*' constitutively expressed gyrA and ldh promoters respectively. This expression cassette for *Streptococcus* species showed differences in sensitivity and expression levels between different species, however the system was not tested in *S. thermophilus* (Xie et al. 2013). Therefore we first set out to validate the xylose induction cassette in *S. thermophilus*.

An alternative version of the induction cassette was constructed by only replacing the xylR promoter with the *S. mutans*' gyrA promoter but left the endogenous *B. megaterium* xylA promoter intact. During the design of the xylose inducible system we considered both versions of the inducible promoter, the natural P_{XylA} promoter found in *Bacillus megaterium* and a hybrid promoter of the highly conserved promoter P_{ldhA} fused with the repressor binding sites of P_{XylA} promoter (FIG. 5). Only a few *Streptococcus* species have been reported to metabolize xylose, and thus the presence of a regulatory machinery to recognize the xylA promoter in the other *Streptococcus* species is not likely. Therefore we constructed both xylose induction systems but only tested the inducibility of mCherry with the P_{ldhA+XylA} system.

In order to determine mCherry inducible expression by xylose, mid-log cultures of cells with the plasmid (pBAV1KT5-XylR-mCherry-P_{ldhA+XylA}) were induced with different concentrations of xylose. Six hours after the induction we measured mCherry fluorescence in the cultures, where we observed substantially higher overall expression levels in cells carrying the plasmid (FIG. 6). It is worth noticing that the system showed a substantial level of basal expression even in the cultures where xylose was not added. This means that the system is 'leaky' and in context of the kill-array this can lead to cell death even before the system is induced with xylose. However, in the subsequent course

of this study we used both versions of the plasmid (pBAV1KT5-XylR-mCherry-P_{ldha+XylA} and pBAV1KT5-XylR-mCherry-P_{xyIA}).

2.3 Design of CRISPR/CAS9 Array

In order to determine if the genomic targeting spacers in a CRISPR array can cause death in *S. thermophilus* LMD-9, we inserted the CRISPR array we designed into the two xylose inducible systems previously constructed (pBAV1KT5-XylR-mCherry-P_{ldha+XylA} and pBAV1KT5-XylR-mCherry-P_{xyIA}). In these plasmids we replaced mCherry with the gBlock containing the CRISPR array (FIG. 7). The variant with the P_{ldha+XylA} promoter was expected to be stronger and have a higher basal activity than the P_{xyIA} (Xie et al. 2013).

2.4 Inhibition of Bacterial Population Growth Using Endogenous Cas9

After we constructed the plasmids in *E. coli*, we transformed the plasmids into *S. thermophilus*. This would allow us to determine if we could cause cell death of a specific bacterial species. Interestingly, bacterial host population size (indicated by growing bacteria and counting colony numbers on agar plates) in *S. thermophilus* exposed to the plasmid containing the strong P_{ldha+XylA} hybrid promoter was 10-fold less when compared to *S. thermophilus* exposed to the plasmid containing the weak, normal P_{xyIA} promoter (FIG. 8; 52 colonies with the strong array expression versus 556 colonies with weak array expression, 10.7-fold difference), the 2 strains having been transformed in parallel using the same batch of electrocompetent *S. thermophilus* cells. This suggests to us that the plasmid carrying the CRISPR array targeting *S. thermophilus* genes is able to kill the cells using the endogenous Cas nuclease and RNase III, thereby inhibiting population growth by 10-fold.

We expect that weak array expression in host cells transformed by the plasmid comprising the P_{xyIA} promoter led to a degree of cell killing, albeit much less than with the strong promoter plasmid. We expect that population growth inhibition that is greater than the observed 10-fold inhibition would be determined if a comparison of the activity of strong array expression was made with *S. thermophilus* that is not exposed to any array-encoding plasmid (such as bacteria directly isolated from gut microbiota). Thus, we believe that array (or single guide RNA) expression in host cells for harnessing endogenous Cas nuclease will be useful for providing effective growth inhibition of target host cells in environmental, medical and other settings mentioned herein. Co-administration of antibiotic may also be useful to enhance the growth inhibition, particularly when one or more antibiotic resistance genes are targeted.

3. Discussion and Outlook

In this study we set out to design a CRISPR-array to specifically kill *S. thermophilus* using the endogenous Cas9 system. In order to gain control over the killing signal we sought to apply an inducible system that can be applied in *S. thermophilus*. The xylose inducible XylR system from *B. megaterium* was previously applied in *S. mutans* (Xie, 2013) but not in *S. thermophilus*. In this study we demonstrated the functionality of the xylR induction system using the designed XylR-mCherry-P_{ldha} circuit in *S. thermophilus*. We found 0.1% wt/vol is sufficient to fully induce the XylR system in *S. thermophilus* (FIG. 6).

In order to observe abundance when co-culturing *S. thermophilus* and *E. coli* we established that supplementation of the culture media with 3 g l⁻¹ of PEA, allows for the selective growth of *S. thermophilus* while limiting the growth of *E. coli* (FIG. 4).

A ST1-CRISPR array, targeting the DNA polymerase III subunit alpha and a tetA like gene in the *S. thermophilus* LMD-9 genome, was placed under the xylose inducible promoter (Xie et al. 2013). Targeting these regions should lead to a double strand break and thus limit *S. thermophilus* viability (FIG. 9). Since the engineered array was designed to target *S. thermophilus* genome using the endogenous CRISPR/Cas machinery to process the encoded CRISPR array, the array is expected to have no influence on growth of unrelated strains such as *E. coli*, even similar targets could be found on its genome. This was successfully tested in a mixed bacterial population (simulating aspects of a human microbiota) as discussed in Example 3.

The demonstration of the ability to inhibit host cell growth on a surface is important and desirable in embodiments where the invention is for treating or preventing diseases or conditions mediated or caused by microbiota as disclosed herein in a human or animal subject. Such microbiota are typically in contact with tissue of the subject (eg, gut tissue) and thus we believe that the demonstration of activity to inhibit growth of a microbiota bacterial species (exemplified by *Streptococcus*) on a surface supports this utility.

Example 2: Specific Microbiota Bacterial Population Growth Inhibition in Different Strains

Example 1 demonstrated specific growth inhibition of *Streptococcus thermophilus* LMD-9. Here we demonstrate growth inhibition can also be obtained in a second strain: *Streptococcus thermophilus* DSM 20617. Methods described in Example 1 were, therefore, applied to the latter strain (except that selective media for *S. thermophilus* DSM 20617 was TH media supplemented with 2.5 g l⁻¹ of 2-phenylethanol (PEA)).

Streptococcus thermophilus DSM 20617 transformed with the CRISPR array plasmids were incubated for recovery in liquid media for a period of 3 hours at 37° C. that would allow for expression of kanamycin resistance. After a recovery period, cells were plated in different selection media in presence of 1% xylose in order to induce cell death, and without xylose as a control (FIG. 10). It is evident that; (1) by xylose induction the growth of *S. thermophilus* can be inhibited (around 10-fold for the 'strong' promoter plasmid versus control), (2) the 'strong' system (pBAV1KT5-XylR-CRISPR-P_{ldha}) results in more growth reduction than the 'weak' system (pBAV1KT5-XylR-CRISPR-P_{xyIA})

Example 3: Selective Bacterial Population Growth Inhibition in a Mixed Consortium of Different Microbiota Species

We next demonstrated selective growth inhibition of a specific bacterial species in a mixed population of three species. We selected species found in gut microbiota of humans and animals (*S. thermophilus* DSM 20617(T), *Lactobacillus lactis* and *E. coli*). We included two gram-positive species (the *S. thermophilus* and *L. lactis*) to see if this would affect the ability for selective killing of the former species; furthermore to increase difficulty (and to more closely simulate situations in microbiota) *L. lactis* was chosen as this is a phylogenetically-related species to *S. thermophilus* (as indicated by high 16s ribosomal RNA sequence identity between the two species). The *S. thermophilus* and *L. lactis* are both *Firmicutes*. Furthermore, to simulate microbiota, a human commensal gut species (*E. coli*) was included.

1. Materials & Methods

Methods as set out in Example 1 were used (except that selective media was TH media supplemented with 2.5 g l⁻¹ of 2-phenylethanol (PEA)).

1.1 Preparation of Electro-Competent *L. lactis* Cells

Overnight cultures of *L. lactis* in TH media supplemented with 0.5 M sucrose and 1% glycine were diluted 100-fold in 5 ml of the same media and grown at 30° C. to an OD₆₀₀ between 0.2-0.7 (approximately 2 hours after inoculation). The cells were collected at 7000×g for 5 min at 4° C. and washed three times with 5 ml of ice cold wash buffer (0.5 M sucrose+10% glycerol). After the cells were washed, they were suspended to an OD₆₀₀ of 15-30 in electroporation buffer (0.5 M sucrose, 10% glycerol and 1 mM MgCl₂). The cells in the electroporation buffer were kept at 4° C. until use (within one hour) or aliquot 50 µl in eppendorf tubes, freezing them in liquid nitrogen and stored at -80° C. for later use.

Electroporation conditions for all species were as described in Example 1.

1.2 Activation of CRISPR Array: Consortium Experiments.

S. thermophilus DSM 20617, *L. lactis* MG1363 and *E. coli* TOP10 were genetically transformed with the plasmid containing the CRISPR array targeting the DNA polymerase III and tetA of *S. thermophilus*. After transformation all cells were grown alone and in co-culture for 3 hours at 37° C. allowing for recovery to develop the antibiotic resistance encoded in the plasmid. We decided to use transformation efficiency as a read out of CRISPR-encoded growth inhibition. Therefore, after allowing the cells for recovery the cultures were plated in TH media, TH supplemented with PEA and MacConkey agar all supplemented with Kanamycin, and induced by 1% xylose.

2. Results

2.0 Phylogenetic Distance Between *L. lactis*, *E. Coli* and *S. thermophilus*

The calculated sequence similarity in the 16S rRNA-encoding DNA sequence of the *S. thermophilus* and *L. lactis* was determined as 83.3%. The following 16S sequences were used: *E. coli*: AB030918.1, *S. thermophilus*: AY188354.1, *L. lactis*: AB030918. The sequences were aligned with needle (http://www.ebi.ac.uk/Tools/psa/emboss_needle/nucleotide.html) with the following parameters: -gapopen 10.0 -gapextend 0.5 -endopen 10.0 -endextend 0.5 -format3 pair -snucleotide1 -snucleotide2. FIG. 11 shows the maximum-likelihood phylogenetic tree of 16S sequences from *S. thermophilus*, *L. lactis* and *E. coli*.

2.1 Growth Condition and Selective Media

S. thermophilus and *L. lactis* are commonly used in combination in many fermented foods and yoghurt. We chose these strains since they are commonly known to be gut microbes that form an intimate association with the host and previous characterizations of the 16S ribosomal RNA region of *S. thermophilus* and *L. lactis* have shown that these organisms are phylogenetically closely related (Ludwig et al., 1995). In parallel we also evaluated the growth of *E. coli* for our mixed population co-culture experiments, since this organism is also commonly found in gut microbe communities. We first set out to establish the bacterial strains and cultivation protocol that would support growth for all strains we planned to use for the co-cultivation experiments. We found that all strains were able to support growth in TH broth at 37° C. (FIG. 3).

Distinguishing the different bacteria from a mixed culture is important in order to determine cell number of the different species. With MacConkey agar is possible to selec-

tively grow *E. coli*, however there is no specific media for selective growth of *S. thermophilus*. PEA agar is a selective medium that is used for the isolation of gram-positive (*S. thermophilus*) from gram-negative (*E. coli*). Additionally, different concentrations of PEA partially inhibit the growth of the different gram positive species and strains, which allow for selection between the other gram-positive bacteria used in this work. Using 2.5 g l⁻¹ of PEA proved to selectively grow *S. thermophilus* while limiting growth of *L. lactis* and *E. coli*.

All strains were transformed with a plasmid that used the vector backbone of pBAV1KT5 that has a kanamycin selection marker; we found that using media supplemented with 30 µg ml⁻¹ of kanamycin was enough to grow the cells while keeping the plasmid.

2.3 Transformation & Selective Growth Inhibition in a Mixed Population

We transformed *S. thermophilus*, *L. lactis* and *E. coli* with plasmid containing the CRISPR array and cultured them in a consortium of all the bacterial species combined in equal parts, which would allow us to determine if we could cause cell death specifically in *S. thermophilus*. We transformed all the species with either the pBAV1KT5-XylR-CRISPR-P_{xylA} or pBAV1KT5-XylR-CRISPR-P_{ldhA+XylA} plasmid.

FIG. 12 shows the selective *S. thermophilus* growth inhibition in a co-culture of *E. coli*, *L. lactis* and *S. thermophilus* harboring either the pBAV1KT5-XylR-CRISPR-P_{xylA} or the pBAV1KT5-XylR-CRISPR-P_{ldhA+XylA} plasmid. No growth difference is observed between *E. coli* harboring the pBAV1KT5-XylR-CRISPR-P_{xylA} or the pBAV1KT5-XylR-CRISPR-P_{ldhA+XylA} plasmid (middle column). However, *S. thermophilus* (selectively grown on TH agar supplemented with 2.5 g l⁻¹ PEA, last column) shows a decrease in transformation efficiency between the pBAV1KT5-XylR-CRISPR-P_{xylA} (strong) or the pBAV1KT5-XylR-CRISPR-P_{ldhA+XylA} (weak) plasmid as we expected. We thus demonstrated a selective growth inhibition of the target *S. thermophilus* sub-population in the mixed population of cells.

REFERENCES

- Barrangou, R., Fremaux, C., Deveau, H., Richards, M., Patrick Boyaval, Moineau, S., . . . Horvath, P. (2007). CRISPR Provides Acquired Resistance Against Viruses in Prokaryotes. *Science*, 315 (March), 1709-1712.
- Bryksin, A. V., & Matsumura, I. (2010). Rational design of a plasmid origin that replicates efficiently in both gram-positive and gram-negative bacteria. *PLoS One*, 5(10), e13244.
- Chan C T Y, Lee J W, Cameron D E, Bashor C J, Collins J J: "Deadman" and "Passcode" microbial kill switches for bacterial containment. *Nat Chem Biol* 2015, 12(Dec.): 1-7.
- Horvath, P., Romero, D. A., Cofut -Monvoisin, A.-C., Richards, M., Deveau, H., Moineau, S., . . . Barrangou, R. (2008). Diversity, activity, and evolution of CRISPR loci in *Streptococcus thermophilus*. *Journal of Bacteriology*, 190(4), 1401-12.
- Ludwig, E. S., Klipper, R., Magrum L., Wose C., & Stackebrandt, E. (1985). The phylogenetic position of *Streptococcus* and *Enterococcus*. *Journal of General Microbiology*, 131, 543-55 1.
- Mercenier, A. (1990). Molecular genetics of *Streptococcus thermophilus*. *FEMS Microbiology Letters*, 87(1-2), 61-77. \

7. Samaržija, D., Antunac, N., & Havranek, J. (2001). Taxonomy, physiology and growth of *Lactococcus lactis*: a review. *Mljekarstvo*, 51(1), 35-48. Retrieved from
8. Sapranauskas, R., Gasiunas, G., Fremaux, C., Barrangou, R., Horvath, P., & Siksnys, V. (2011). The *Streptococcus thermophilus* CRISPR/Cas system provides immunity in *Escherichia coli*. *Nucleic Acids Research*, 39(21), 9275-9282.
9. Somkuti, G. A., & Steinberg, D. H. (1988). Genetic transformation of *Streptococcus thermophilus* by electroporation. *Biochimie*, 70(4), 579-585
10. Sorg, R. A., Kuipers, O. P., & Veening, J.-W. (2014). Gene expression platform for synthetic biology in the human pathogen *Streptococcus pneumoniae*. *ACS Synthetic Biology*, 4(3), 228-239.
11. Suvorov, a. (1988). Transformation of group A streptococci by electroporation. *FEMS Microbiology Letters*, 56(1), 95-99.
12. Xie, Z., Qi, F., & Merritt, J. (2013). Development of a tunable wide-range gene induction system useful for the study of streptococcal toxin-antitoxin systems. *Applied and Environmental Microbiology*, 79(20), 6375-84.
13. Zhang, X. Z., & Zhang, Y. H. P. (2011). Simple, fast and high-efficiency transformation system for directed evolution of cellulase in *Bacillus subtilis*. *Microbial Biotechnology*, 4(1), 98-105.

Example 4: Altering the Ratio of *Clostridium Dificile* in a Mixed Gut Microbiota Population

Alteration of the ratio of bacteria will be performed according to the present example, which is described by reference to knocking-down *Clostridium difcile* bacteria in a mixed gut microbiota sample. The sample will contain *Bacteroides* and metronidazole (MTZ)-resistant *C difcile* strain 630 sub-populations. Ex vivo the mixed population is combined with a population of carrier bacteria (*Lactobacillus acidophilus* La-14 and/or La-5) that have been engineered to contain CRISPR arrays.

Each CRISPR array is comprised on a plasmid that is compatible with the carrier bacterium and *C difcile* cells. The array is comprised by a *Bacteroides thetaiotamicron* CTnDot transposon that also comprises oriT, an intDOT sequence, a tetQ-rteA-rteB operon, rteC and the operon xis2c-xis2d-orf3-exc. In one experiment, mob and tra operons are excluded (instead relying on these supplied by *Bacteroides* cells to which the transposons are transferred in the mixture combined with the carrier bacteria). In another experiment, the mob and tra operons are included in the transposons.

Protein translocation across the cytoplasmic membrane is an essential process in all bacteria. The Sec system, comprising at its core an ATPase, SecA, and a membrane channel, SecYEG, is responsible for the majority of this protein transport. A second parallel Sec system has been described in a number of Gram-positive species. This accessory Sec system is characterized by the presence of a second copy of the energizing ATPase, SecA2; where it has been studied, SecA2 is responsible for the translocation of a subset of Sec substrates. In common with many pathogenic Gram-positive species, *Clostridium difcile* possesses two copies of SecA. Export of the S-layer proteins (SLPs) and an additional cell wall protein (CwpV) is dependent on SecA2. Accumulation of the cytoplasmic precursor of the SLPs SlpA and other cell wall proteins is observed in cells expressing dominant-negative secA1 or secA2 alleles, concomitant with a decrease in the levels of mature SLPs in the

cell wall. Furthermore, expression of either dominant-negative allele or antisense RNA knockdown of SecA1 or SecA2 dramatically impairs growth, indicating that both Sec systems are essential in *C. difcile*.

C. difcile Strain 630 (epidemic type X) has a single circular chromosome with 4,290,252 bp (G+C content=29.06%) and a circular plasmid with 7,881 bp (G+C content=27.9%). The whole genome has been sequenced and found that 11% of the genome consists of mobile genetic elements such as conjugative transposons. These elements provide *C. difcile* with the genes responsible for its antimicrobial resistance, virulence, host interaction and the production of surface structures. For example, the cdeA gene of *C. difcile* produces a multidrug efflux pump which was shown to be homologous to known efflux transporters in the multidrug and toxic compound extrusion (MATE) family. The protein facilitates energy-dependent and sodium-coupled efflux of drugs from cells. In addition, the cme gene in *C. difcile* has been shown to provide multidrug resistance in other bacteria.

The array comprises a R1-S1-R1' CRISPR unit (spacer flanked by two CRISPR repeats) for targeting a sequence in an essential gene (SecA2) of *C difcile* cells. In another experiment, targeting is to the cdeA gene in the presence of MTZ and optionally one or more other anti-*C difcile* antibiotics. Each spacer (S) comprises a 20mer nucleotide sequence of the SecA or cdeA gene, wherein the sequence comprises a PAM of a *C difcile* strain 630 CRISPR/Cas system that is cognate to the repeat sequences. Each repeat is identical to a *C difcile* strain 630 repeat.

The repeats function with Cas that is endogenous to the *C difcile* cells in the mixed population. The mixed population of bacteria is retrieved as an ex vivo sample from a stool sample of a human patient suffering from *C difcile* infection. The mixed population is mixed with the carrier bacteria in vitro and incubated at 37 degrees centigrade under anaerobic conditions to simulate gut conditions in the presence of tetracycline. It is expected that transposons containing the CRISPR arrays will be transferred to *Bacteroides* and *C difcile* cells in the mixture. Furthermore, it is expected that the target sites in the latter cells will be cut by Cas nuclease action, thus reducing the proportion of *C difcile* in the mixed population (and increasing the ratio of *Bacteroides* versus *C difcile*).

In a follow-on experiment, a drink is produced comprising the carrier bacteria and this is consumed by the human patient once or twice for several consecutive days with or without an ant-acid. The patient is also administered with tetracycline during the treatment period. It is expected that stool analysis will reveal that the proportion of *C difcile* in the stool samples will reduce (and the ratio of *Bacteroides* versus *C difcile* will increase).

Example 5: Vector-Encoded System for Selective Species & Strain Growth Inhibition in A Mixed Bacterial Consortium

In Example 3 we surprisingly established the possibility of harnessing endogenous Cas nuclease activity in host bacteria for selective population growth inhibition in a mixed consortium of different species. We next explored the possibility of instead using vector-encoded Cas activity for selective population growth inhibition in a mixed consortium of different species. We demonstrated selective growth inhibition of a specific bacterial species in a mixed population of three different species, and further including a strain alternative to the target bacteria. We could surprisingly show

selective growth inhibition of just the target strain of the predetermined target species. Furthermore, the alternative strain was not targeted by the vector-encoded CRISPR/Cas system, which was desirable for establishing the fine specificity of such vector-borne systems in a mixed bacterial consortium that mimicked human or animal gut microbiota elements.

We selected species found in gut microbiota of humans and animals (*Bacillus subtilis*, *Lactobacillus lactis* and *E. coli*). We included two strains of the human commensal gut species, *E. coli*. We thought it of interest to see if we could distinguish between closely related strains that nevertheless had sequence differences that we could use to target killing in one strain, but not the other. This was of interest as some strains of *E. coli* in microbiota are desirable, whereas others may be undesirable (eg. pathogenic to humans or animals) and thus could be targets for Cas modification to knock-down that strain.

1. Material and Methods

1.1. Plasmids and Strains

See Tables 7 and 8. All strains were cultivated in Todd-Hewitt broth (TH) (T1438 Sigma-Aldrich), in aerobic conditions and at 37° C., unless elsewhere indicated. The strains were stored in 25% glycerol at -80° C.

The self-targeting sgRNA-Cas9 complex was tightly regulated by a theophylline riboswitch and the AraC/P_{BAD} expression system respectively. Tight regulation of Cas9 is desired in order to be carried stably in *E. coli*. The plasmid contained the exogenous Cas9 from *Streptococcus pyogenes* with a single guide RNA (sgRNA) targeting *E. coli*'s K-12 strains. Therefore K-12 derived strains TOP10 was susceptible to double strand self-cleavage and consequent death when the system was activated. *E. coli* strains like Nissle don't have the same target sequence therefore they were unaffected by the sgRNA-Cas9 activity. See Tables 9-11 below, which show sequences used in Example 9. We chose a target sequence (ribosomal RNA-encoding sequence) that is conserved in the target cells and present in multiple copies (7 copies), which increased the chances of cutting host cell genomes in multiple places to promote killing using a single gRNA design.

FIG. 13 shows regulators controlling the expression of spCas9 and the self-targeting sgRNA targeting the ribosomal RNA subunit 16s.

1.2. Differential Growth Media

All strains were grown on TH media at 37° C. for 20 hours. Selective media for *B. subtilis* was TH media supplemented with 2.5 g l⁻¹ of 2-phenylethanol (PEA). PEA was added to the media and autoclaved at 121° C. for 15 minutes at 15 psi. Agar plates were prepared by adding 1.5% (wt/vol) agar to the corresponding media.

1.3. Cloning

E. coli (One Shot® ThermoFischer TOP10 Chemically Competent cells) was used in all subcloning procedures. PCR was carried out using Phusion™ polymerase. All PCR products were purified with Nucleospin™ Gel and PCR Clean-up by Macherey-Nagel™ following the manufacturer's protocol. The purified fragments were digested with restriction enzyme DpnI in 1× FD buffer with 1 µl enzyme in a total volume of 34 µl. The digested reaction was again purified with Nucleospin Gel and PCR Clean-up by Macherey-Nagel following the manufacturer's protocol. Gibson assembly was performed in 10 µl reactions following the manufacturer's protocol (NewEngland Biolab).

Plasmid DNA was prepared using Qiagen kits according to the manufacturer's instructions. Modifications for Gram-

positive strains included growing bacteria in a medium supplemented with 0.5% glycine and lysozyme to facilitate cell lysis.

1.4. Transformation

1.4.1 Electro-Competent *E. coli* Cells and Transformation

Commercially electrocompetent cells were used for cloning and the experiments (One Shot® ThermoFischer TOP10 electropotent *E. coli*). Electroporation was done using standard settings: 1800 V, 25 µF and 200 SI using an Electro Cell Manipulator (BTX Harvard Apparatus ECM630). Following the pulse, 1 ml LB-SOC media was added and the cells were incubated at 37° C. for 1 hour. The transformed cells were plated in LB-agar containing the corresponding antibiotics.

1.5. Activation of sgRNA-Cas9 in *E. coli* and Consortium Experiments.

E. coli TOP10 and Nissle both with the plasmid containing the sgRNA targeting the ribosomal RNA-encoding sequence of K-12 derived strains and the other bacteria were grown overnight in 3 ml of TH broth. The next day the cells were diluted to ~OD 0.5 and next 10-fold serially diluted in TH media and using a 96-well replicator (Mettler Toledo Liquidator™ 96) 4 µL volume drops were spotted on TH agar, TH agar with inducers (1% arabinose and 2 mM theophylline), TH agar supplemented with 2.5 g l⁻¹ PEA and MacConkey agar supplemented with 1% maltose. The plates were incubated for 20 h at 37° C. and the colony forming units (CFU) were calculated from triplicate measurements.

2. Results

2.1 Specific Targeting of *E. coli* Strains Using an Exogenous CRISPR-Cas9 System

We first tested if the system could differentiate between two *E. coli* strains by introducing the killing system in both *E. coli* TOP10 and Nissle.

2.1 Targeting of *E. coli* Using an Exogenous CRISPR-Cas9 System in a Mixed Culture

Serial dilutions of overnight cultures were done in duplicate for both *E. coli* strains, *B. subtilis*, *L. lactis*, and in triplicate for the mixed cultures. All strains were grown at 37° C. for 20 hours in selective plates with and without the inducers. Induction of the system activates the sgRNA-Cas9 targeting K-12 derived strains, while leaving intact the other bacteria.

Distinguishing the different bacteria from a mixed culture is important in order to determine cell numbers of the different species and determine the specific removal of a species. MacConkey agar selectively grows *E. coli*, PEA agar is a selective medium that is used for the isolation of gram-positive (*B. subtilis*) from gram-negative (*E. coli*). Additionally, we found that different concentrations of PEA partially inhibit the growth of other gram positives. 2.5 g l⁻¹ of PEA proved to selectively grow *B. subtilis* while limiting growth of *E. coli* and *L. lactis*.

FIG. 14 shows specific targeting of *E. coli* strain by the inducible, exogenous, vector-borne CRISPR-Cas system. The sgRNA target the genome of K-12 derived *E. coli* strain *E. coli* TOP10, while the other *E. coli* strain tested was unaffected.

FIG. 15 shows spot assay with serial dilutions of individual bacterial species used in this study and mixed culture in TH agar without induction of the CRISPR-Cas9 system.

FIG. 16 shows a spot assay of the dilution 10^3 on different selective media. TH with 2.5 g l^{-1} PEA is a selective media for *B. subtilis* alone. MacConkey supplemented with maltose is a selective and differential culture medium for bacteria designed to selectively isolate Gram-negative and enteric bacilli and differentiate them based on maltose fermentation. Therefore TOP10 Δ malK mutant makes white colonies on the plates while Nissle makes pink colonies; A is *E coli* Δ malK, B is *E coli* Nissle, C is *B subtilis*, D is *L lactis*, E is mixed culture; the images at MacConkey-/B and E appear pink; the images at MacConkey+/B and E appear pink. FIG. 17 shows selective growth of the bacteria used in this study on different media and selective plates. It can be seen that we clearly, selectively killed the target *E coli* strain ("*E coli*" on x-axis in FIG. 17) in the mixed population, whereas the other related strain ("*E coli*-Nissle") was not similarly killed

Killing of the target strain in the mixed population was 1000-fold in this experiment.

REFERENCES

- Zhang, X. Z., & Zhang, Y. H. P. (2011). Simple, fast and high-efficiency transformation system for directed evolution of cellulase in *Bacillus subtilis*. *Microbial Biotechnology*, 4(1), 98-105. <http://doi.org/10.1111/j.1751-7915.2010.00230.x>
- Wegmann, U., O'Connell-Motherway, M., Zomer, A., Buist, G., Shearman, C., Canchaya, C., . . . Kok, J. (2007). Complete genome sequence of the prototype lactic acid bacterium *Lactococcus lactis* subsp. *cremoris* MG1363. *Journal of Bacteriology*, 189(8), 3256-70. <http://doi.org/10.1128/JB.0.01768-06>

TABLE 1

EXAMPLE BACTERIA				
Abiotrophia	Acidocella	Actinomyces	Alkaliimnicola	Aquaspirillum
Abiotrophia	Acidocella	Actinomyces bovis	Alkaliimnicola	Aquaspirillum
defectiva	aminolytica	Actinomyces	ehrlrichii	polymorphum
Acaricomes	Acidocella facilis	denticolens	Alkaliphilus	Aquaspirillum
Acaricomes	Acidomonas	Actinomyces	Alkaliphilus	putridiconchylium
phytoseiuli	Acidomonas	europaeus	orelandii	Aquaspirillum
Acetitomaculum	methanolica	Actinomyces	Alkaliphilus	serpens
Acetitomaculum	Acidothermus	georgiae	transvaalensis	Aquimarina
ruminis	Acidothermus	Actinomyces	Allochrochromatium	Aquimarina
Acetivibrio	cellulolyticus	gerencseriae	Allochrochromatium	latercula
Acetivibrio	Acidovorax	Actinomyces	vinosum	Arcanobacterium
cellulolyticus	Acidovorax	hordeovulneris	Alloiococcus	Arcanobacterium
Acetivibrio	anthurii	Actinomyces	Alloiococcus otitis	haemolyticum
ethanolgignens	Acidovorax caeni	howellii	Allokutzneria	Arcanobacterium
Acetivibrio	Acidovorax	Actinomyces	Allokutzneria albata	pyogenes
multivorans	cattleyae	hyovaginalis	Altererythrobacter	Archangium
Acetoanaerobium	Acidovorax citrulli	Actinomyces	Altererythrobacter	Archangium
Acetoanaerobium	Acidovorax	israelii	ishigakiensis	gephyra
noterae	defluvii	Actinomyces	Altermonas	Arcobacter
Acetobacter	Acidovorax	johnsonii	Altermonas	Arcobacter butzleri
Acetobacter acetii	delafieldii	Actinomyces	haloplanktis	Arcobacter
Acetobacter	Acidovorax facilis	meyeri	Altermonas	cryaerophilus
cerevisiae	Acidovorax	Actinomyces	macleodii	Arcobacter
Acetobacter	konjaci	naeslundii	Alysiella	halophilus
cibinongensis	Acidovorax	Actinomyces neuui	Alysiella crassa	Arcobacter
Acetobacter	temperans	Actinomyces	Alysiella filiformis	nitrofigilis
estunensis	Acidovorax	odontolyticus	Aminobacter	Arcobacter
Acetobacter	valerianellae	Actinomyces oris	Aminobacter	skirrowii
fabarum	Acinetobacter	Actinomyces	aganoensis	Arhodomonas
Acetobacter	Acinetobacter	raidingae	Aminobacter	Arhodomonas
ghanensis	baumannii	Actinomyces	aminovorans	aquaeolei
Acetobacter	Acinetobacter	slackii	Aminobacter	Arsenophonus
indonesiensis	baylyi	Actinomyces	niiyataensis	Arsenophonus
Acetobacter	Acinetobacter	turicensis	Aminobacterium	nasoniae
lovaniensis	bouvetii	Actinomyces	Aminobacterium	Arthrobacter
Acetobacter	Acinetobacter	viscosus	mobile	Arthrobacter agilis
malorum	calcoaceticus	Actinoplanes	Aminomonas	Arthrobacter albus
Acetobacter	Acinetobacter	Actinoplanes	Aminomonas	Arthrobacter
nitrogenifigens	germeri	auranticolor	paucivorans	aurescens
Acetobacter oeni	Acinetobacter	Actinoplanes	Ammoniphilus	Arthrobacter
Acetobacter	haemolyticus	brasiliensis	Ammoniphilus	chlorophenolicus
orientalis	Acinetobacter	Actinoplanes	oxalaticus	Arthrobacter
Acetobacter	johnsonii	consettensis	Ammoniphilus	citreus
orleanensis	Acinetobacter junii	Actinoplanes	oxalivorans	Arthrobacter
Acetobacter	Acinetobacter	deccanensis	Amphibacillus	crystallopoietes
pasteurianus	lwofii	Actinoplanes	Amphibacillus	Arthrobacter
Acetobacter	Acinetobacter	derwentensis	xylanus	cumminsii
pomorum	parvus	Actinoplanes	Amphritea	Arthrobacter
Acetobacter	Acinetobacter	digitatis	Amphritea balenae	globiformis
senegalensis	radioresistens	Actinoplanes	Amphritea japonica	Arthrobacter
Acetobacter	Acinetobacter	durhamensis	Amycolatopsis	histidinovorans
xylinus	schindleri	Actinoplanes	Amycolatopsis alba	Arthrobacter ilicis
Acetobacterium	Acinetobacter soli	ferrugineus	Amycolatopsis	Arthrobacter luteus
Acetobacterium	Acinetobacter	Actinoplanes	albidoflavus	Arthrobacter
bakii	tandooi	globisporus	Amycolatopsis	methyloctrophus
Acetobacterium	Acinetobacter	Actinoplanes	azurea	Arthrobacter
carbinolicum	tjembergiae	humidus	Amycolatopsis	mysorensis

TABLE 1-continued

EXAMPLE BACTERIA				
<i>Acetobacterium</i>	<i>Acinetobacter</i>	<i>Actinoplanes</i>	<i>coloradensis</i>	<i>Arthrobacter</i>
<i>dehalogenans</i>	<i>towneri</i>	<i>italicus</i>	<i>Amycolatopsis</i>	<i>nicotianae</i>
<i>Acetobacterium</i>	<i>Acinetobacter</i>	<i>Actinoplanes</i>	<i>lurida</i>	<i>Arthrobacter</i>
<i>fimetarium</i>	<i>ursingii</i>	<i>liguriensis</i>	<i>Amycolatopsis</i>	<i>nicotinovorans</i>
<i>Acetobacterium</i>	<i>Acinetobacter</i>	<i>Actinoplanes</i>	<i>mediterranei</i>	<i>Arthrobacter</i>
<i>malicum</i>	<i>venetianus</i>	<i>lobatus</i>	<i>Amycolatopsis</i>	<i>oxydans</i>
<i>Acetobacterium</i>	<i>Acrocarpospora</i>	<i>Actinoplanes</i>	<i>rifamycinica</i>	<i>Arthrobacter</i>
<i>paludosum</i>	<i>Acrocarpospora</i>	<i>missouriensis</i>	<i>Amycolatopsis</i>	<i>pascens</i>
<i>Acetobacterium</i>	<i>corrugata</i>	<i>Actinoplanes</i>	<i>rubida</i>	<i>Arthrobacter</i>
<i>tundrae</i>	<i>Acrocarpospora</i>	<i>palleronii</i>	<i>Amycolatopsis</i>	<i>phenanthrenivorans</i>
<i>Acetobacterium</i>	<i>macrocephala</i>	<i>Actinoplanes</i>	<i>sulphurea</i>	<i>Arthrobacter</i>
<i>wieringae</i>	<i>Acrocarpospora</i>	<i>philippinensis</i>	<i>Amycolatopsis</i>	<i>polychromogenes</i>
<i>Acetobacterium</i>	<i>pleiomorpha</i>	<i>Actinoplanes</i>	<i>tolypomycina</i>	<i>Arthrobacter</i>
<i>woodii</i>	<i>Actibacter</i>	<i>rectilineatus</i>	<i>Anabaena</i>	<i>protophormiae</i>
<i>Acetofilamentum</i>	<i>Actibacter</i>	<i>Actinoplanes</i>	<i>Anabaena cylindrica</i>	<i>Arthrobacter</i>
<i>Acetofilamentum</i>	<i>sediminis</i>	<i>regularis</i>	<i>Anabaena flosaquae</i>	<i>psychrolactophilus</i>
<i>rigidum</i>	<i>Actinoalloteichus</i>	<i>Actinoplanes</i>	<i>Anabaena variabilis</i>	<i>Arthrobacter</i>
<i>Acetohalobium</i>	<i>Actinoalloteichus</i>	<i>teichomyceticus</i>	<i>Anaeroarcus</i>	<i>ramosus</i>
<i>Acetohalobium</i>	<i>cyanogriseus</i>	<i>Actinoplanes</i>	<i>Anaeroarcus</i>	<i>Arthrobacter</i>
<i>arabaticum</i>	<i>Actinoalloteichus</i>	<i>utahensis</i>	<i>burkinensis</i>	<i>sulfonivorans</i>
<i>Acetomicrobium</i>	<i>hymeniacidonis</i>	<i>Actinopolyspora</i>	<i>Anaerobaculum</i>	<i>Arthrobacter</i>
<i>Acetomicrobium</i>	<i>Actinoalloteichus</i>	<i>Actinopolyspora</i>	<i>Anaerobaculum</i>	<i>sulfureus</i>
<i>faecale</i>	<i>spitiensis</i>	<i>halophila</i>	<i>mobile</i>	<i>Arthrobacter</i>
<i>Acetomicrobium</i>	<i>Actinobacillus</i>	<i>Actinopolyspora</i>	<i>Anaerobiospirillum</i>	<i>uratoxydans</i>
<i>flavidum</i>	<i>Actinobacillus</i>	<i>mortivallis</i>	<i>Anaerobiospirillum</i>	<i>Arthrobacter</i>
<i>Acetonema</i>	<i>capsulatus</i>	<i>Actinosynnema</i>	<i>succiniciproducens</i>	<i>ureafaciens</i>
<i>Acetonema longum</i>	<i>Actinobacillus</i>	<i>Actinosynnema</i>	<i>Anaerobiospirillum</i>	<i>Arthrobacter</i>
<i>Acetothermus</i>	<i>delphinicola</i>	<i>mirum</i>	<i>thomasi</i>	<i>viscosus</i>
<i>Acetothermus</i>	<i>Actinobacillus</i>	<i>Actinotalea</i>	<i>Anaerococcus</i>	<i>Arthrobacter</i>
<i>paucivorans</i>	<i>hominis</i>	<i>Actinotalea</i>	<i>Anaerococcus</i>	<i>woluwenis</i>
<i>Acholeplasma</i>	<i>Actinobacillus</i>	<i>fermentans</i>	<i>hydrogenalis</i>	<i>Asaia</i>
<i>Acholeplasma</i>	<i>indolicus</i>	<i>Aerococcus</i>	<i>Anaerococcus</i>	<i>Asaia bogorensis</i>
<i>axanthum</i>	<i>Actinobacillus</i>	<i>Aerococcus</i>	<i>lactolyticus</i>	<i>Asanoa</i>
<i>Acholeplasma</i>	<i>lignieresii</i>	<i>sanguinicola</i>	<i>Anaerococcus</i>	<i>Asanoa ferruginea</i>
<i>brassicae</i>	<i>Actinobacillus</i>	<i>Aerococcus urinae</i>	<i>prevotii</i>	<i>Asticcacaulis</i>
<i>Acholeplasma</i>	<i>minor</i>	<i>Aerococcus</i>	<i>Anaerococcus</i>	<i>Asticcacaulis</i>
<i>cavigenitalium</i>	<i>Actinobacillus</i>	<i>urinaeequi</i>	<i>tetradus</i>	<i>biprosthecium</i>
<i>Acholeplasma</i>	<i>muris</i>	<i>Aerococcus</i>	<i>Anaerococcus</i>	<i>Asticcacaulis</i>
<i>equifetale</i>	<i>Actinobacillus</i>	<i>urinaehominis</i>	<i>vaginalis</i>	<i>excentricus</i>
<i>Acholeplasma</i>	<i>pleuropneumoniae</i>	<i>Aerococcus</i>	<i>Anaerofustis</i>	<i>Atopobacter</i>
<i>granularum</i>	<i>Actinobacillus</i>	<i>viridans</i>	<i>Anaerofustis</i>	<i>Atopobacter</i>
<i>Acholeplasma</i>	<i>porcinus</i>	<i>Aeromicrobium</i>	<i>stercorihominis</i>	<i>phocae</i>
<i>hippikon</i>	<i>Actinobacillus</i>	<i>Aeromicrobium</i>	<i>Anaeromusa</i>	<i>Atopobium</i>
<i>Acholeplasma</i>	<i>rossii</i>	<i>erythreum</i>	<i>Anaeromusa</i>	<i>Atopobium fossor</i>
<i>laidlawii</i>	<i>Actinobacillus</i>	<i>Aeromonas</i>	<i>acidaminophila</i>	<i>Atopobium</i>
<i>Acholeplasma</i>	<i>scotiae</i>	<i>Aeromonas</i>	<i>Anaeromyxobacter</i>	<i>minutum</i>
<i>modicum</i>	<i>Actinobacillus</i>	<i>allosaccharophila</i>	<i>Anaeromyxobacter</i>	<i>Atopobium</i>
<i>Acholeplasma</i>	<i>seminis</i>	<i>Aeromonas</i>	<i>dehalogenans</i>	<i>parvulum</i>
<i>morum</i>	<i>Actinobacillus</i>	<i>bestiarum</i>	<i>Anaerorhabdus</i>	<i>Atopobium rimae</i>
<i>Acholeplasma</i>	<i>succinogenes</i>	<i>Aeromonas caviae</i>	<i>Anaerorhabdus</i>	<i>Atopobium vaginae</i>
<i>multilocale</i>	<i>Actinobacillus</i>	<i>Aeromonas</i>	<i>furcosa</i>	<i>Aureobacterium</i>
<i>Acholeplasma</i>	<i>suis</i>	<i>encheleia</i>	<i>Anaerosinus</i>	<i>Aureobacterium</i>
<i>oculi</i>	<i>Actinobacillus</i>	<i>Aeromonas</i>	<i>Anaerosinus</i>	<i>barkeri</i>
<i>Acholeplasma</i>	<i>ureae</i>	<i>enteropelogenes</i>	<i>glycerini</i>	<i>Aurobacterium</i>
<i>palmae</i>	<i>Actinobaculum</i>	<i>Aeromonas</i>	<i>Anaerovirgula</i>	<i>Aurobacterium</i>
<i>Acholeplasma</i>	<i>Actinobaculum</i>	<i>eucrenophila</i>	<i>Anaerovirgula</i>	<i>liquefaciens</i>
<i>parvum</i>	<i>massiliense</i>	<i>Aeromonas</i>	<i>multivorans</i>	<i>Avibacterium</i>
<i>Acholeplasma</i>	<i>Actinobaculum</i>	<i>ichthiosmia</i>	<i>Ancalomicrobium</i>	<i>Avibacterium avium</i>
<i>pleciae</i>	<i>Actinobaculum</i>	<i>Aeromonas</i>	<i>Ancalomicrobium</i>	<i>Avibacterium</i>
<i>Acholeplasma</i>	<i>Actinobaculum</i>	<i>jandaei</i>	<i>adetum</i>	<i>gallinarum</i>
<i>vituli</i>	<i>suis</i>	<i>Aeromonas media</i>	<i>Ancylobacter</i>	<i>Avibacterium</i>
<i>Achromobacter</i>	<i>Actinomyces</i>	<i>Aeromonas</i>	<i>Ancylobacter</i>	<i>paragallinarum</i>
<i>Achromobacter</i>	<i>urinale</i>	<i>popoffii</i>	<i>aquaticus</i>	<i>Avibacterium</i>
<i>denitrificans</i>	<i>Actinocatenispora</i>	<i>Aeromonas sobria</i>	<i>Aneurinibacillus</i>	<i>volantium</i>
<i>Achromobacter</i>	<i>Actinocatenispora</i>	<i>Aeromonas veronii</i>	<i>Aneurinibacillus</i>	<i>Azoarcus</i>
<i>insolitus</i>	<i>rupis</i>	<i>Agrobacterium</i>	<i>aneurinilyticus</i>	<i>Azoarcus indigenus</i>
<i>Achromobacter</i>	<i>Actinocatenispora</i>	<i>Agrobacterium</i>	<i>Aneurinibacillus</i>	<i>Azoarcus</i>
<i>piechaudii</i>	<i>thailandica</i>	<i>gelatinovorum</i>	<i>migulanus</i>	<i>tolulyticus</i>
<i>Achromobacter</i>	<i>Actinocatenispora</i>	<i>Agrococcus</i>	<i>Aneurinibacillus</i>	<i>Azoarcus</i>
<i>ruhlandii</i>	<i>sera</i>	<i>Agrococcus</i>	<i>thermoaerophilus</i>	<i>toluovorans</i>
<i>Achromobacter</i>	<i>Actinocorallia</i>	<i>citreus</i>	<i>Angiococcus</i>	<i>Azohydromonas</i>
<i>spanius</i>	<i>Actinocorallia</i>	<i>Agrococcus</i>	<i>Angiococcus</i>	<i>Azohydromonas</i>
<i>Acidaminobacter</i>	<i>aurantiaca</i>	<i>jenensis</i>	<i>disciformis</i>	<i>australiana</i>
<i>Acidaminobacter</i>	<i>Actinocorallia</i>	<i>Agromonas</i>	<i>Angulomicrobium</i>	<i>Azohydromonas</i>
<i>hydrogeniformans</i>	<i>aurea</i>	<i>Agromonas</i>	<i>Angulomicrobium</i>	<i>lata</i>
<i>Acidaminococcus</i>	<i>Actinocorallia</i>	<i>oligotrophica</i>	<i>tetraedrale</i>	<i>Azomonas</i>
<i>Acidaminococcus</i>	<i>cavernae</i>	<i>Agromyces</i>	<i>Anoxybacillus</i>	<i>Azomonas agilis</i>
<i>fermentans</i>	<i>Actinocorallia</i>	<i>Agromyces</i>	<i>Anoxybacillus</i>	<i>Azomonas insignis</i>

TABLE 1-continued

EXAMPLE BACTERIA				
<i>Acidaminococcus</i>	<i>glomerata</i>	<i>fucosus</i>	<i>pushchinoensis</i>	<i>Azomonas</i>
<i>intestini</i>	<i>Actinocorallia</i>	<i>Agromyces</i>	<i>Aquabacterium</i>	<i>macrocytogenes</i>
<i>Acidicaldus</i>	<i>herbida</i>	<i>hippuratus</i>	<i>Aquabacterium</i>	<i>Azorhizobium</i>
<i>Acidicaldus</i>	<i>Actinocorallia</i>	<i>Agromyces</i>	<i>commune</i>	<i>Azorhizobium</i>
<i>organivorans</i>	<i>libanotica</i>	<i>luteolus</i>	<i>Aquabacterium</i>	<i>caulinodans</i>
<i>Acidimicrobium</i>	<i>Actinocorallia</i>	<i>Agromyces</i>	<i>parvum</i>	<i>Azorhizophilus</i>
<i>Acidimicrobium</i>	<i>longicatena</i>	<i>mediolanus</i>		<i>Azorhizophilus</i>
<i>ferrooxidans</i>	<i>Actinomadura</i>	<i>Agromyces</i>		<i>paspali</i>
<i>Acidiphilium</i>	<i>Actinomadura alba</i>	<i>ramosus</i>		<i>Azospirillum</i>
<i>Acidiphilium</i>	<i>Actinomadura</i>	<i>Agromyces</i>		<i>Azospirillum</i>
<i>acidophilum</i>	<i>atramentaria</i>	<i>rhizospherae</i>		<i>brasilense</i>
<i>Acidiphilium</i>	<i>Actinomadura</i>	<i>Akkermansia</i>		<i>Azospirillum</i>
<i>angustum</i>	<i>bangladeshensis</i>	<i>Akkermansia</i>		<i>halopraeferens</i>
<i>Acidiphilium</i>	<i>Actinomadura</i>	<i>muciniphila</i>		<i>Azospirillum</i>
<i>cryptum</i>	<i>catellatispora</i>	<i>Albidiferax</i>		<i>irakense</i>
<i>Acidiphilium</i>	<i>Actinomadura</i>	<i>Albidiferax</i>		<i>Azotobacter</i>
<i>multivorum</i>	<i>chibensis</i>	<i>ferrireducens</i>		<i>Azotobacter</i>
<i>Acidiphilium</i>	<i>Actinomadura</i>	<i>Albidovulum</i>		<i>beijerinckii</i>
<i>organovorum</i>	<i>chokoviensis</i>	<i>Albidovulum</i>		<i>Azotobacter</i>
<i>Acidiphilium</i>	<i>Actinomadura</i>	<i>inexpectatum</i>		<i>chroococcum</i>
<i>rubrum</i>	<i>citrea</i>	<i>Alcaligenes</i>		<i>Azotobacter</i>
<i>Acidisoma</i>	<i>Actinomadura</i>	<i>Alcaligenes</i>		<i>nigricans</i>
<i>Acidisoma</i>	<i>coerulea</i>	<i>denitrificans</i>		<i>Azotobacter</i>
<i>sibiricum</i>	<i>Actinomadura</i>	<i>Alcaligenes</i>		<i>salinestris</i>
<i>Acidisoma tundrae</i>	<i>echinospora</i>	<i>faecalis</i>		<i>Azotobacter</i>
<i>Acidisphaera</i>	<i>Actinomadura</i>	<i>Alcanivorax</i>		<i>vinelandii</i>
<i>Acidisphaera</i>	<i>fibrosa</i>	<i>Alcanivorax</i>		
<i>rubrifaciens</i>	<i>Actinomadura</i>	<i>borkumensis</i>		
<i>Acidithiobacillus</i>	<i>formosensis</i>	<i>Alcanivorax</i>		
<i>Acidithiobacillus</i>	<i>Actinomadura</i>	<i>jadensis</i>		
<i>albertensis</i>	<i>hibisca</i>	<i>Algicola</i>		
<i>Acidithiobacillus</i>	<i>Actinomadura</i>	<i>Algicola</i>		
<i>caldus</i>	<i>kijaniata</i>	<i>bacteriolytica</i>		
<i>Acidithiobacillus</i>	<i>Actinomadura</i>	<i>Alicyclobacillus</i>		
<i>ferrooxidans</i>	<i>Actinomadura</i>	<i>Alicyclobacillus</i>		
<i>Acidithiobacillus</i>	<i>Actinomadura</i>	<i>disulfidooxidans</i>		
<i>thiooxidans</i>	<i>livida</i>	<i>Alicyclobacillus</i>		
<i>Acidobacterium</i>	<i>Actinomadura</i>	<i>sendaiensis</i>		
<i>Acidobacterium</i>	<i>luteofluorescens</i>	<i>Alicyclobacillus</i>		
<i>capsulatum</i>	<i>Actinomadura</i>	<i>vulcanalis</i>		
	<i>macra</i>	<i>Alisshewanella</i>		
	<i>Actinomadura</i>	<i>Alisshewanella</i>		
	<i>madurae</i>	<i>fetalis</i>		
	<i>Actinomadura</i>	<i>Alkalibacillus</i>		
	<i>oligospora</i>	<i>Alkalibacillus</i>		
	<i>Actinomadura</i>	<i>haloalkaliphilus</i>		
	<i>pelletieri</i>			
	<i>Actinomadura</i>			
	<i>rubrobrunea</i>			
	<i>Actinomadura</i>			
	<i>rugatobispora</i>			
	<i>Actinomadura</i>			
	<i>umbrina</i>			
	<i>Actinomadura</i>			
	<i>verrucosospora</i>			
	<i>Actinomadura</i>			
	<i>vinacea</i>			
	<i>Actinomadura</i>			
	<i>viridilutea</i>			
	<i>Actinomadura</i>			
	<i>viridis</i>			
	<i>Actinomadura</i>			
	<i>yumaensis</i>			
Bacillus	Bacteroides	Bibersteinia	Borrelia	Brevinema
[see below]	<i>Bacteroides</i>	<i>Bibersteinia trehalosi</i>	<i>Borrelia afzelii</i>	<i>Brevinema</i>
Bacteriovorax	<i>caccae</i>	Bifidobacterium	<i>Borrelia americana</i>	<i>andersonii</i>
Bacteriovorax	<i>Bacteroides</i>	<i>Bifidobacterium</i>	<i>Borrelia</i>	Brevundimonas
<i>stolpii</i>	<i>coagulans</i>	<i>adolescentis</i>	<i>burgdorferi</i>	<i>Brevundimonas</i>
	<i>Bacteroides</i>	<i>Bifidobacterium</i>	<i>Borrelia</i>	<i>alba</i>
	<i>eggerthii</i>	<i>angulatum</i>	<i>carolinensis</i>	<i>Brevundimonas</i>
	<i>Bacteroides</i>	<i>Bifidobacterium</i>	<i>Borrelia coriaceae</i>	<i>aurantiaca</i>
	<i>fragilis</i>	<i>animalis</i>	<i>Borrelia garinii</i>	<i>Brevundimonas</i>
	<i>Bacteroides</i>	<i>Bifidobacterium</i>	<i>Borrelia japonica</i>	<i>diminuta</i>
	<i>galacturonicus</i>	<i>asteroides</i>	Bosea	<i>Brevundimonas</i>
	<i>Bacteroides</i>	<i>Bifidobacterium</i>	<i>Bosea</i>	<i>intermedia</i>
	<i>helcogenes</i>	<i>bifidum</i>	<i>minatitlanensis</i>	<i>Brevundimonas</i>

TABLE 1-continued

EXAMPLE BACTERIA

<i>Bacteroides ovatus</i>	<i>Bifidobacterium boum</i>	<i>Bosea thiooxidans</i>	<i>subvibrioides</i>
<i>Bacteroides pectinophilus</i>	<i>Bifidobacterium breve</i>	Brachybacterium	<i>Brevundimonas vancomneytii</i>
<i>Bacteroides pyogenes</i>	<i>Bifidobacterium caterulatum</i>	<i>Brachybacterium alimentarium</i>	<i>Brevundimonas variabilis</i>
<i>Bacteroides salyersiae</i>	<i>Bifidobacterium choerinum</i>	<i>Brachybacterium faecium</i>	<i>Brevundimonas vesicularis</i>
<i>Bacteroides stercoris</i>	<i>Bifidobacterium coryneforme</i>	<i>Brachybacterium paraconglomeratum</i>	Brochothrix
<i>Bacteroides suis</i>	<i>Bifidobacterium cuniculi</i>	<i>Brachybacterium rhamnosum</i>	<i>Brochothrix campestris</i>
<i>Bacteroides tectus</i>	<i>Bifidobacterium dentium</i>	<i>Brachybacterium tyrofermentans</i>	<i>Brochothrix thermosphacta</i>
<i>Bacteroides thetaiotaomicron</i>	<i>Bifidobacterium gallicum</i>	Brachyspira	Brucella
<i>Bacteroides uniformis</i>	<i>Bifidobacterium gallinarum</i>	<i>Brachyspira alvinipulli</i>	<i>Brucella canis</i>
<i>Bacteroides ureolyticus</i>	<i>Bifidobacterium indicum</i>	<i>Brachyspira brachyspira</i>	<i>Brucella neotomae</i>
<i>Bacteroides vulgatus</i>	<i>Bifidobacterium longum</i>	<i>Brachyspira hyodysenteriae</i>	Bryobacter
Balnearium	<i>Bifidobacterium magnum</i>	<i>Brachyspira innocens</i>	<i>Bryobacter aggregatus</i>
Balnearium lithotrophicum	<i>Bifidobacterium merycicum</i>	<i>Brachyspira pilosicoli</i>	Burkholderia
Balneatrix	<i>Bifidobacterium minimum</i>	Bradyrhizobium	<i>Burkholderia ambifaria</i>
<i>Balneatrix alpica</i>	<i>Bifidobacterium pseudocatenulatum</i>	<i>Bradyrhizobium canariense</i>	<i>Burkholderia andropogonis</i>
Balneola	<i>Bifidobacterium pseudolongum</i>	<i>Bradyrhizobium elkanii</i>	<i>Burkholderia anthina</i>
<i>Balneola vulgaris</i>	<i>Bifidobacterium pullorum</i>	<i>Bradyrhizobium japonicum</i>	<i>Burkholderia caledonica</i>
Barnesiella	<i>Bifidobacterium ruminantium</i>	<i>Bradyrhizobium liaoningense</i>	<i>Burkholderia caryophylli</i>
<i>Barnesiella viscericola</i>	<i>Bifidobacterium saeculare</i>	Brenneria	<i>Burkholderia cenocepacia</i>
Bartonella	<i>Bifidobacterium subtile</i>	<i>Brenneria alni</i>	<i>Burkholderia cepacia</i>
<i>Bartonella alsatica</i>	<i>Bifidobacterium thermophilum</i>	<i>Brenneria quercina</i>	<i>Burkholderia cocovenenans</i>
<i>Bartonella bacilliformis</i>	<i>Bifidobacterium bilophila</i>	<i>Brenneria quercina salicis</i>	<i>Burkholderia dolosa</i>
<i>Bartonella clarridgeiae</i>	<i>Bifidobacterium biostraticola</i>	Brevibacillus	<i>Burkholderia fungorum</i>
<i>Bartonella doshaiae</i>	<i>Bifidobacterium biostraticola tofi</i>	<i>Brevibacillus agrif</i>	<i>Burkholderia glumae</i>
<i>Bartonella elizabethae</i>	Bizionia	<i>Brevibacillus borstelensis</i>	<i>Burkholderia gramini</i>
<i>Bartonella grahamii</i>	<i>Bizionia argentinensis</i>	<i>Brevibacillus brevis</i>	<i>Burkholderia kururiensis</i>
<i>Bartonella henselae</i>	Blastobacter	<i>Brevibacillus invocatus</i>	<i>Burkholderia multivorans</i>
<i>Bartonella rochalimae</i>	<i>Blastobacter capsulatus</i>	<i>Brevibacillus laterosporus</i>	<i>Burkholderia phenazinium</i>
<i>Bartonella vinsonii</i>	<i>Blastobacter denitrificans</i>	<i>Brevibacillus parabrevis</i>	<i>Burkholderia plantarii</i>
Bavariococcus	Blastococcus	<i>Brevibacillus reuszeri</i>	<i>Burkholderia pyrrocinia</i>
<i>Bavariococcus seileri</i>	<i>Blastococcus saxosidens</i>	Brevibacterium	<i>Burkholderia silvaticola</i>
Bdellovibrio	<i>Blastococcus aggregatus</i>	<i>Brevibacterium abidum</i>	<i>Burkholderia stabilis</i>
<i>Bdellovibrio bacteriovorus</i>	<i>Blastococcus saxosidens</i>	<i>Brevibacterium album</i>	<i>Burkholderia thailandensis</i>
<i>Bdellovibrio exovorax</i>	<i>Blastochloris viridis</i>	<i>Brevibacterium aurantiacum</i>	<i>Burkholderia tropica</i>
Beggiatoa	<i>Blastomonas natatoria</i>	<i>Brevibacterium celere</i>	<i>Burkholderia unamae</i>
<i>Beggiatoa alba</i>	<i>Blastopirellula marina</i>	<i>Brevibacterium epidermidis</i>	<i>Burkholderia vietnamiensis</i>
Beijerinckia	<i>Blastopirellula blautia</i>	<i>Brevibacterium frigiditolerans</i>	Buttiauxella
<i>Beijerinckia derxii</i>	<i>Blastopirellula blautia coccoides</i>	<i>Brevibacterium halotolerans</i>	<i>Buttiauxella agrestis</i>
<i>Beijerinckia fluminensis</i>	<i>Blastopirellula blautia producta</i>	<i>Brevibacterium iodinum</i>	<i>Buttiauxella brennerae</i>
<i>Beijerinckia indica</i>	<i>Blastopirellula bogoriella</i>	<i>Brevibacterium linens</i>	<i>Buttiauxella ferrugutiae</i>
<i>Beijerinckia mobilis</i>	<i>Blastopirellula caseolytica</i>	<i>Brevibacterium lyticum</i>	<i>Buttiauxella gaviniae</i>
Belliella	<i>Blastopirellula bordetella</i>		
<i>Belliella baltica</i>	<i>Blastopirellula bronchiseptica</i>		
Bellilinea	<i>Blastopirellula bordetella hinzii</i>		
<i>Bellilinea caldifistulae</i>	<i>Blastopirellula bordetella holmesii</i>		
Belnapia	<i>Blastopirellula bordetella parapertussis</i>		
<i>Belnapia moabensis</i>	<i>Blastopirellula bordetella pertussis</i>		

TABLE 1-continued

EXAMPLE BACTERIA			
Bergeriella	<i>Bordetella petrii</i>	<i>Brevibacterium</i>	<i>Buttiauxella</i>
<i>Bergeriella</i>	<i>Bordetella trematum</i>	<i>mcbrellneri</i>	<i>izardii</i>
<i>denitrificans</i>		<i>Brevibacterium</i>	<i>Buttiauxella</i>
Beutenbergia		<i>otitidis</i>	<i>noackiae</i>
<i>Beutenbergia</i>		<i>Brevibacterium</i>	<i>Buttiauxella</i>
<i>cavernae</i>		<i>oxydans</i>	<i>warmboldiae</i>
		<i>Brevibacterium</i>	Butyrivibrio
		<i>paucivorans</i>	Butyrivibrio
		<i>Brevibacterium</i>	<i>fibrisolvens</i>
		<i>stationis</i>	Butyrivibrio
			<i>hungatei</i>
			Butyrivibrio
			<i>proteoclasticus</i>
<hr/>			
Bacillus			
<i>B. acidicer</i>	<i>B. aminovorans</i>	<i>B. glucanolyticus</i>	<i>B. lautus</i>
<i>B. acidicola</i>	<i>B. amylolyticus</i>	<i>B. gordonae</i>	<i>B. lehenis</i>
<i>B. acidiproducens</i>	<i>B. andreesenii</i>	<i>B. gottheilii</i>	<i>B. lentimorbus</i>
<i>B. acidocaldarius</i>	<i>B. aneurinilyticus</i>	<i>B. graminis</i>	<i>B. lentus</i>
<i>B. acidoterrestris</i>	<i>B. anthracis</i>	<i>B. halmopalus</i>	<i>B. licheniformis</i>
<i>B. aeolius</i>	<i>B. aquimaris</i>	<i>B. haloalkaliphilus</i>	<i>B. ligniniphilus</i>
<i>B. aerius</i>	<i>B. arenosi</i>	<i>B. halocharae</i>	<i>B. litoralis</i>
<i>B. aerophilus</i>	<i>B. arseniciselenatis</i>	<i>B. halodenitrificans</i>	<i>B. locisalis</i>
<i>B. agaradhaerens</i>	<i>B. arsenicus</i>	<i>B. haloduvans</i>	<i>B. luciferensis</i>
<i>B. agri</i>	<i>B. aurantiacus</i>	<i>B. halophilus</i>	<i>B. luteolus</i>
<i>B. aidingensis</i>	<i>B. arvi</i>	<i>B. halosaccharovorans</i>	<i>B. luteus</i>
<i>B. akibai</i>	<i>B. aryabhatai</i>	<i>B. hemicellulosilyticus</i>	<i>B. macauensis</i>
<i>B. alcalophilus</i>	<i>B. asahii</i>	<i>B. hemicentroti</i>	<i>B. macerans</i>
<i>B. algicola</i>	<i>B. atrophaeus</i>	<i>B. herbersteinensis</i>	<i>B. macquariensis</i>
<i>B. alginolyticus</i>	<i>B. axarquiensis</i>	<i>B. horikoshii</i>	<i>B. macyae</i>
<i>B. alkalidiazotrophicus</i>	<i>B. azotofixans</i>	<i>B. horneckiae</i>	<i>B. malacitensis</i>
<i>B. alkalinitrilicus</i>	<i>B. azotoformans</i>	<i>B. horti</i>	<i>B. mammaryticus</i>
<i>B. alkalisediminis</i>	<i>B. badius</i>	<i>B. huizhouensis</i>	<i>B. marisflavi</i>
<i>B. alkalitelluris</i>	<i>B. barbaricus</i>	<i>B. humi</i>	<i>B. marismortui</i>
<i>B. altitudinis</i>	<i>B. bataviensis</i>	<i>B. hwajinpoensis</i>	<i>B. marmarensis</i>
<i>B. alveayuensis</i>	<i>B. beijingensis</i>	<i>B. idriensis</i>	<i>B. massiliensis</i>
<i>B. alvei</i>	<i>B. benzoovorans</i>	<i>B. indicus</i>	<i>B. megaterium</i>
<i>B. amyloliquefaciens</i>	<i>B. beringensis</i>	<i>B. infantis</i>	<i>B. mesonae</i>
<i>B.a. subsp. amyloliquefaciens</i>	<i>B. berkeleyi</i>	<i>B. infernus</i>	<i>B. methanolicus</i>
<i>B.a. subsp. plantarum</i>	<i>B. beveridgei</i>	<i>B. insolitus</i>	<i>B. methylotrophicus</i>
<i>B. dipsosauri</i>	<i>B. bogoriensis</i>	<i>B. invictae</i>	<i>B. migulanus</i>
<i>B. drentensis</i>	<i>B. boroniphilus</i>	<i>B. iranensis</i>	<i>B. mojavensis</i>
<i>B. edaphicus</i>	<i>B. borstelensis</i>	<i>B. isabelliae</i>	<i>B. mucilaginosus</i>
<i>B. chimensis</i>	<i>B. brevis Migula</i>	<i>B. isronensis</i>	<i>B. muralis</i>
<i>B. eiseniae</i>	<i>B. butanolivorans</i>	<i>B. jeotgali</i>	<i>B. murimartini</i>
<i>B. enclensis</i>	<i>B. canaveralius</i>	<i>B. kaustophilus</i>	<i>B. mycoides</i>
<i>B. endophyticus</i>	<i>B. carboniphilus</i>	<i>B. kobensis</i>	<i>B. naganensis</i>
<i>B. endoradicis</i>	<i>B. cecembensis</i>	<i>B. kochii</i>	<i>B. nanhaiensis</i>
<i>B. farraginis</i>	<i>B. cellulosityticus</i>	<i>B. kokeshiiformis</i>	<i>B. nanhaisediminis</i>
<i>B. fastidiosus</i>	<i>B. centrosporus</i>	<i>B. koreensis</i>	<i>B. nealsonii</i>
<i>B. fengquensis</i>	<i>B. cereus</i>	<i>B. korlensis</i>	<i>B. neidei</i>
<i>B. firmus</i>	<i>B. chagannorensis</i>	<i>B. kribbensis</i>	<i>B. neizhouensis</i>
<i>B. flexus</i>	<i>B. chitinolyticus</i>	<i>B. krulwichiae</i>	<i>B. niabensis</i>
<i>B. foraminis</i>	<i>B. chondroitinus</i>	<i>B. laevolacticus</i>	<i>B. niacini</i>
<i>B. fordii</i>	<i>B. choshinensis</i>	<i>B. larvae</i>	<i>B. novalis</i>
<i>B. formosus</i>	<i>B. chungangensis</i>	<i>B. laterosporus</i>	<i>B. oceanisediminis</i>
<i>B. fortis</i>	<i>B. cibi</i>	<i>B. salexigens</i>	<i>B. odysseyi</i>
<i>B. fumarioli</i>	<i>B. circulans</i>	<i>B. saliphilus</i>	<i>B. okhensis</i>
<i>B. fimiculus</i>	<i>B. clarkii</i>	<i>B. schlegelii</i>	<i>B. okuhidensis</i>
<i>B. fusiformis</i>	<i>B. clausii</i>	<i>B. sediminis</i>	<i>B. oleronius</i>
<i>B. galactophilus</i>	<i>B. coagulans</i>	<i>B. selenatarsenatis</i>	<i>B. oryzaecorticis</i>
<i>B. galactosidilyticus</i>	<i>B. coahuilensis</i>	<i>B. selenitireducens</i>	<i>B. oshimensis</i>
<i>B. galliensis</i>	<i>B. cohuii</i>	<i>B. seohaeanensis</i>	<i>B. pabuli</i>
<i>B. gelatini</i>	<i>B. composti</i>	<i>B. shacheensis</i>	<i>B. pakistanensis</i>
<i>B. gibsonii</i>	<i>B. curdlanolyticus</i>	<i>B. shackletonii</i>	<i>B. pallidus</i>
<i>B. ginsengi</i>	<i>B. cycloheptanicus</i>	<i>B. siamensis</i>	<i>B. pallidus</i>
<i>B. ginsengihumi</i>	<i>B. cytotoxicus</i>	<i>B. silvestris</i>	<i>B. panacisoli</i>
<i>B. ginsengisoli</i>	<i>B. daliensis</i>	<i>B. simplex</i>	<i>B. panaciterrae</i>
<i>B. globisporus</i>	<i>B. decisifrontis</i>	<i>B. siralis</i>	<i>B. pantothenicus</i>
(eg. <i>B.g. subsp.</i>	<i>B. decolorationis</i>	<i>B. smithii</i>	<i>B. parabravis</i>
<i>Globisporus</i> ;	<i>B. deserti</i>	<i>B. soli</i>	<i>B. paraflexus</i>
or <i>B.g. subsp.</i>		<i>B. solimangrovi</i>	<i>B. pasteurii</i>
<i>Marinus</i>)		<i>B. solisalsi</i>	<i>B. patagoniensis</i>
		<i>B. songklensis</i>	
		<i>B. sonorensis</i>	
		<i>B. sphaericus</i>	
		<i>B. sporothermodurans</i>	
		<i>B. stearothermophilus</i>	
		<i>B. taeanensis</i>	
		<i>B. tequilensis</i>	
		<i>B. thermantarcticus</i>	
		<i>B. thermoaerophilus</i>	
		<i>B. thermoamylovorans</i>	
		<i>B. thermocatenuatus</i>	
		<i>B. thermocloacae</i>	
		<i>B. thermocopriae</i>	
		<i>B. thermodenitrificans</i>	
		<i>B. thermoglucosidasius</i>	
		<i>B. thermolactis</i>	
		<i>B. thermoleovorans</i>	
		<i>B. thermophilus</i>	
		<i>B. thermoruber</i>	
		<i>B. thermosphaericus</i>	
		<i>B. thiaminolyticus</i>	
		<i>B. thioparans</i>	
		<i>B. thuringiensis</i>	
		<i>B. tianshenii</i>	
		<i>B. tryoxylicola</i>	
		<i>B. tusciae</i>	
		<i>B. validus</i>	
		<i>B. vallismortis</i>	
		<i>B. vedderi</i>	
		<i>B. velezensis</i>	
		<i>B. vietnamensis</i>	
		<i>B. vireti</i>	
		<i>B. vulcani</i>	
		<i>B. wakoensis</i>	
		<i>B. weihenstephanensis</i>	
		<i>B. xiamenensis</i>	
		<i>B. xiaoxiensis</i>	
		<i>B. zhanjiangensis</i>	
		<i>B. peoriae</i>	
		<i>B. persepolensis</i>	
		<i>B. persicus</i>	
		<i>B. pervagus</i>	
		<i>B. plakortidis</i>	
		<i>B. pocheonensis</i>	
		<i>B. polygoni</i>	
		<i>B. polymyxa</i>	
		<i>B. popilliae</i>	
		<i>B. pseudocalophilus</i>	
		<i>B. pseudofirmus</i>	
		<i>B. pseudomycooides</i>	
		<i>B. psychrodurans</i>	
		<i>B. psychrophilus</i>	
		<i>B. psychrosaccharolyticus</i>	
		<i>B. psychrotolerans</i>	
		<i>B. pulvificiens</i>	
		<i>B. pumilus</i>	
		<i>B. purgationiresistens</i>	
		<i>B. pycnus</i>	
		<i>B. qingdaonensis</i>	
		<i>B. qingshengii</i>	
		<i>B. reuszeri</i>	
		<i>B. rhizosphaerae</i>	
		<i>B. rigui</i>	
		<i>B. ruris</i>	
		<i>B. safensis</i>	
		<i>B. salarius</i>	

TABLE 1-continued

EXAMPLE BACTERIA				
	<i>B. stratosphericus</i>			
	<i>B. subterraneus</i>			
	<i>B. subtilis</i>			
	(eg. B.s. subsp.			
	<i>Inaquosorum</i> ;			
	or B.s. subsp.			
	<i>Spizizeni</i> ;			
	or B.s. subsp.			
	<i>Subtilis</i>)			
Caenimonas	Campylobacter	Cardiobacterium	Catenuloplanes	Curtobacterium
<i>Caenimonas koreensis</i>	<i>Campylobacter coli</i>	<i>Cardiobacterium</i>	<i>Catenuloplanes</i>	<i>Curtobacterium</i>
Caldalkalibacillus	<i>Campylobacter</i>	<i>hominis</i>	<i>atrovinosus</i>	<i>albidum</i>
<i>Caldalkalibacillus</i>	<i>concisus</i>	Carnimonas	<i>Catenuloplanes</i>	<i>Curtobacterium</i>
<i>uzonensis</i>	<i>Campylobacter</i>	<i>Carnimonas</i>	<i>castaneus</i>	<i>citreus</i>
Caldanaerobacter	<i>curvus</i>	<i>nigrificans</i>	<i>Catenuloplanes</i>	
<i>Caldanaerobacter</i>	<i>Campylobacter fetus</i>	Carnobacterium	<i>crispus</i>	
<i>subterraneus</i>	<i>Campylobacter</i>	<i>Carnobacterium</i>	<i>Catenuloplanes</i>	
Caldanaerobius	<i>gracilis</i>	<i>alterfunditum</i>	<i>indicus</i>	
<i>Caldanaerobius</i>	<i>Campylobacter</i>	<i>Carnobacterium</i>	<i>Catenuloplanes</i>	
<i>fijiensis</i>	<i>helveticus</i>	<i>divergens</i>	<i>japonicus</i>	
<i>Caldanaerobius</i>	<i>Campylobacter</i>	<i>Carnobacterium</i>	<i>Catenuloplanes</i>	
<i>polysaccharolyticus</i>	<i>hominis</i>	<i>funditum</i>	<i>nepalensis</i>	
<i>Caldanaerobius zeae</i>	<i>Campylobacter</i>	<i>Carnobacterium</i>	<i>Catenuloplanes</i>	
Caldanaerovirga	<i>hyointestinalis</i>	<i>gallinarum</i>	<i>niger</i>	
<i>Caldanaerovirga</i>	<i>Campylobacter</i>	<i>Carnobacterium</i>	Chryseobacterium	
<i>acetigignens</i>	<i>jejuni</i>	<i>maltaromaticum</i>	Chryseobacterium	
Caldicellulosiruptor	<i>Campylobacter lari</i>	<i>Carnobacterium</i>	<i>balustinum</i>	
<i>Caldicellulosiruptor</i>	<i>mucosalis</i>	<i>mobile</i>	Citrobacter	
<i>besicii</i>	<i>Campylobacter</i>	<i>Carnobacterium</i>	<i>C. amalonaticus</i>	
<i>Caldicellulosiruptor</i>	<i>rectus</i>	<i>viridans</i>	<i>C. braakii</i>	
<i>kristjanssonii</i>	<i>Campylobacter</i>	Caryophanon	<i>C. diversus</i>	
<i>Caldicellulosiruptor</i>	<i>showae</i>	<i>Caryophanon</i>	<i>C. farmeri</i>	
<i>owensensis</i>	<i>Campylobacter</i>	<i>latum</i>	<i>C. freundii</i>	
	<i>Campylobacter</i>	<i>Caryophanon</i>	<i>C. gillenii</i>	
	<i>sputorum</i>	<i>tenue</i>	<i>C. koseri</i>	
	<i>Campylobacter</i>	Catellatospora	<i>C. murliniaie</i>	
	<i>upsaliensis</i>	<i>Catellatospora</i>	<i>C. pasteurii</i> ^[1]	
	Capnocytophaga	<i>citrea</i>	<i>C. rodentium</i>	
	<i>Capnocytophaga</i>	<i>Catellatospora</i>	<i>C. sedlakii</i>	
	<i>canimorsus</i>	<i>methionotrophica</i>	<i>C. werkmanii</i>	
	<i>Capnocytophaga</i>	Catenococcus	<i>C. youngae</i>	
	<i>cynodegmi</i>	<i>Catenococcus</i>	Clostridium	
	<i>Capnocytophaga</i>	<i>thiocycli</i>	(see below)	
	<i>gingivalis</i>		Coccochloris	
	<i>Capnocytophaga</i>		<i>Coccochloris</i>	
	<i>granulosa</i>		<i>elabens</i>	
	<i>Capnocytophaga</i>		Corynebacterium	
	<i>haemolytica</i>		<i>Corynebacterium</i>	
	<i>Capnocytophaga</i>		<i>flavescens</i>	
	<i>ochracea</i>		<i>Corynebacterium</i>	
	<i>Capnocytophaga</i>		<i>variabile</i>	
	<i>sputigena</i>			
Clostridium				
<i>Clostridium absonum</i> , <i>Clostridium aceticum</i> , <i>Clostridium acetireducens</i> , <i>Clostridium acetobutylicum</i> ,				
<i>Clostridium acidisoli</i> , <i>Clostridium aciditolerans</i> , <i>Clostridium acidurici</i> , <i>Clostridium aerotolerans</i> ,				
<i>Clostridium aestuarii</i> , <i>Clostridium akagii</i> , <i>Clostridium aldenense</i> , <i>Clostridium aldrichii</i> , <i>Clostridium</i>				
<i>algidicarni</i> , <i>Clostridium algidixylanolyticum</i> , <i>Clostridium algifaecis</i> , <i>Clostridium algoriphilum</i> , <i>Clostridium</i>				
<i>alkalicellulosi</i> , <i>Clostridium aminophilum</i> , <i>Clostridium aminovalericum</i> , <i>Clostridium amygdalinum</i> ,				
<i>Clostridium amylolyticum</i> , <i>Clostridium arbuti</i> , <i>Clostridium arcticum</i> , <i>Clostridium argentinense</i> ,				
<i>Clostridium asparagiforme</i> , <i>Clostridium aurantibutyricum</i> , <i>Clostridium autoethanogenum</i> , <i>Clostridium</i>				
<i>baratii</i> , <i>Clostridium barkeri</i> , <i>Clostridium bartlettii</i> , <i>Clostridium bejerinckii</i> , <i>Clostridium bifementans</i> ,				
<i>Clostridium bolteaie</i> , <i>Clostridium bornimense</i> , <i>Clostridium botulinum</i> , <i>Clostridium bowmanii</i> , <i>Clostridium</i>				
<i>bryantii</i> , <i>Clostridium butyricum</i> , <i>Clostridium cadaveris</i> , <i>Clostridium caenicola</i> , <i>Clostridium</i>				
<i>caminithermale</i> , <i>Clostridium carboxidivorans</i> , <i>Clostridium carnis</i> , <i>Clostridium cavendishii</i> , <i>Clostridium</i>				
<i>celatum</i> , <i>Clostridium celerecrescens</i> , <i>Clostridium cellobioparum</i> , <i>Clostridium cellulofementans</i> ,				
<i>Clostridium cellulolyticum</i> , <i>Clostridium cellulosi</i> , <i>Clostridium cellulovorans</i> , <i>Clostridium chartatabidum</i> ,				
<i>Clostridium chauvoei</i> , <i>Clostridium chromiireducens</i> , <i>Clostridium citroniae</i> , <i>Clostridium clariflavum</i> ,				
<i>Clostridium clostridioforme</i> , <i>Clostridium coccoides</i> , <i>Clostridium cochlearium</i> , <i>Clostridium colletant</i> ,				
<i>Clostridium colicanis</i> , <i>Clostridium colinum</i> , <i>Clostridium collagenovorans</i> , <i>Clostridium cylindrosporum</i> ,				
<i>Clostridium difficile</i> , <i>Clostridium diolis</i> , <i>Clostridium disporicum</i> , <i>Clostridium drakei</i> , <i>Clostridium durum</i> ,				
<i>Clostridium estertheticum</i> , <i>Clostridium estertheticum estertheticum</i> , <i>Clostridium estertheticum</i>				
<i>laramiense</i> , <i>Clostridium fallax</i> , <i>Clostridium felsineum</i> , <i>Clostridium fervidum</i> , <i>Clostridium fimetarium</i> ,				
<i>Clostridium formicaceticum</i> , <i>Clostridium frigidicarnis</i> , <i>Clostridium frigoris</i> , <i>Clostridium ganghwense</i> ,				
<i>Clostridium gasigenes</i> , <i>Clostridium ghonii</i> , <i>Clostridium glycolicum</i> , <i>Clostridium glycyrrhizinilyticum</i> ,				
<i>Clostridium grantii</i> , <i>Clostridium haemolyticum</i> , <i>Clostridium halophilum</i> , <i>Clostridium hastiforme</i> ,				
<i>Clostridium hathewayi</i> , <i>Clostridium herbivorans</i> , <i>Clostridium hiranonis</i> , <i>Clostridium histolyticum</i> ,				
<i>Clostridium homopropionicum</i> , <i>Clostridium huakuii</i> , <i>Clostridium hungatei</i> , <i>Clostridium hydrogeniformans</i> ,				

TABLE 1-continued

EXAMPLE BACTERIA

Clostridium hydroxybenzoicum, *Clostridium hylemonae*, *Clostridium jejuense*, *Clostridium indolis*, *Clostridium innocuum*, *Clostridium intestinale*, *Clostridium irregulare*, *Clostridium isatidis*, *Clostridium josui*, *Clostridium kluyveri*, *Clostridium lactatifermentans*, *Clostridium lacusfryxellense*, *Clostridium laramiense*, *Clostridium lavalense*, *Clostridium lentocellum*, *Clostridium lentoputrescens*, *Clostridium leptum*, *Clostridium limosum*, *Clostridium litorale*, *Clostridium lituseburense*, *Clostridium ljungdahlii*, *Clostridium lortetii*, *Clostridium lundense*, *Clostridium magnum*, *Clostridium malenominatum*, *Clostridium manganotii*, *Clostridium mayombeii*, *Clostridium methoxybenzovorans*, *Clostridium methylpentosum*, *Clostridium neopropionicum*, *Clostridium nexile*, *Clostridium nitrophenolicum*, *Clostridium novyi*, *Clostridium oceanicum*, *Clostridium orbiscindens*, *Clostridium oroticum*, *Clostridium oxalicum*, *Clostridium papyrosolvans*, *Clostridium paradoxum*, *Clostridium parapaperfringens* (Alias: *C. welchii*), *Clostridium paraputrificum*, *Clostridium pascui*, *Clostridium pasteurianum*, *Clostridium peptidivorans*, *Clostridium perenne*, *Clostridium perfringens*, *Clostridium pfennigii*, *Clostridium phytofermentans*, *Clostridium pūliforme*, *Clostridium polysaccharolyticum*, *Clostridium populeti*, *Clostridium propionicum*, *Clostridium proteoclasticum*, *Clostridium proteolyticum*, *Clostridium psychrophilum*, *Clostridium puniceum*, *Clostridium purinilyticum*, *Clostridium putrefaciens*, *Clostridium putrificum*, *Clostridium quercicolum*, *Clostridium quini*, *Clostridium ramosum*, *Clostridium rectum*, *Clostridium roseum*, *Clostridium saccharobutylicum*, *Clostridium saccharogumia*, *Clostridium saccharolyticum*, *Clostridium saccharoperbutylacetonicum*, *Clostridium sardiniense*, *Clostridium sartagoforme*, *Clostridium scatologenes*, *Clostridium schirmacherense*, *Clostridium scindens*, *Clostridium septicum*, *Clostridium sordellii*, *Clostridium sphenoides*, *Clostridium spiroforme*, *Clostridium sporogenes*, *Clostridium sporosphaeroides*, *Clostridium stercorarium*, *Clostridium stercorarium leptospartum*, *Clostridium stercorarium stercorarium*, *Clostridium stercorarium thermolacticum*, *Clostridium sticklandii*, *Clostridium straminisolvans*, *Clostridium subterminale*, *Clostridium sufflavum*, *Clostridium sulfidigenes*, *Clostridium symbiosum*, *Clostridium tagluense*, *Clostridium tepidiprofundii*, *Clostridium termitidis*, *Clostridium tertium*, *Clostridium tetani*, *Clostridium tetanomorphum*, *Clostridium thermacetium*, *Clostridium thermautotrophicum*, *Clostridium thermoalcaliphilum*, *Clostridium thermobutyricum*, *Clostridium thermocellum*, *Clostridium thermocopriae*, *Clostridium thermohydrosulfuricum*, *Clostridium thermolacticum*, *Clostridium thermopalmarium*, *Clostridium thermopapyrolyticum*, *Clostridium thermosaccharolyticum*, *Clostridium thermosuccinogenes*, *Clostridium thermosulfurigenes*, *Clostridium thiosulfatireducens*, *Clostridium tyrobutyricum*, *Clostridium uliginosum*, *Clostridium ultunense*, *Clostridium villosum*, *Clostridium vincentii*, *Clostridium viride*, *Clostridium xylanolyticum*, *Clostridium xylanovorans*

<i>Dactylosporangium</i>	<i>Deinococcus</i>	<i>Delftia</i>	<i>Echinicola</i>
<i>Dactylosporangium aurantiacum</i>	<i>Deinococcus aerius</i>	<i>Delftia acidovorans</i>	<i>Echinicola pacifica</i>
<i>Dactylosporangium fulvum</i>	<i>Deinococcus apachensis</i>	<i>Desulfovibrio desulfuricans</i>	<i>Echinicola vietnamensis</i>
<i>Dactylosporangium matsuzakiense</i>	<i>Deinococcus aquaticus</i>	<i>Diplococcus</i>	
<i>Dactylosporangium roseum</i>	<i>Deinococcus aquatilis</i>	<i>Diplococcus pneumoniae</i>	
<i>Dactylosporangium thailandense</i>	<i>Deinococcus caeni</i>		
<i>Dactylosporangium vinaceum</i>	<i>Deinococcus radiodurans</i>		
	<i>Deinococcus radiophilus</i>		
<i>Enterobacter</i>	<i>Enterobacter kobei</i>	<i>Faecalibacterium</i>	<i>Flavobacterium</i>
<i>E. aerogenes</i>	<i>E. ludwigii</i>	<i>Faecalibacterium prausnitzii</i>	<i>Flavobacterium antarcticum</i>
<i>E. amnigenus</i>	<i>E. mori</i>	<i>Fangia</i>	<i>Flavobacterium aquatile</i>
<i>E. agglomerans</i>	<i>E. nimipressuralis</i>	<i>Fangia hongkongensis</i>	<i>Flavobacterium aquidunense</i>
<i>E. arachidis</i>	<i>E. oryzae</i>	<i>Fastidiosipila</i>	<i>Flavobacterium balustinum</i>
<i>E. asburiae</i>	<i>E. pulveris</i>	<i>Fastidiosipila sanguinis</i>	<i>Flavobacterium croceum</i>
<i>E. cancerogenus</i>	<i>E. pyrinus</i>	<i>Fusobacterium</i>	<i>Flavobacterium cucumis</i>
<i>E. cloacae</i>	<i>E. radicincitans</i>	<i>Fusobacterium nucleatum</i>	<i>Flavobacterium daejeonense</i>
<i>E. cowanii</i>	<i>E. taylorae</i>		<i>Flavobacterium defluvii</i>
<i>E. dissolvans</i>	<i>E. turicensis</i>		<i>Flavobacterium degerlachei</i>
<i>E. gergoviae</i>	<i>E. sakazakii</i>		<i>Flavobacterium denitrificans</i>
<i>E. helveticus</i>	<i>Enterobacter soli</i>		<i>Flavobacterium filum</i>
<i>E. hormaechei</i>	<i>Enterococcus</i>		<i>Flavobacterium flevense</i>
<i>E. intermedius</i>	<i>Enterococcus durans</i>		<i>Flavobacterium frigidarium</i>
	<i>Enterococcus faecalis</i>		<i>Flavobacterium mizutaii</i>
	<i>Enterococcus faecium</i>		
	<i>Erwinia</i>		
	<i>Erwinia hapontici</i>		
	<i>Escherichia</i>		
	<i>Escherichia coli</i>		

TABLE 1-continued

EXAMPLE BACTERIA				<i>Flavobacterium okeanoikoites</i>
Gaetbulibacter	Haemophilus	Ideonella	Janibacter	
<i>Gaetbulibacter saemankumensis</i>	<i>Haemophilus aegyptius</i>	<i>Ideonella azotifigens</i>	<i>Janibacter anophelis</i>	
Gallibacterium	<i>Haemophilus aphrophilus</i>	Idiomarina	<i>Janibacter corallicola</i>	
<i>Gallibacterium anatis</i>	<i>Haemophilus felis</i>	<i>Idiomarina abyssalis</i>	<i>Janibacter limosus</i>	
Gallicola	<i>Haemophilus gallinarum</i>	<i>Idiomarina baltica</i>	<i>Janibacter melonis</i>	
<i>Gallicola barnesae</i>	<i>Haemophilus haemolyticus</i>	<i>Idiomarina fontislapidosi</i>	<i>Janibacter terrae</i>	
Garciella	<i>Haemophilus haemolyticus</i>	<i>Idiomarina loihiensis</i>	Jannaschia	
<i>Garciella nitratireducens</i>	<i>Haemophilus paracuniculus</i>	<i>Idiomarina ramblicola</i>	<i>Jannaschia cystaugens</i>	
Geobacillus	<i>Haemophilus parahaemolyticus</i>	<i>Idiomarina seostinensis</i>	<i>Jannaschia helgolandensis</i>	
<i>Geobacillus thermoglucosidasius</i>	<i>Haemophilus parainfluenzae</i>	<i>Idiomarina zobellii</i>	<i>Jannaschia pohangensis</i>	
<i>Geobacillus stearothermophilus</i>	<i>Haemophilus parasuis</i>	Ignatzschineria	<i>Jannaschia rubra</i>	
Geobacter	<i>Haemophilus hydrogenophilus</i>	<i>Ignatzschineria larvae</i>	Janthinobacterium	
<i>Geobacter bemidjensis</i>	<i>Haemophilus pittmaniae</i>	<i>Ignatzschineria larvae</i>	<i>Janthinobacterium agaricidammosum</i>	
<i>Geobacter brememis</i>	Hafnia	Ignavigranum	<i>Janthinobacterium alimentarius</i>	
<i>Geobacter chapellei</i>	<i>Hafnia alvei</i>	<i>Ignavigranum ruoffiae</i>	<i>Jeotgaliococcus halotolerans</i>	
<i>Geobacter grbiciae</i>	Hahella	Illumatobacter		
<i>Geobacter hydrogenophilus</i>	<i>Hahella ganghwensis</i>	<i>Illumatobacter fluminis</i>		
<i>Geobacter lovleyi</i>	<i>Hahella halalkalibacillus</i>	Ilyobacter		
<i>Geobacter metallireducens</i>	<i>Hahella halalkalibacillus</i>	<i>Ilyobacter delafieldii</i>		
<i>Geobacter pelophilus</i>	<i>Halalkalibacillus halophilus</i>	<i>Ilyobacter insuetus</i>		
<i>Geobacter pickeringii</i>	Helicobacter	<i>Ilyobacter polytropus</i>		
<i>Geobacter sulfurireducens</i>	<i>Helicobacter pylori</i>	<i>Ilyobacter tartaricus</i>		
Geodermatophilus				
<i>Geodermatophilus obscurus</i>				
Gluconacetobacter				
<i>Gluconacetobacter xylinus</i>				
Gordonia				
<i>Gordonia rubripertincta</i>				
Kaistia	Labedella	<i>Listeria ivanovii</i>	Micrococcus	Nesterenkonia
<i>Kaistia adipata</i>	<i>Labedella gwakjiensis</i>	<i>L. marthii</i>	<i>Micrococcus luteus</i>	<i>Nesterenkonia holobia</i>
<i>Kaistia soli</i>	Labrenzia	<i>L. monocytogenes</i>	<i>Micrococcus lylae</i>	Nocardia
Kangiella	<i>Labrenzia aggregata</i>	<i>L. newyorkensis</i>	Moraxella	<i>Nocardia argentinensis</i>
<i>Kangiella aquimarina</i>	<i>Labrenzia alba</i>	<i>L. riparia</i>	<i>Moraxella bovis</i>	<i>Nocardia corallina</i>
<i>Kangiella koreensis</i>	Labrenzia	<i>L. rocourtiae</i>	<i>Moraxella nonliquefaciens</i>	<i>Nocardia otitidiscaviarum</i>
Kerstersia	<i>Labrenzia alexandrii</i>	<i>L. seeligeri</i>	Moraxella	
<i>Kerstersia gyiorum</i>	<i>Labrenzia marina</i>	<i>L. weihenstephanensis</i>	<i>Moraxella osloensis</i>	
Kiloniella	Labrys	<i>L. welshimeri</i>	Nakamurella	
<i>Kiloniella laminariae</i>	<i>Labrys methylaminiphilus</i>	Listonella	<i>Nakamurella multipartita</i>	
Klebsiella	<i>Labrys miyagiensis</i>	<i>Listonella anguillarum</i>	Nannocystis	
<i>K. granulomatis</i>	<i>Labrys monachus</i>	Macrococcus	<i>Nannocystis pusilla</i>	
<i>K. oxytoca</i>	<i>Labrys okinawensis</i>	<i>Macrococcus bovicus</i>	Natranaerobius	
<i>K. pneumoniae</i>	Labrys	Marinobacter	<i>Natranaerobius thermophilus</i>	
<i>K. terrigena</i>	<i>Labrys portucalensis</i>	<i>Marinobacter flavimaris</i>	<i>Natranaerobius trueperi</i>	
<i>K. variicola</i>	Lactobacillus	<i>Marinobacter ammontaphilum</i>	Naxibacter	
Kluyvera	[see below]	Meiothermus	<i>Naxibacter alkalitolerans</i>	
<i>Kluyvera ascorbata</i>	Laceyella	<i>Meiothermus ruber</i>	Neisseria	
Kocuria	<i>Laceyella putida</i>	Methylophilus	<i>Neisseria cinerea</i>	
<i>Kocuria roasea</i>	Lechevalieria	<i>Methylophilus methylotrophus</i>	<i>Neisseria denitrificans</i>	
<i>Kocuria varians</i>	<i>Lechevalieria aerocolonigenes</i>	Legionella	<i>Neisseria gonorrhoeae</i>	
Kurthia	Legionella	[see below]	<i>Neisseria lactamica</i>	
<i>Kurthia zopfii</i>	[see below]	Listeria	<i>Neisseria mucosa</i>	
	<i>L. aquatica</i>	<i>Listeria ammontaphilum</i>	<i>Neisseria sica</i>	
	<i>L. booriae</i>	<i>Listeria cornellensis</i>		
	<i>L. floridensis</i>	<i>Listeria fleischmannii</i>		
		<i>Listeria liquefaciens</i>		

TABLE 1-continued

EXAMPLE BACTERIA				
	<i>L. grandensis</i>	<i>Microbacterium</i>	<i>Neisseria subflava</i>	
	<i>L. grayi</i>	<i>oxydans</i>	<i>Neptunomonas</i>	
	<i>L. imocua</i>		<i>Neptunomonas japonica</i>	
<hr/>				
Lactobacillus				
<i>L. acetotolerans</i>	<i>L. cateniformis</i>	<i>L. mali</i>	<i>L. parakefiri</i>	<i>L. sakei</i>
<i>L. acidifarinae</i>	<i>L. cetii</i>	<i>L. manihotivorans</i>	<i>L. paralimentarius</i>	<i>L. salivarius</i>
<i>L. acidipiscis</i>	<i>L. coleohominis</i>	<i>L. mindensis</i>	<i>L. paraplantarum</i>	<i>L. sanfranciscensis</i>
<i>L. acidophilus</i>	<i>L. collinoides</i>	<i>L. mucosae</i>	<i>L. pentosus</i>	<i>L. satsumensis</i>
<i>Lactobacillus agilis</i>	<i>L. composti</i>	<i>L. murinus</i>	<i>L. perolens</i>	<i>L. secaliphilus</i>
<i>L. algidus</i>	<i>L. concavus</i>	<i>L. nagelii</i>	<i>L. plantarum</i>	<i>L. sharpeae</i>
<i>L. alimentarius</i>	<i>L. coryniformis</i>	<i>L. namuwensis</i>	<i>L. pontis</i>	<i>L. siliginis</i>
<i>L. amylolyticus</i>	<i>L. crispatus</i>	<i>L. nantensis</i>	<i>L. protectus</i>	<i>L. spicheri</i>
<i>L. amylophilus</i>	<i>L. crustorum</i>	<i>L. oligofermentans</i>	<i>L. psittaci</i>	<i>L. subcicus</i>
<i>L. amylotrophicus</i>	<i>L. curvatus</i>	<i>L. oris</i>	<i>L. reminii</i>	<i>L. thailandensis</i>
<i>L. amylovorus</i>	<i>L. delbrueckii</i>	<i>L. panis</i>	<i>L. reuteri</i>	<i>L. ulunensis</i>
<i>L. animalis</i>	subsp. <i>bulgaricus</i>	<i>L. pantheris</i>	<i>L. rhammosus</i>	<i>L. vaccinostrercus</i>
<i>L. antri</i>	<i>L. delbrueckii</i>	<i>L. parabrevis</i>	<i>L. rimae</i>	<i>L. vaginalis</i>
<i>L. apodemi</i>	subsp. <i>delbrueckii</i>	<i>L. parabuchneri</i>	<i>L. rogosae</i>	<i>L. versmoldensis</i>
<i>L. avitarius</i>	<i>L. delbrueckii</i>	<i>L. paracasei</i>	<i>L. rossiae</i>	<i>L. vini</i>
<i>L. bifementans</i>	subsp. <i>lactis</i>	<i>L. paracollinoides</i>	<i>L. ruminis</i>	<i>L. vitulinus</i>
<i>L. brevis</i>	<i>L. dextrinicus</i>	<i>L. parafarraginis</i>	<i>L. saerimmeri</i>	<i>L. zaeae</i>
<i>L. buchneri</i>	<i>L. diolivorans</i>	<i>L. homohiochii</i>	<i>L. jensenii</i>	<i>L. zymae</i>
<i>L. camelliae</i>	<i>L. equi</i>	<i>L. iners</i>	<i>L. johnsonii</i>	<i>L. gastricus</i>
<i>L. casei</i>	<i>L. equigenerosi</i>	<i>L. ingluviei</i>	<i>L. kalixensis</i>	<i>L. ghanensis</i>
<i>L. kitasatonis</i>	<i>L. farraginis</i>	<i>L. intestinalis</i>	<i>L. kefranofermentans</i>	<i>L. graminis</i>
<i>L. kunkeei</i>	<i>L. farciminis</i>	<i>L. fuchuensis</i>	<i>L. kefiri</i>	<i>L. hammersii</i>
<i>L. leichmannii</i>	<i>L. fermentum</i>	<i>L. gallinarum</i>	<i>L. kimchii</i>	<i>L. hamsteri</i>
<i>L. lindneri</i>	<i>L. formicis</i>	<i>L. gasseri</i>	<i>L. helveticus</i>	<i>L. harbinensis</i>
<i>L. malefermentans</i>	<i>L. fructivorans</i>		<i>L. hilgardii</i>	<i>L. hayakitensis</i>
	<i>L. frumenti</i>			
<hr/>				
Legionella				
<i>Legionella adelaiddensis</i>	<i>Legionella drancourtii</i>	<i>Candidatus Legionella jeonii</i>	<i>Legionella quinlivanii</i>	
<i>Legionella anisa</i>	<i>Legionella dresdenensis</i>	<i>Legionella jordanis</i>	<i>Legionella rowbothamii</i>	
<i>Legionella belliardensis</i>	<i>Legionella drozanskii</i>	<i>Legionella lansingensis</i>	<i>Legionella rubrilucens</i>	
<i>Legionella birminghamensis</i>	<i>Legionella dumoffii</i>	<i>Legionella londiniensis</i>	<i>Legionella sainthelensi</i>	
<i>Legionella bozemanae</i>	<i>Legionella erythra</i>	<i>Legionella longbeachae</i>	<i>Legionella santicrucis</i>	
<i>Legionella brunensis</i>	<i>Legionella faiyfeldensis</i>	<i>Legionella lytica</i>	<i>Legionella shakespearei</i>	
<i>Legionella busanensis</i>	<i>Legionella fallonii</i>	<i>Legionella maceachernii</i>	<i>Legionella spiritensis</i>	
<i>Legionella cardiaca</i>	<i>Legionella feeleii</i>	<i>Legionella massiliensis</i>	<i>Legionella steelei</i>	
<i>Legionella cherrii</i>	<i>Legionella geestiana</i>	<i>Legionella micdadei</i>	<i>Legionella steigerwaltii</i>	
<i>Legionella cincinnatiensis</i>	<i>Legionella genomospecies</i>	<i>Legionella monrovia</i>	<i>Legionella taurenensis</i>	
<i>Legionella clemsonensis</i>	<i>Legionella gormanii</i>	<i>Legionella moravica</i>	<i>Legionella tucsonensis</i>	
<i>Legionella donaldsonii</i>	<i>Legionella gratiana</i>	<i>Legionella nagasakiensis</i>	<i>Legionella tunisiensis</i>	
	<i>Legionella gresilensis</i>	<i>Legionella nautarum</i>	<i>Legionella wadsworthii</i>	
	<i>Legionella hackeliae</i>	<i>Legionella norrlandica</i>	<i>Legionella waltersii</i>	
	<i>Legionella impletisoli</i>	<i>Legionella oakridgensis</i>	<i>Legionella worsteiensis</i>	
	<i>Legionella israelensis</i>	<i>Legionella parisiensis</i>	<i>Legionella yabuuchiae</i>	
	<i>Legionella jamestowniensis</i>	<i>Legionella pittsburghensis</i>		
		<i>Legionella pneumophila</i>		
		<i>Legionella quateirensis</i>		
<hr/>				
Oceanibulbus	Paenibacillus	Prevotella	Quadriflora	
<i>Oceanibulbus indolifex</i>	<i>Paenibacillus thiaminolyticus</i>	<i>Prevotella albensis</i>	<i>Quadriflora granulorum</i>	
<i>Oceanicaulis</i>	<i>Pantoea</i>	<i>Prevotella amnii</i>	<i>Quadriflora cococcus</i>	
<i>Oceanicaulis alexandrii</i>	<i>Pantoea agglomerans</i>	<i>Prevotella bergensis</i>	<i>Quadriflora australiensis</i>	

TABLE 1-continued

EXAMPLE BACTERIA				
<i>Oceanicola</i>	<i>Paracoccus</i>	<i>Prevotella bivia</i>	<i>Quinella</i>	
<i>Oceanicola batsensis</i>	<i>Paracoccus</i>	<i>Prevotella brevis</i>	<i>Quinella ovalis</i>	
<i>Oceanicola</i>	<i>alcaliphilus</i>	<i>Prevotella</i>	<i>Ralstonia</i>	
<i>granulosus</i>	<i>Paucimonas</i>	<i>bryantii</i>	<i>Ralstonia</i>	
<i>Oceanicola</i>	<i>Paucimonas</i>	<i>Prevotella buccae</i>	<i>eutropha</i>	
<i>nanhaiensis</i>	<i>lemoinei</i>	<i>Prevotella</i>	<i>Ralstonia</i>	
<i>Oceanimonas</i>	<i>Pectobacterium</i>	<i>buccalis</i>	<i>insidiosa</i>	
<i>Oceanimonas</i>	<i>Pectobacterium</i>	<i>Prevotella copri</i>	<i>Ralstonia</i>	
<i>baumannii</i>	<i>avoidearum</i>	<i>Prevotella</i>	<i>mannitolilytica</i>	
<i>Oceaniserpentilla</i>	<i>Pectobacterium</i>	<i>dentalis</i>	<i>Ralstonia</i>	
<i>Oceaniserpentilla</i>	<i>atrosepticum</i>	<i>Prevotella</i>	<i>pickettii</i>	
<i>haliotis</i>	<i>Pectobacterium</i>	<i>denticola</i>	<i>Ralstonia</i>	
<i>Oceanisphaera</i>	<i>betavascularum</i>	<i>Prevotella disiens</i>	<i>pseudosolanacearum</i>	
<i>Oceanisphaera</i>	<i>Pectobacterium</i>	<i>Prevotella</i>	<i>Ralstonia syzygii</i>	
<i>donghaensis</i>	<i>cacticida</i>	<i>histicola</i>	<i>Ralstonia</i>	
<i>Oceanisphaera</i>	<i>Pectobacterium</i>	<i>Prevotella</i>	<i>solanacearum</i>	
<i>litoralis</i>	<i>carnegieana</i>	<i>intermedia</i>	<i>Ramlibacter</i>	
<i>Oceanithermus</i>	<i>Pectobacterium</i>	<i>Prevotella</i>	<i>Ramlibacter</i>	
<i>Oceanithermus</i>	<i>carotovorum</i>	<i>maculosa</i>	<i>henchirensis</i>	
<i>desulfurans</i>	<i>Pectobacterium</i>	<i>Prevotella</i>	<i>Ramlibacter</i>	
<i>Oceanithermus</i>	<i>chrysanthemii</i>	<i>marshii</i>	<i>tataouinensis</i>	
<i>profundus</i>	<i>Pectobacterium</i>	<i>Prevotella</i>	<i>Raoultella</i>	
<i>Oceanobacillus</i>	<i>cyripedii</i>	<i>melaninogenica</i>	<i>Raoultella</i>	
<i>Oceanobacillus caeni</i>	<i>Pectobacterium</i>	<i>Prevotella micans</i>	<i>ornithinolytica</i>	
<i>Oceanospirillum</i>	<i>rhapontici</i>	<i>Prevotella</i>	<i>Raoultella</i>	
<i>Oceanospirillum</i>	<i>Pectobacterium</i>	<i>multiformis</i>	<i>planticola</i>	
<i>linum</i>	<i>wasabiae</i>	<i>Prevotella</i>	<i>Raoultella</i>	
	<i>Planococcus</i>	<i>nigrescens</i>	<i>terrigena</i>	
	<i>Planococcus</i>	<i>Prevotella oralis</i>	<i>Rathayibacter</i>	
	<i>citreus</i>	<i>Prevotella oris</i>	<i>Rathayibacter</i>	
	<i>Planomicrobium</i>	<i>Prevotella</i>	<i>caricis</i>	
	<i>Planomicrobium</i>	<i>oulorum</i>	<i>Rathayibacter</i>	
	<i>okeanokoites</i>	<i>Prevotella pallens</i>	<i>festucae</i>	
	<i>Plesiomonas</i>	<i>Prevotella salivae</i>	<i>Rathayibacter</i>	
	<i>Plesiomonas</i>	<i>Prevotella</i>	<i>iranicus</i>	
	<i>shigelloides</i>	<i>stercorea</i>	<i>Rathayibacter</i>	
	<i>Proteus</i>	<i>Prevotella</i>	<i>rathayi</i>	
	<i>Proteus vulgaris</i>	<i>tannerae</i>	<i>Rathayibacter</i>	
		<i>Prevotella</i>	<i>toxicus</i>	
		<i>timonensis</i>	<i>Rathayibacter</i>	
		<i>Prevotella</i>	<i>tritici</i>	
		<i>veroralis</i>	<i>Rhodobacter</i>	
		<i>Providencia</i>	<i>Rhodobacter</i>	
		<i>Providencia</i>	<i>sphaeroides</i>	
		<i>stuartii</i>	<i>Ruegeria</i>	
		<i>Pseudomonas</i>	<i>Ruegeria</i>	
		<i>Pseudomonas</i>	<i>gelatinovorans</i>	
		<i>aeruginosa</i>		
		<i>Pseudomonas</i>		
		<i>alcaligenes</i>		
		<i>Pseudomonas</i>		
		<i>anguillispetica</i>		
		<i>Pseudomonas</i>		
		<i>fluorescens</i>		
		<i>Pseudoalteromonas</i>		
		<i>haloplanktis</i>		
		<i>Pseudomonas</i>		
		<i>mendocina</i>		
		<i>Pseudomonas</i>		
		<i>pseudoalcaligenes</i>		
		<i>Pseudomonas</i>		
		<i>putida</i>		
		<i>Pseudomonas</i>		
		<i>tutzeri</i>		
		<i>Pseudomonas</i>		
		<i>syringae</i>		
		<i>Psychrobacter</i>		
		<i>Psychrobacter</i>		
		<i>faecalis</i>		
		<i>Psychrobacter</i>		
		<i>phenylpyruvicus</i>		
<i>Saccharococcus</i>	<i>Sagittula</i>	<i>Sanguibacter</i>	<i>Stenotrophomonas</i>	<i>Tatlockia</i>
<i>Saccharococcus</i>	<i>Sagittula stellata</i>	<i>Sanguibacter</i>	<i>Stenotrophomonas</i>	<i>Tatlockia</i>
<i>thermophilus</i>	<i>Salegentibacter</i>	<i>keddiei</i>	<i>maltophila</i>	<i>maceachernii</i>
<i>Saccharomonospora</i>	<i>Salegentibacter</i>	<i>Sanguibacter</i>	<i>Streptococcus</i>	<i>Tatlockia</i>
<i>Saccharomonospora</i>	<i>salegens</i>	<i>suarezii</i>	[also see below]	<i>micdadei</i>
<i>azurea</i>	<i>Salimicrobium</i>	<i>Saprosira</i>	<i>Streptomyces</i>	<i>Tenacibaculum</i>

TABLE 1-continued

EXAMPLE BACTERIA				
<i>Saccharomonospora cyanea</i>	<i>Salimicrobium album</i>	<i>Saprospira grandis</i>	<i>Streptomyces achromogenes</i>	<i>Tenacibaculum amylolyticum</i>
<i>Saccharomonospora viridis</i>	<i>Salinibacter ruber</i>	<i>Sarcina maxima</i>	<i>Streptomyces cesalbus</i>	<i>Tenacibaculum discolor</i>
<i>Saccharophagus degradans</i>	<i>Salinicoccus alkaliphilus</i>	<i>Sarcina ventriculi</i>	<i>Streptomyces cescaepitosus</i>	<i>Tenacibaculum gallaicum</i>
<i>Saccharopolyspora erythraea</i>	<i>Salinicoccus hispanicus</i>	<i>Serratia</i>	<i>Streptomyces cesdiastaticus</i>	<i>Tenacibaculum lutimaris</i>
<i>Saccharopolyspora gregorii</i>	<i>Salinicoccus roseus</i>	<i>Serratia fonticola</i>	<i>Streptomyces cesexfoliatus</i>	<i>Tenacibaculum mesophilum</i>
<i>Saccharopolyspora hirsuta</i>	<i>Salinispora arenicola</i>	<i>Serratia marcescens</i>	<i>Streptomyces fimbriatus</i>	<i>Tenacibaculum skagerrakense</i>
<i>Saccharopolyspora hordei</i>	<i>Salinispora tropica</i>	<i>Sphaerotilus natans</i>	<i>Streptomyces fulvisimus</i>	<i>Tepidanaerobacter tepidanaerobacter</i>
<i>Saccharopolyspora rectivirgula</i>	<i>Salinivibrio costicola</i>	<i>Sphingobacterium multivorum</i>	<i>Streptomyces fradiae</i>	<i>Tepidanaerobacter syntrophicus</i>
<i>Saccharopolyspora spinosa</i>	<i>Salmonella bongori</i>	<i>Staphylococcus</i>	<i>Streptomyces griseus</i>	<i>Tepidibacter tepidibacter</i>
<i>Saccharopolyspora taberi</i>	<i>Salmonella enterica</i>	[see below]	<i>Streptomyces griseoruber</i>	<i>formicigenes</i>
<i>Saccharothrix australiensis</i>	<i>Salmonella subterranea</i>		<i>Streptomyces griseus</i>	<i>Tepidibacter thalassicus</i>
<i>Saccharothrix coeruleofusca</i>	<i>Salmonella typhi</i>		<i>Streptomyces lavendulae</i>	<i>Thermus thermophilus</i>
<i>Saccharothrix espanaensis</i>			<i>Streptomyces aquaticus</i>	
<i>Saccharothrix longispora</i>			<i>Streptomyces phaeochromogenes</i>	
<i>Saccharothrix mutabilis</i>			<i>Streptomyces thermodiastaticus</i>	
<i>Saccharothrix syringae</i>			<i>Streptomyces tubercidicus</i>	
<i>Saccharothrix tangerinus</i>				
<i>Saccharothrix texasensis</i>				
<i>Staphylococcus S. arlettae</i>	<i>S. equorum</i>	<i>S. microti</i>		<i>S. schleiferi</i>
<i>S. agnetis</i>	<i>S. felis</i>	<i>S. muscae</i>		<i>S. sciuri</i>
<i>S. aureus</i>	<i>S. fleurettii</i>	<i>S. nepalensis</i>		<i>S. simiae</i>
<i>S. auricularis</i>	<i>S. gallinarum</i>	<i>S. pasteurii</i>		<i>S. simulans</i>
<i>S. capitis</i>	<i>S. haemolyticus</i>	<i>S. petrasii</i>		<i>S. stepanovicii</i>
<i>S. caprae</i>	<i>S. hominis</i>	<i>S. pettenkoferi</i>		<i>S. succinus</i>
<i>S. carnosus</i>	<i>S. hyicus</i>	<i>S. piscifermentans</i>		<i>S. vitulinus</i>
<i>S. caseolyticus</i>	<i>S. intermedius</i>	<i>S. pseudintermedius</i>		<i>S. warneri</i>
<i>S. chromogenes</i>	<i>S. kloosii</i>	<i>S. pseudolugdunensis</i>		<i>S. xylosus</i>
<i>S. cohnii</i>	<i>S. leei</i>	<i>S. pulvereri</i>		
<i>S. condimenti</i>	<i>S. lentus</i>	<i>S. rostri</i>		
<i>S. delphini</i>	<i>S. lugdunensis</i>	<i>S. saccharolyticus</i>		
<i>S. devriesei</i>	<i>S. lutrae</i>	<i>S. saprophyticus</i>		
<i>S. epidermidis</i>	<i>S. lyticans</i>			
	<i>S. massiliensis</i>			
<i>Streptococcus Streptococcus agalactiae</i>	<i>Streptococcus infantarius</i>	<i>Streptococcus orisratti</i>	<i>Streptococcus thermophilus</i>	
<i>Streptococcus anginosus</i>	<i>Streptococcus iniae</i>	<i>Streptococcus parasanguinis</i>	<i>Streptococcus sanguinis</i>	
<i>Streptococcus bovis</i>	<i>Streptococcus intermedius</i>	<i>Streptococcus peroris</i>	<i>Streptococcus sobrinus</i>	
<i>Streptococcus canis</i>	<i>Streptococcus lactarius</i>	<i>Streptococcus pneumoniae</i>	<i>Streptococcus suis</i>	
<i>Streptococcus constellatus</i>	<i>Streptococcus milleri</i>	<i>Streptococcus pseudopneumoniae</i>	<i>Streptococcus uberis</i>	
<i>Streptococcus downei</i>	<i>Streptococcus mitis</i>	<i>Streptococcus pyogenes</i>	<i>Streptococcus vestibularis</i>	
<i>Streptococcus dysgalactiae</i>	<i>Streptococcus mutans</i>	<i>Streptococcus ratti</i>	<i>Streptococcus viridans</i>	
<i>Streptococcus equines</i>	<i>Streptococcus mutans</i>	<i>Streptococcus salivarius</i>	<i>Streptococcus zooepidemicus</i>	
<i>Streptococcus faecalis</i>	<i>Streptococcus oralis</i>			
<i>Streptococcus ferus</i>	<i>Streptococcus tigurinus</i>			
<i>Uliginosibacterium Uliginosibacterium gangwonense</i>	<i>Vagococcus carniphilus</i>	<i>Vibrio aerogenes</i>	<i>Virgibacillus halodemitrificans</i>	<i>Xanthobacter agilis</i>
<i>Ulvibacter</i>	<i>Vagococcus elongatus</i>	<i>Vibrio aestuarianus</i>	<i>Virgibacillus pantothenicus</i>	<i>Xanthobacter aminoxidans</i>
<i>Ulvibacter litoralis</i>	<i>Vagococcus fessus</i>	<i>Vibrio albensis</i>	<i>Weissella</i>	<i>Xanthobacter autotrophicus</i>
<i>Umezawaea</i>	<i>Vagococcus fluvialis</i>	<i>alginolyticus</i>	<i>Weissella cibaria</i>	
<i>Umezawaea tangerina</i>				

TABLE 1-continued

EXAMPLE BACTERIA				
Undibacterium	<i>Vagococcus lutrae</i>	<i>Vibrio campbellii</i>	<i>Weissella confusa</i>	<i>Xanthobacter flavus</i>
<i>Undibacterium pigrum</i>	<i>Vagococcus salmoninarum</i>	<i>Vibrio cholerae</i>	<i>Weissella halotolerans</i>	<i>Xanthobacter tagetidis</i>
Ureaplasma	Variovorax	<i>Vibrio cincinnatiensis</i>	<i>Weissella hellenica</i>	<i>Xanthobacter viscosus</i>
<i>Ureaplasma urealyticum</i>	Variovorax	<i>Vibrio coralliilyticus</i>	<i>Weissella kandleri</i>	Xanthomonas
Ureibacillus	<i>boronicumulans</i>	<i>Vibrio cyclitrophicus</i>	<i>Weissella korensis</i>	<i>Xanthomonas albilineans</i>
<i>Ureibacillus composti</i>	Variovorax	<i>Vibrio diazotrophicus</i>	<i>Weissella minor</i>	<i>Xanthomonas alfalfae</i>
<i>Ureibacillus dokdonensis</i>	<i>dokdonensis</i>	<i>Vibrio fluvialis</i>	<i>Weissella paramesenteroides</i>	<i>Xanthomonas arboricola</i>
<i>Ureibacillus suwonensis</i>	Variovorax	<i>Vibrio furnissii</i>	<i>Weissella soli</i>	<i>Xanthomonas axonopodis</i>
<i>Ureibacillus terrenus</i>	<i>paradoxus</i>	<i>Vibrio gazogenes</i>	<i>Weissella thailandensis</i>	<i>Xanthomonas campestris</i>
<i>Ureibacillus thermophilus</i>	Variovorax <i>soli</i>	<i>Vibrio haliotocoli</i>	<i>Weissella viridescens</i>	<i>Xanthomonas citri</i>
<i>Ureibacillus thermosphaericus</i>	Veillonella	<i>Vibrio harveyi</i>	Williamsia	<i>Xanthomonas codiaei</i>
	<i>Veillonella atypica</i>	<i>Vibrio ichthyenteri</i>	<i>Williamsia marianensis</i>	<i>Xanthomonas cucurbitae</i>
	<i>Veillonella caviae</i>	<i>Vibrio mediterranei</i>	<i>Williamsia metschnikovii</i>	<i>Xanthomonas euvesicatoria</i>
	<i>Veillonella criceti</i>	<i>Vibrio montpellierensis</i>	<i>Vibrio mytili</i>	<i>Xanthomonas fragariae</i>
	<i>Veillonella dispar</i>	<i>Veillonella parvula</i>	<i>Vibrio natriegens</i>	<i>Xanthomonas fuscans</i>
	<i>Veillonella</i>	<i>Veillonella ratti</i>	<i>Vibrio navarrensis</i>	<i>Xanthomonas gardneri</i>
	<i>Veillonella rodentium</i>	<i>Veillonella</i>	<i>Vibrio nereis</i>	<i>Xanthomonas hortorum</i>
	Venenivibrio	<i>Veillonella</i>	<i>Vibrio nigripulchritudo</i>	<i>Xanthomonas hyacinthi</i>
	Venenivibrio	rodentium	<i>Vibrio ordalii</i>	<i>Xanthomonas perforans</i>
	<i>stagnispumantis</i>	Venenivibrio	<i>Vibrio orientalis</i>	<i>Xanthomonas phaseoli</i>
	Verminephrobacter	<i>stagnispumantis</i>	<i>Vibrio paraahaemolyticus</i>	<i>Xanthomonas pisi</i>
	<i>Verminephrobacter eiseniae</i>	Verminephrobacter	<i>Vibrio pectenicida</i>	<i>Xanthomonas populi</i>
	Verrucomicrobium	<i>Verminephrobacter eiseniae</i>	<i>Vibrio penaeicida</i>	<i>Xanthomonas theicola</i>
	<i>Verrucomicrobium spinosum</i>	Verrucomicrobium	<i>Vibrio proteolyticus</i>	<i>Xanthomonas translucens</i>
		<i>Verrucomicrobium spinosum</i>	<i>Vibrio shilonii</i>	<i>Xanthomonas vesicatoria</i>
			<i>Vibrio splendidus</i>	Xylella
			<i>Vibrio tubiashii</i>	<i>Xylella fastidiosa</i>
			<i>Vibrio vulnificus</i>	Xylophilus
				<i>Xylophilus ampelinus</i>
Xenophilus	Yangia	<i>Yersinia mollaretii</i>	Zooshikella	Zobellella
<i>Xenophilus azovorans</i>	<i>Yangia pacifica</i>	<i>Yersinia philomiragia</i>	<i>Zooshikella ganghwensis</i>	<i>Zobellella denitrificans</i>
Xenorhabdus	Yaniella	<i>Yersinia pseudotuberculosis</i>	Zunongwangia	<i>Zobellella taiwanensis</i>
<i>Xenorhabdus beddingii</i>	<i>Yaniella flava</i>	<i>Yersinia rohdei</i>	<i>Zunongwangia profunda</i>	<i>Zeaxanthinibacter zeaxanthinibacter</i>
<i>Xenorhabdus bovienii</i>	Yaniella	<i>Yersinia ruckeri</i>	Zymbacter	<i>Zeaxanthinibacter enoshimensis</i>
<i>Xenorhabdus cabanillasii</i>	<i>halotolerans</i>	Yokenella	<i>Zymbacter palmae</i>	Zihengliuella
<i>Xenorhabdus doucetiae</i>	Yeosuana	<i>Yokenella regensburgei</i>	Zymomonas	<i>Zihengliuella halotolerans</i>
<i>Xenorhabdus griffinae</i>	<i>Yeosuana aromativorans</i>	Yonghaparkia	<i>Zymomonas mobilis</i>	Xylanibacterium
<i>Xenorhabdus hominickii</i>	Yersinia	alkaliphila	Zymophilus	<i>Xylanibacterium ulmi</i>
<i>Xenorhabdus koppenhoeferi</i>	<i>Yersinia aldovae</i>	Zavarzinia	<i>Zymophilus paucivorans</i>	
<i>Xenorhabdus nematophila</i>	<i>Yersinia bercovieri</i>	<i>Zavarzinia raffinosivorans</i>	<i>Zymophilus</i>	
<i>Xenorhabdus poinarii</i>	Yersinia			
Xylanibacter	<i>enterocolitica</i>			
<i>Xylanibacter oryzae</i>	Yersinia			
	<i>frederiksenii</i>			
	<i>Yersinia intermedia</i>			
	<i>Yersinia kristensenii</i>			

TABLE 2

MEDICAMENTS			
ACE inhibitors with calcium channel blocking agents	decongestants	respiratory agents	otic agents
ACE inhibitors with thiazides	dermatological agents	sex hormones	renin inhibitors
adamantane antivirals	diagnostic	topical agents	respiratory agents
adrenal cortical steroids	radiopharmaceuticals	uncategorized agents	respiratory inhalant products
adrenal corticosteroid inhibitors	diarylquinolines	vaginal agents	rifamycin derivatives
adrenergic bronchodilators	dibenzazepine	mitotic inhibitors	salicylates
agents for hypertensive emergencies	anticonvulsants	monoamine oxidase inhibitors	sclerosing agents
agents for pulmonary hypertension	digestive enzymes	mouth and throat products	second generation cephalosporins
aldosterone receptor antagonists	dipeptidyl peptidase 4 inhibitors	mTOR inhibitors	selective estrogen receptor modulators
alkylating agents	diuretics	mucolytics	selective immunosuppressants
allergens	dopaminergic	multikinase inhibitors	selective phosphodiesterase-4 inhibitors
alpha-glucosidase inhibitors	antiparkinsonism agents	muscle relaxants	selective serotonin reuptake inhibitors
alternative medicines	drugs used in alcohol dependence	mydriatics	serotonin-norepinephrine reuptake inhibitors
amebicides	echinocandins	narcotic analgesic combinations	serotonergic neuroenteric modulators
aminoglycosides	EGFR inhibitors	narcotic analgesics	sex hormone combinations
aminopenicillins	estrogen receptor antagonists	nasal anti-infectives	sex hormones
aminosalicylates	expectorants	nasal antihistamines and decongestants	SGLT-2 inhibitors
AMPA receptor antagonists	factor Xa inhibitors	nasal lubricants and irrigations	skeletal muscle relaxant combinations
amylin analogs	fatty acid derivative	nasal preparations	relaxants
analgesic combinations	anticonvulsants	nasal steroids	smoking cessation agents
analgesics	fibric acid derivatives	natural penicillins	somatostatin and somatostatin analogs
androgens and anabolic steroids	first generation cephalosporins	neprilysin inhibitors	spermicides
angiotensin converting enzyme inhibitors	cephalosporins	neuraminidase inhibitors	statins
angiotensin II inhibitors with calcium channel blockers	fourth generation cephalosporins	neuromuscular blocking agents	sterile irrigating solutions
angiotensin II inhibitors with thiazides	functional bowel disorder agents	neuronal potassium channel openers	streptomyces derivatives
angiotensin receptor blockers	gallstone solubilizing agents	next generation cephalosporins	succinimide anticonvulsants
angiotensin receptor blockers and neprilysin inhibitors	gamma-aminobutyric acid analogs	nicotinic acid derivatives	sulfonamides
anorectal preparations	gamma-aminobutyric acid reuptake inhibitors	NK1 receptor antagonists	sulfonylureas
anorexiant	gastrointestinal agents	NNRTIs	synthetic ovulation stimulants
antacids	general anesthetics	non-cardioselective beta blockers	tetracyclic antidepressants
anthelmintics	genitourinary tract agents	contrast media	tetracyclines
anti-angiogenic ophthalmic agents	GI stimulants	non-ionic iodinated contrast media	therapeutic radiopharmaceuticals
anti-CTLA-4 monoclonal antibodies	glucocorticoids	non-sulfonylureas	therapeutic vaccines
anti-infectives	glucose elevating agents	non-steroidal anti-inflammatory agents	thiazide diuretics
Anti-PD-1 monoclonal antibodies	glycopeptide antibiotics	NS5A inhibitors	thiazolidinediones
antiadrenergic agents (central) with thiazides	glycoprotein platelet inhibitors	nucleoside reverse transcriptase inhibitors (NRTIs)	thioxanthenes
antiadrenergic agents (peripheral) with thiazides	glycylcyclines	nutraceutical products	third generation cephalosporins
antiadrenergic agents, centrally acting	gonadotropin releasing hormones	nutritional products	thrombin inhibitors
antiadrenergic agents, peripherally acting	gonadotropin-releasing hormone antagonists	ophthalmic anesthetics	thrombolytics
antiandrogens	gonadotropins	ophthalmic anti-infectives	thyroid drugs
antianginal agents	group I antiarrhythmics	ophthalmic anti-inflammatory agents	TNF alpha inhibitors
antiarrhythmic agents	group II antiarrhythmics	ophthalmic antihistamines and decongestants	tocolytic agents
antiasthmatic combinations	group III antiarrhythmics	ophthalmic diagnostic agents	topical acne agents
antibiotics/antineoplastics	group IV antiarrhythmics	ophthalmic glaucoma agents	topical anesthetics
anticholinergic antiemetics	group V antiarrhythmics	ophthalmic lubricants and irrigations	topical anti-rosacea agents
anticholinergic antiparkinson agents	growth hormone receptor blockers	ophthalmic preparations	topical antibiotics
anticholinergic bronchodilators	growth hormones		topical antifungals
anticholinergic chronotropic agents	guanylate cyclase-C agonists		topical antihistamines
anticholinergics/antispasmodics	H. pylori eradication agents		topical antineoplastics
anticoagulant reversal agents	H2 antagonists		topical antipsoriasis
anticoagulants	Hedgehog pathway inhibitors		topical antivirals
anticonvulsants	hematopoietic stem cell mobilizer		topical astringents
antidepressants	heparin antagonists		
antidiabetic agents	heparins		
antidiabetic combinations	HER2 inhibitors		
antidiarrheals			
antidiuretic hormones			

TABLE 2-continued

MEDICAMENTS			
antidotes	herbal products	ophthalmic steroids	topical debriding agents
antiemetic/antivertigo agents	histone deacetylase inhibitors	ophthalmic steroids with anti-infectives	topical depigmenting agents
antifungals	hormones	ophthalmic surgical agents	topical emollients
antigonadotropic agents	hormones/antineoplastics	oral nutritional supplements	topical keratolytics
antigout agents	hydantoin	immunostimulants	topical non-steroidal anti-inflammatories
antihistamines	immunoglobulins	immunosuppressants	topical photochemotherapeutics
antihyperlipidemic agents	immunologic agents	otic anesthetics	topical rubefacient
antihyperlipidemic combinations	immunostimulants	otic anti-infectives	topical steroids
antihypertensive combinations	immunostimulatory agents	otic preparations	topical steroids with anti-infectives
antihyperuricemic agents	impotence agents	otic steroids	triazine
antimalarial agents	in vivo diagnostic biologicals	otic steroids with anti-infectives	anticonvulsants
antimalarial combinations	incretin mimetics	oxazolindione anticonvulsants	tricyclic antidepressants
antimalarial quinolines	inhaled anti-infectives	antibiotics	trifunctional monoclonal antibodies
antimetabolites	inhaled corticosteroids	parathyroid hormone and analogs	ultrasound contrast media
antimigraine agents	inotropic agents	PARP inhibitors	upper respiratory combinations
antineoplastic detoxifying agents	insulin	PCSK9 inhibitors	urea anticonvulsants
antineoplastic interferons	insulin-like growth factor	penicillinase resistant penicillins	urea cycle disorder agents
antineoplastics	integrase strand transfer inhibitor	penicillins	urinary anti-infectives
antiparkinson agents	interferons	peripheral opioid receptor antagonists	urinary antispasmodics
antiplatelet agents	interleukin inhibitors	peripheral opioid receptor mixed agonists/antagonists	urinary pH modifiers
antipseudomonal penicillins	interleukins	peripheral vasodilators	uterotonic agents
antipsoriasis	intravenous nutritional products	peripherally acting antiobesity agents	vaccine combinations
antipsychotics	iodinated contrast media	phenothiazine antiemetics	vaginal anti-infectives
antirheumatics	ionic iodinated contrast media	phenothiazine antipsychotics	vaginal preparations
antiseptic and germicides	iron products	phenothiazine antipsychotics	vasodilators
antithyroid agents	ketolides	phenothiazine antipsychotics	vasopressin antagonists
antitoxins and antivenins	laxatives	phenylpiperazine antidepressants	vasopressors
antituberculosis agents	leprostatics	phosphate binders	VEGF/VEGFR inhibitors
antituberculosis combinations	leukotriene modifiers	plasma expanders	viral vaccines
antitussives	lincomycin derivatives	platelet aggregation inhibitors	viscosupplementation agents
antiviral agents	local injectable anesthetics	platelet-stimulating agents	vitamin and mineral combinations
antiviral boosters	local injectable anesthetics with corticosteroids	polyenes	vitamins
antiviral combinations	loop diuretics	potassium sparing diuretics with thiazides	5-alpha-reductase inhibitors
antiviral interferons	lung surfactants	potassium-sparing diuretics	5-aminosalicylates
anxiolytics, sedatives, and hypnotics	lymphatic staining agents	probiotics	5HT3 receptor antagonists
aromatase inhibitors	lysosomal enzymes	progesterone receptor modulators	chloride channel activators
atypical antipsychotics	macrolide derivatives	progestins	cholesterol absorption inhibitors
azole antifungals	macrolides	prolactin inhibitors	cholinergic muscle stimulants
bacterial vaccines	magnetic resonance imaging contrast media	prostaglandin D2 antagonists	cholinesterase inhibitors
barbiturate anticonvulsants	mast cell stabilizers	protease inhibitors	CNS stimulants
barbiturates	medical gas	protease-activated receptor-1 antagonists	coagulation modifiers
BCR-ABL tyrosine kinase inhibitors	meglitinides	proteasome inhibitors	colony stimulating factors
benzodiazepine anticonvulsants	metabolic agents	proton pump inhibitors	contraceptives
benzodiazepines	metabolic agents	psoralens	corticotropin
beta blockers with calcium channel blockers	methylxanthines	psychotherapeutic agents	coumarins and indandiones
beta blockers with thiazides	mineralocorticoids	psychotherapeutic agents	cox-2 inhibitors
beta-adrenergic blocking agents	minerals and electrolytes	purine nucleosides	
beta-lactamase inhibitors	agents	pyrrolidine anticonvulsants	
bile acid sequestrants	analgesics		
biologicals	antibiotics		
bisphosphonates	anticonvulsants		
bone morphogenetic proteins	antidepressants		
bone resorption inhibitors	antidiabetic agents		
bronchodilator combinations	antiemetics		
bronchodilators	antifungals		
calcimimetics	antihyperlipidemic agents		
calcineurin inhibitors	antihypertensive combinations		
calcitonin	antimalarials		
calcium channel blocking agents	antineoplastics		
carbamate anticonvulsants	antiparkinson agents		
carbapenems			
carbonic anhydrase inhibitors			

TABLE 2-continued

MEDICAMENTS	
antipsychotic agents	quinolones
antituberculosis agents	radiocontrast agents
antivirals	radiologic adjuncts
anxiolytics, sedatives and hypnotics	radiologic agents
bone resorption inhibitors	radiologic
cardiovascular agents	conjugating agents
central nervous system agents	radiopharmaceuticals
coagulation modifiers	recombinant
diagnostic dyes	human
diuretics	erythropoietins
genitourinary tract agents	anticonvulsants
GI agents	cardiac stressing agents
hormones	cardioselective beta blockers
metabolic agents	cardiovascular agents
ophthalmic agents	catecholamines
	CD20 monoclonal antibodies
	CD30 monoclonal antibodies
	CD33 monoclonal antibodies
	CD38 monoclonal antibodies
	CD52 monoclonal antibodies
	central nervous system agents
	cephalosporins
	cephalosporins/beta- lactamase inhibitors
	cerumenolytics
	CFTR combinations
	CFTR potentiators
	chelating agents
	chemokine receptor antagonist

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 10

<210> SEQ ID NO 1

<211> LENGTH: 112

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 1

Arg Val Lys Phe Ser Arg Ser Ala Asp Ala Pro Ala Tyr Lys Gln Gly
1 5 10 15

Gln Asn Gln Leu Tyr Asn Glu Leu Asn Leu Gly Arg Arg Glu Glu Tyr
20 25 30

Asp Val Leu Asp Lys Arg Arg Gly Arg Asp Pro Glu Met Gly Gly Lys
35 40 45

Pro Arg Arg Lys Asn Pro Gln Glu Gly Leu Tyr Asn Glu Leu Gln Lys
50 55 60

Asp Lys Met Ala Glu Ala Tyr Ser Glu Ile Gly Met Lys Gly Glu Arg
65 70 75 80

Arg Arg Gly Lys Gly His Asp Gly Leu Tyr Gln Gly Leu Ser Thr Ala
85 90 95

-continued

Thr Lys Asp Thr Tyr Asp Ala Leu His Met Gln Ala Leu Pro Pro Arg
 100 105 110

<210> SEQ ID NO 2
 <211> LENGTH: 336
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 2

```

agagtgaagt tcagcaggag cgcagacgcc cccgcgtaca agcagggcca gaaccagctc    60
tataacgagc tcaatctagg acgaagagag gagtacgatg ttttggacaa gagacgtggc    120
cgggaccctg agatgggggg aaagccgaga aggaagaacc ctcaggaagg cctgtacaat    180
gaactgcaga aagataagat gccggaggcc tacagtgaga ttgggatgaa aggcgagcgc    240
cggaggggca aggggcaaga tggcctttac cagggctca gtacagccac caaggacacc    300
tacgacgcc ttcacatgca ggcctgccc cctcgc    336
    
```

<210> SEQ ID NO 3
 <211> LENGTH: 726
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 3

```

gacatccaga tgacacagac tacatcctcc ctgtctgcct ctctgggaga cagagtcacc    60
atcagttgca gggcaagtca ggacattagt aaatatttaa attggtatca gcagaaacca    120
gatggaactg ttaaactcct gatctaccat acatcaagat tacactcagg agtccatca    180
aggttcagtg gcagtgggtc tggaacagat tattctctca ccattagcaa cctggagcaa    240
gaagatattg ccacttactt ttgccaacag ggtaatacgc ttccgtacac gttcggaggg    300
gggaccaage tggagatcac aggtggcggt ggctcgggcy gtggtgggtc gggtgccggc    360
ggatctgagg taaaactgca ggagtcagga cctggcctgg tggcgcctc acagagcctg    420
tccgtcacat gactgtctc aggggtctca ttaccgact atggtgtaag ctggattcgc    480
cagcctccac gaaagggctt ggagtggctg ggagtaatat ggggtagtga aaccacatac    540
tataattcag ctctcaaate cagactgacc atcatcaagg acaactccaa gagccaagtt    600
ttcttaaaaa tgaacagtct gcaaactgat gacacagcca tttactactg tgccaaacat    660
tattactacg gtggtagcta tgctatggac tactggggcc aaggaacctc agtcaccgtc    720
tcctca    726
    
```

<210> SEQ ID NO 4
 <211> LENGTH: 242
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 4

```

Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly
1           5           10           15
Asp Arg Val Thr Ile Ser Cys Arg Ala Ser Gln Asp Ile Ser Lys Tyr
20          25          30
Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile
    
```

-continued

35					40					45					
Tyr	His	Thr	Ser	Arg	Leu	His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
50					55					60					
Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu	Gln
65					70					75					80
Glu	Asp	Ile	Ala	Thr	Tyr	Phe	Cys	Gln	Gln	Gly	Asn	Thr	Leu	Pro	Tyr
			85						90					95	
Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Thr	Gly	Gly	Gly	Gly	Ser
		100						105						110	
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Lys	Leu	Gln	Glu
		115					120					125			
Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln	Ser	Leu	Ser	Val	Thr	Cys
		130					135					140			
Thr	Val	Ser	Gly	Val	Ser	Leu	Pro	Asp	Tyr	Gly	Val	Ser	Trp	Ile	Arg
145					150					155					160
Gln	Pro	Pro	Arg	Lys	Gly	Leu	Glu	Trp	Leu	Gly	Val	Ile	Trp	Gly	Ser
				165					170						175
Glu	Thr	Thr	Tyr	Tyr	Asn	Ser	Ala	Leu	Lys	Ser	Arg	Leu	Thr	Ile	Ile
			180						185					190	
Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu	Lys	Met	Asn	Ser	Leu	Gln
		195					200					205			
Thr	Asp	Asp	Thr	Ala	Ile	Tyr	Tyr	Cys	Ala	Lys	His	Tyr	Tyr	Tyr	Gly
		210					215					220			
Gly	Ser	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Ser	Val	Thr	Val
225					230					235					240

Ser Ser

<210> SEQ ID NO 5
 <211> LENGTH: 72
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 5

atctacatct gggcgccctt ggccgggact tgtggggctcc ttctcctgtc actgggtatc 60

accctttact gc 72

<210> SEQ ID NO 6
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 6

Ile Trp Ala Pro Leu Ala Gly Thr Cys Gly Val Leu Leu Leu Ser Leu
1 5 10 15Val Ile Thr Leu Tyr Cys
20

<210> SEQ ID NO 7
 <211> LENGTH: 135
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 7

-continued

```
accacgacgc cagcgccgcg accaccaaca cggcgccca ccatcgcgtc gcagccctg      60
tcctcgccc cagaggcgtg ccggccagcg gcgggggcg cagtgcacac gaggggctg      120
gacttcgctt gtgat                                          135
```

```
<210> SEQ ID NO 8
<211> LENGTH: 47
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

```
<400> SEQUENCE: 8
Thr Thr Thr Pro Ala Pro Arg Pro Pro Thr Pro Ala Pro Thr Ile Ala
1           5           10           15
Ser Gln Pro Leu Ser Leu Arg Pro Glu Ala Cys Arg Pro Ala Ala Gly
          20           25           30
Gly Ala Val His Thr Arg Gly Leu Asp Phe Ala Cys Asp Ile Tyr
          35           40           45
```

```
<210> SEQ ID NO 9
<211> LENGTH: 126
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

```
<400> SEQUENCE: 9
aaacggggca gaaagaaact cctgtatata ttcaacaac catttatgag accagtacaa      60
actactcaag aggaagatgg ctgtagctgc cgatttccag aagaagaaga aggaggatgt      120
gaactg                                          126
```

```
<210> SEQ ID NO 10
<211> LENGTH: 42
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

```
<400> SEQUENCE: 10
Lys Arg Gly Arg Lys Lys Leu Leu Tyr Ile Phe Lys Gln Pro Phe Met
1           5           10           15
Arg Pro Val Gln Thr Thr Gln Glu Glu Asp Gly Cys Ser Cys Arg Phe
          20           25           30
Pro Glu Glu Glu Glu Gly Gly Cys Glu Leu
          35           40
```

What is claimed is:

1. A method for enhancing efficacy of an immunotherapy of a cancer selected from the group consisting of melanoma, lung cancer, and colon cancer in a human or animal patient, wherein the method comprises increasing the relative proportion of a sub-population of bacteria comprising *Akkermansia* in a gut microbiota of the patient by administering a bacterial transplant to the patient, wherein an effective amount of the immunotherapy is administered to the patient, wherein the immunotherapy is selected from the group consisting of an immune checkpoint therapy, an adoptive cell therapy, a CAR-T (chimeric antigen receptor T) cell therapy, and a TIL (Tumor-infiltrating lymphocyte) therapy, wherein the bacterial transplant comprises *Akkermansia*, and

wherein increasing the relative proportion of the sub-population of bacteria modulates immune cells in the patient, whereby the efficacy of the immunotherapy is enhanced for treatment of the cancer in the patient.

2. The method of claim 1, wherein the immunotherapy is an immune checkpoint therapy comprising administration of an immune checkpoint modulator.

3. The method of claim 2, wherein the immune checkpoint modulator is a CTLA4 (Cytotoxic T-Lymphocyte Associated Protein 4) inhibitor.

4. The method of claim 3, wherein the immune checkpoint modulator is an antibody.

5. The method of claim 4, wherein the immune checkpoint modulator is ipilimumab or tremelimumab.

6. The method of claim 1, wherein the bacterial transplant comprises *Akkermansia muciphila*.

7. The method of claim 1, wherein increasing the relative proportion of the sub-population of bacteria modulates immune cells selected from CD8+ cells, tumor infiltrating lymphocytes (TILs), CD4+ cells, Treg cells and memory cells and Th17 cells. 5

8. The method of claim 1, wherein the immune cells comprise CD8+ cells.

9. The method of claim 8, wherein the CD8+ cells are CD8+ cells of the patient. 10

10. The method of claim 1, wherein the immune cells are upregulated or expanded in the patient.

11. The method of claim 1, wherein the immune cells comprise memory cells selected from the group consisting of central memory T-cells (TCM), effector memory T-cells (TEM), stem cell memory cells (TSCM), and memory effector T-cells (Teff). 15

12. The method of claim 1, wherein the immune cells comprise memory cells selected from the group consisting of CD45RO+CD62L+ and CD25+ CD45RA- CD45RO+ CD127+ cells. 20

13. The method of claim 1, wherein the immune cells comprise administered cells via adoptive cell therapy.

14. The method of claim 1, wherein the method further comprises administering a purine nucleoside. 25

15. The method of claim 1, wherein the cancer is a colon cancer.

16. The method of claim 1, wherein the cancer is melanoma or lung cancer. 30

* * * * *